Asymptotic Results for the Two-parameter Poisson-Dirichlet **Distribution**

Shui Feng[∗] Department of Mathematics and Statistics McMaster University Hamilton, Ontario Canada L8S 4K1 shuifeng@mcmaster.ca

Fuqing Gao† School of Mathematics and Statistics Wuhan University Wuhan 430072, China fqgao@whu.edu.cn

Abstract

The two-parameter Poisson-Dirichlet distribution is the law of a sequence of decreasing nonnegative random variables with total sum one. It can be constructed from stable and Gamma subordinators with the two-parameters, α and θ , corresponding to the stable component and Gamma component respectively. The moderate deviation principles are established for the two-parameter Poisson-Dirichlet distribution and the corresponding homozygosity when θ approaches infinity, and the large deviation principle is established for the two-parameter Poisson-Dirichlet distribution when both α and θ approach zero.

Key words: Poisson-Dirichlet distribution, two-parameter Poisson-Dirichlet distribution, GEM representation, homozygosity, large deviations, moderate deviations.

AMS 1991 subject classifications: Primary: 60F10; Secondary: 92D10.

1 Introduction

For α in $(0, 1)$ and $\theta > -\alpha$, let $U_k, k = 1, 2, \dots$, be a sequence of independent random variables such that U_k has $Beta(1 - \alpha, \theta + k\alpha)$ distribution. Set

$$
X_1^{\alpha,\theta} = U_1, \ X_n^{\alpha,\theta} = (1 - U_1) \cdots (1 - U_{n-1}) U_n, \ n \ge 2. \tag{1.1}
$$

[∗]Research supported by the Natural Science and Engineering Research Council of Canada

[†]Research supported by the NSF of China(No.10571139).

Then with probability one

$$
\sum_{k=1}^{\infty} X_k^{\alpha,\theta} = 1,
$$

and the law of $(X_1^{\alpha,\theta})$ $\lambda_1^{\alpha,\theta}, X_2^{\alpha,\theta}, \dots$ is called the two-parameter GEM distribution.

Let $\mathbf{P}(\alpha,\theta) = (P_1(\alpha,\theta), P_2(\alpha,\theta), \cdots)$ denote the descending order statistic of $(X_1^{\alpha,\theta})$ $X_1^{\alpha,\theta}, X_2^{\alpha,\theta}, \cdots$). The law of $\mathbf{P}(\alpha, \theta)$ is called the two-parameter Poisson-Dirichlet distribution and is denoted by $\Pi_{\alpha,\theta}$. The well-known one-parameter Poisson-Dirichlet distribution corresponds to $\alpha = 0$.

For each integer $m \geq 2$, taking a random sample of size m from a population with the twoparameter Poisson-Dirichlet distribution. Given the population proportion $\mathbf{p} = (p_1, p_2, \ldots)$, the probability that all samples are of the same type is given by

$$
H_m(\mathbf{p}) = \sum_{i=1}^{\infty} p_i^m,
$$

which is referred to as the homozygosity of order m .

The main properties of the two-parameter Poisson-Dirichlet distribution are studied in Pitman and Yor [\[17\]](#page-18-0) including relations to subordinators, Markov chains, Brownian motion and Brownian bridges. The detailed calculations of moments and parameter estimations were carried out in Carlton [\[2\]](#page-17-0). In [\[6\]](#page-17-1) and the references therein one can find connections between two-parameter Poisson-Dirichlet distribution and models in physics including mean-field spin glasses, random map models, fragmentation, and returns of a random walk to origin. The two-parameter Poisson-Dirichlet distribution has also been used in macroeconomics and finance([\[1\]](#page-17-2)).

Many properties of the one-parameter Poisson-Dirichlet distribution have generalizations in the two-parameter setting including but not limited to the sampling formula (cf. [\[8\]](#page-17-3), [\[16\]](#page-18-1)), the Markov-Krein identity (cf. $[7]$, $[18]$), and subordinator representation (cf. $[13]$, $[17]$). Recently, a large deviation principle (henceforth, LDP)is established in [\[9\]](#page-17-6) for the two-parameter Poisson-Dirichlet distribution when θ goes to infinity. This is a generalization to the LDP result for the one-parameter Poisson-Dirichlet distribution in [\[3\]](#page-17-7). Our first result here establishes the corresponding moderate deviation principle (henceforth, MDP). This can be viewed as a generalization of the MDP result in [\[11\]](#page-17-8) to the two-parameter setting. The MDP for the homozygosity is also established generalizing corresponding result in [\[11\]](#page-17-8). In order to apply the Campbell's theorem, we turn to a representation of the two-parameter Poisson-Dirichlet distribution obtained in [\[16\]](#page-18-1).

When $\alpha = 0$, the one-parameter Poisson-Dirichlet distribution converges to $\delta_{(1,0,...)}$ as θ goes to zero. The corresponding LDP is established in [\[10\]](#page-17-9) where a structure called "energy ladder" is revealed. Our second main result generalizes this result to the two-parameter Poisson-Dirichlet distribution when both α and θ go to zero. It turns out that the large deviation speed will depend on α if it converges to zero at a slower speed than that of θ .

The current paper is organized as follows. Distributional results are derived in Section 2 using the change of measure formula and the subordinator representation. Section 3 is dedicated to establishing the MDP for $\Pi_{\alpha,\theta}$ when θ goes to infinity. The large θ MDP for the homozygosity is established in Section 4. In Section 5 we prove the LDP for $\Pi_{\alpha,\theta}$ when both α and θ go to zero.

2 Marginal Distributions

In this section, we derive the marginal distributions of the two-parameter Poisson-Dirichlet distribution. The basic tools are the change of measure formula and the subordinator representation. For general concepts and theorems on MDP and LDP, we will refer to [\[5\]](#page-17-10).

¿From now on, the parameter θ will be assumed to be positive and α is in $(0, 1)$. Let $\{\rho_s, s \geq 0\}$ be a subordinator, an increasing process with stationary independent increment, with no drift component. The Laplace transform of ρ_s is then given by

$$
E\left(\exp(-\lambda \rho_s)\right) = \exp\left\{ s \int_0^\infty (e^{-\lambda x} - 1)\Lambda(dx) \right\}, \lambda \ge 0,
$$
\n(2.1)

where Λ is the Lévy measure on $(0, +\infty)$ describing the distribution of the jump sizes. Let $V_1(\rho_s) \ge$ $V_2(\rho_s) \geq \cdots$ denote the jump sizes of $\{\rho_s, s \geq 0\}$ over $[0, s)$ in decreasing order.

If

$$
\Lambda(dx) = c_{\alpha} x^{-(1+\alpha)} dx.
$$

for some $c_{\alpha} > 0$, then the subordinator is called a stable subordinator with index α and is denoted by ${\tau_s, s \ge 0}$. Without loss of generality, we choose $c_{\alpha} = \frac{\alpha}{\Gamma(1-\alpha)}$ in this paper.

The next result is from [\[17\]](#page-18-0).

Proposition 2.1 (Pitman and Yor). Let $\{\sigma_s : s \geq 0\}$ and $\{\gamma_s : s \geq 0\}$ be two independent subordinators with respective Lévy measures $\alpha Cx^{-(\alpha+1)}e^{-x}dx$ and $x^{-1}e^{-x}dx$ for some $C > 0$. Let

$$
\zeta(\alpha,\theta) = \frac{\gamma_{\theta/\alpha}}{C\Gamma(1-\alpha)}.
$$

Then $T = T(\alpha, \theta) = \sigma_{\zeta(\alpha, \theta)}$, and

$$
\left(\frac{V_1(T)}{T}, \frac{V_2(T)}{T}, \dots\right)
$$

are independent with respective laws the $Gamma(\theta, 1)$ distribution and the two-parameter Poisson-Dirichlet distribution $\Pi_{\alpha,\theta}$.

Let $E_{\alpha,\theta}$ denote the expectation with respect to $\Pi_{\alpha,\theta}$. For $n \geq 1$, set

$$
C_{\alpha,\theta} = \frac{\Gamma(\theta + 1)}{\Gamma(\frac{\theta}{\alpha} + 1)},\tag{2.2}
$$

$$
C_{\alpha,\theta,n} = \frac{\Gamma(\theta+1)\Gamma(\frac{\theta}{\alpha}+n)\alpha^{n-1}}{\Gamma(\theta+n\alpha)\Gamma(\frac{\theta}{\alpha}+1)\Gamma(1-\alpha)^n}.
$$
\n(2.3)

The following change of measure formula is obtained in [\[15\]](#page-18-3).

Proposition 2.2 (Perman, Pitman and Yor). For any bounded measurable function f on \mathbb{R}^{∞}_+ ,

$$
E_{\alpha,\theta}(f(P_1,P_2,\ldots)) = C_{\alpha,\theta} E\left(\tau_1^{-\theta} f\left(\frac{V_1(\tau_1)}{\tau_1},\frac{V_2(\tau_1)}{\tau_1},\ldots\right)\right),\tag{2.4}
$$

where the law of

$$
\left(\frac{V_1(\tau_1)}{\tau_1}, \frac{V_2(\tau_1)}{\tau_1}, \cdots\right)
$$

is $\Pi_{\alpha,0}$.

Now we are ready to derive the following distributional results.

Theorem 2.3 For each $\beta > 0$, define

$$
g_{\alpha,\beta}(x) = P(P_1(\alpha,\beta) \le x). \tag{2.5}
$$

Then for any $n \geq 1$, the joint density function of $(P_1(\alpha, \theta), \cdots, P_n(\alpha, \theta))$ is given by

$$
h_{\alpha,\theta,n}(p_1,\dots,p_n) = C_{\alpha,\theta,n} \frac{\left(1 - \sum_{i=1}^n p_i\right)^{\theta+n\alpha-1}}{\prod_{i=1}^n p_i^{1+\alpha}} g_{\alpha,\theta+n\alpha} \left(\frac{p_n}{1 - \sum_{i=1}^n p_i}\right). \tag{2.6}
$$

Proof By Proposition [2.2](#page-2-0) and Perman's formula (cf. [\[14\]](#page-18-4)), for any non-negative product measurable function f and any any $n > 1$, the joint density function of $(\tau_1, \frac{V_1(\tau_1)}{\tau_1})$ $\frac{(\tau_1)}{\tau_1}, \cdots, \frac{V_n(\tau_1)}{\tau_1}$ $\frac{(\tau_1)}{\tau_1}$ is given by

$$
\phi_n(t, p_1, \cdots, p_n) = (c_\alpha)^{n-1} \hat{p}_n^{-1} (p_1 \cdots p_{n-1})^{-(1+\alpha)} t^{-(\theta + (n-1)\alpha)} \phi_1(t\hat{p}_n, p_n/\hat{p}_n).
$$
 (2.7)

where

$$
\hat{p}_n = 1 - p_1 - \dots - p_{n-1},\tag{2.8}
$$

and $\phi_1(t, u)$ satisfies

$$
\phi_1(t, u) = c_{\alpha} t^{-\alpha} u^{-(1+\alpha)} \int_0^{\frac{u}{1-u}\wedge 1} \phi_1(t(1-u), v) dv.
$$
\n(2.9)

Integrating out the t coordinate, it follows from (2.9) that

$$
h_{\alpha,\theta,n}(p_1,\dots,p_n) \qquad (2.10)
$$
\n
$$
= C_{\alpha,\theta}(c_{\alpha})^{n-1}\hat{p}_n^{-1}(p_1\cdots p_{n-1})^{-(1+\alpha)} \int_0^{\infty} t^{-(\theta+(n-1)\alpha)} \phi_1(t\hat{p}_n, p_n/\hat{p}_n) dt
$$
\n
$$
= C_{\alpha,\theta}(c_{\alpha})^{n-1}\hat{p}_n^{\theta+(n-1)\alpha-2}(p_1\cdots p_{n-1})^{-(1+\alpha)} \int_0^{\infty} s^{-(\theta+(n-1)\alpha)} \phi_1(s, p_n/\hat{p}_n) ds
$$
\n
$$
= C_{\alpha,\theta}(c_{\alpha})^n \frac{\hat{p}_n^{\theta+n\alpha-1}}{(p_1\cdots p_{n-1}p_n)^{(1+\alpha)}} \int_0^{\frac{p_n}{\hat{p}_{n+1}}\wedge 1} dx \int_0^{\infty} s^{-(\theta+n\alpha)} \phi_1(s(1-p_n/\hat{p}_n), x) ds
$$
\n
$$
= C_{\alpha,\theta}(c_{\alpha})^n \frac{(\hat{p}_{n+1})^{\theta+n\alpha-1}}{(p_1\cdots p_{n-1}p_n)^{(1+\alpha)}} \int_0^{\frac{p_n}{\hat{p}_{n+1}}\wedge 1} dx \int_0^{\infty} u^{-(\theta+n\alpha)} \phi_1(u,x) du
$$
\n
$$
= \frac{C_{\alpha,\theta}(c_{\alpha})^n}{C_{\alpha,\theta+n\alpha}} \frac{(\hat{p}_{n+1})^{\theta+n\alpha-1}}{(p_1\cdots p_{n-1}p_n)^{(1+\alpha)}} g_{\alpha,\theta+n\alpha} \left(\frac{p_n}{1-\sum_{i=1}^n p_i}\right),
$$
\n(2.10)

which leads to [\(2.6\)](#page-3-1).

Remark. This result appears in Handa [\[12\]](#page-17-11) where a different proof is used.

Theorem 2.4 For any $s > 0$,

$$
\nu_{\alpha,\theta,1}(s) = P(V_1(T) \le s)
$$
\n
$$
= \left(1 + c_{\alpha}s^{-\alpha} \int_1^{\infty} z^{-(1+\alpha)} e^{-sz} dz\right)^{-\theta/\alpha}.
$$
\n(2.11)

Proof For each $s > 0$, it follows from Proposition [2.1](#page-2-1) and the property of the Poisson random measure that

$$
\nu_{\alpha,\theta,1}(s) = E(P(V_1(T) \le s | \zeta(\alpha, \theta)))
$$

=
$$
E\left(\exp\left\{-\alpha C\zeta(\alpha, \theta)\int_s^{\infty} x^{-(\alpha+1)}e^{-x}dx\right\}\right)
$$

=
$$
E\left(\exp\left\{-c_{\alpha}\gamma_{\theta/\alpha}s^{-\alpha}\int_1^{\infty} z^{-(\alpha+1)}e^{-sz}dz\right\}\right)
$$

which leads to (2.11) .

3 Moderate Deviations for the two-parameter Poisson-Dirichlet Distribution

By theorem 6.1 in [\[12\]](#page-17-11), when θ goes to infinity $\mathbf{P}(\alpha, \theta) = (P_1(\alpha, \theta), P_2(\alpha, \theta), \cdots)$ approaches a non-trivial random sequence when scaled by a factor of θ and shifted by

$$
\beta(\alpha, \theta) = \log \theta - (\alpha + 1) \log \log \theta - \log \Gamma(1 - \alpha).
$$

In [\[9\]](#page-17-6), the LDP has been established associated with the limit

$$
\lim_{\theta \to \infty} \mathbf{P}(\alpha, \theta) = (0, 0, \ldots).
$$

Replacing the scaling factor by $a(\theta)$ satisfying

$$
\lim_{\theta \to \infty} \frac{a(\theta)}{\theta} = 0, \quad \lim_{\theta \to \infty} a(\theta) = \infty,
$$
\n(3.1)

we still have

 \Box

 \Box

$$
\lim_{\theta \to \infty} a(\theta) \left(\mathbf{P}(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta} (1, 1, \ldots) \right) \to (0, 0, \ldots). \tag{3.2}
$$

The LDP associated with [\(3.2\)](#page-5-0) is called the MDP for $\mathbf{P}(\alpha, \theta) = (P_1(\alpha, \theta), P_2(\alpha, \theta) \cdots)$. This MDP will be established in this section through a series of lemmas.

The first lemma establishes the MDP for $V_1(T)/\theta$.

Lemma 3.1 The MDP holds for $V_1(T)/\theta$ with speed $\frac{a(\theta)}{\theta}$ and rate function

$$
J_1(x) = \begin{cases} x, & x \ge 0 \\ \infty, & otherwise. \end{cases}
$$

Proof For any fixed x , we have

$$
P\left\{a(\theta)\left(\frac{V_1(T) - \beta(\alpha, \theta)}{\theta}\right) \le x\right\} = P\left(V_1(T) < \frac{\theta}{a(\theta)}x + \beta(\alpha, \theta)\right). \tag{3.3}
$$

Assume that

$$
\lim_{\theta \to \infty} \left[\frac{\theta}{a(\theta)} x + \beta(\alpha, \theta) \right] = +\infty.
$$

Then it follows from [\(2.11\)](#page-4-0) that

$$
P\left(V_1(T) < \frac{\theta}{a(\theta)}x + \beta(\alpha, \theta)\right) \sim \left(1 + \frac{c_\alpha}{(\frac{\theta}{a(\theta)}x + \beta(\alpha, \theta))^{\alpha+1}}e^{-(\frac{\theta}{a(\theta)}x + \beta(\alpha, \theta))}\right)^{-\theta/\alpha}
$$

Therefore

$$
\limsup_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left(a(\theta) \left(\frac{V_1(T) - \beta(\alpha, \theta)}{\theta} \right) \leq x \right)
$$
\n
$$
= \lim_{\theta \to \infty} \frac{a(\theta)}{\theta} \log \left(1 + \frac{c_{\alpha} (\log \theta)^{\alpha+1} \Gamma(1-\alpha)}{\theta(\frac{\theta}{a(\theta)} x + \beta(\alpha, \theta))^{\alpha+1}} e^{-\frac{\theta}{a(\theta)} x} \right)^{-\theta/\alpha}
$$
\n
$$
= \begin{cases} 0, & x \geq 0 \\ -\infty, & x < 0 \end{cases} (3.4)
$$

If there exists a subsequence θ' such that the $\lim_{\theta'\to\infty}$ $\left(\frac{\theta'}{a(\theta)}\right)$ $\frac{\theta'}{a(\theta')}x+\beta(\alpha,\theta')$ exists in $[-\infty,+\infty)$, then x must be strictly negative. Since, by Theorem [2.4,](#page-4-1) $V_1(\hat{T})$ converges to infinity as θ converges to infinity, it follows that

$$
\limsup_{\theta' \to \infty} \frac{a(\theta')}{\theta'} \log P\left(a(\theta')\left(\frac{V_1(T) - \beta(\alpha, \theta')}{\theta'}\right) \le x\right) = -\infty.
$$
\n(3.5)

Putting [\(3.4\)](#page-5-1) and [\(3.5\)](#page-5-2) together, one gets

$$
\lim_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P\left(a(\theta) \left(\frac{V_1(T) - \beta(\alpha, \theta)}{\theta}\right) \le x\right) = 0, \ x \ge 0,
$$
\n(3.6)

and

$$
\limsup_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P\left(a(\theta) \left(\frac{V_1(T) - \beta(\alpha, \theta)}{\theta}\right) \le x\right) = -\infty, \ x < 0. \tag{3.7}
$$

For $x \geq 0$, it follows from (3.3) and (2.11) that

$$
\limsup_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left(a(\theta) \left(\frac{V_1(T) - \beta(\alpha, \theta)}{\theta} \right) \geq x \right)
$$
\n
$$
= \limsup_{\theta \to \infty} \frac{a(\theta)}{\theta} \log \left(1 - \left(1 + \frac{c_{\alpha} e^{-\left(\frac{\theta}{a(\theta)} x + \beta(\alpha, \theta) \right)}}{\theta \left(\frac{\theta}{a(\theta)} x + \beta(\alpha, \theta) \right)^{\alpha + 1}} \right)^{-\theta/\alpha} \right)
$$
\n
$$
= \limsup_{\theta \to \infty} \frac{a(\theta)}{\theta} \log \left(1 + \frac{c_{\alpha} e^{-\left(\frac{\theta}{a(\theta)} x + \beta(\alpha, \theta) \right)}}{\theta \left(\frac{\theta}{a(\theta)} x + \beta(\alpha, \theta) \right)^{\alpha + 1}} \right)^{-\theta/\alpha} = -x.
$$
\n(3.8)

A combination of [\(3.8\)](#page-6-0) and [\(3.7\)](#page-6-1) implies that the laws of $a(\theta) \left(\frac{V_1(T) - \beta(\alpha, \theta)}{\theta} \right)$ is exponentially tight.

Similarly, we can get that for $x > 0$ and $\delta > 0$ with $x - \delta > 0$,

$$
\lim_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left(a(\theta) \left(\frac{V_1(T) - \beta(\alpha, \theta)}{\theta} \right) \in (x - \delta, x + \delta) \right)
$$
\n
$$
= \lim_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left(a(\theta) \left(\frac{V_1(T) - \beta(\alpha, \theta)}{\theta} \right) \in (x - \delta, x + \delta) \right)
$$
\n
$$
= -x + \delta. \tag{3.9}
$$

The equality [\(3.6\)](#page-6-2) combined with [\(3.7\)](#page-6-1) implies that

$$
\lim_{\delta \to 0} \limsup_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left(a(\theta) \left(\frac{V_1(T) - \beta(\alpha, \theta)}{\theta} \right) \in (-\delta, \delta) \right)
$$
\n
$$
= \lim_{\delta \to 0} \liminf_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left(a(\theta) \left(\frac{V_1(T) - \beta(\alpha, \theta)}{\theta} \right) \in (-\delta, \delta) \right)
$$
\n
$$
= 0. \tag{3.10}
$$

The lemma now follows from [\(3.9\)](#page-6-3), [\(3.10\)](#page-6-4), and the exponential tightness.

 \Box

Set

$$
\gamma(\theta) = \frac{a(\theta)\beta(\alpha,\theta)}{\theta},
$$

and, without loss of generality, we can assume that

$$
\lim_{\theta \to \infty} \gamma(\theta) = c \in [0, +\infty].
$$

It is clear that

$$
\frac{a(\theta)}{\gamma^2(\theta)} = \frac{\theta^2}{a(\theta)\beta^2(\alpha,\theta)} \to \infty, \quad \theta \to \infty.
$$
 (3.11)

If $c < \infty$, it follows from Corollary 3.1 in [\[11\]](#page-17-8) that for any $L > 0$

$$
\limsup_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left\{ \gamma(\theta) \Big| \frac{\theta}{T} - 1 \Big| \ge L \right\} = -\infty. \tag{3.12}
$$

For $c = \infty$, and any $1 > \delta > 0$

$$
\left\{\gamma(\theta)|\frac{\theta}{T} - 1| \ge L\right\} \subset \left\{\gamma(\theta)|\frac{T}{\theta} - 1| \ge L(1 - \delta)\right\} \bigcup \left\{\left|\frac{T}{\theta} - 1\right| \ge \delta\right\}.
$$
\n(3.13)

Since $\gamma(\theta) \leq \beta(\alpha, \theta)$ for large θ and $\lim_{\theta \to \infty} \frac{\beta(\alpha, \theta)}{\sqrt{\theta}} = 0$, it follows from the MDP (Theorem 3.2) in [\[11\]](#page-17-8)) for T/θ , and [\(3.11\)](#page-7-0) that

$$
\limsup_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left\{ \gamma(\theta) | \frac{T}{\theta} - 1 | \ge (1 - \delta)L \right\}
$$
\n
$$
= \limsup_{\theta \to \infty} \frac{a(\theta)}{\gamma^2(\theta)} \frac{\gamma^2(\theta)}{\theta} \log P \left\{ \gamma(\theta) | \frac{T}{\theta} - 1 | \ge (1 - \delta)L \right\} = -\infty,
$$
\n(3.14)

which combined with Corollary 3.1 in [\[11\]](#page-17-8) and (3.13) shows that (3.12) still holds in this case. Therefore $a(\theta)(P_1(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta})$ $\frac{\partial(\alpha,\theta)}{\partial(\theta)}$ and $\frac{\theta}{T}a(\theta)(\frac{V_1(T)-\beta(\alpha,\theta)}{\theta})$ are exponentially equivalent.

Since $\frac{\theta}{T}a(\theta)(\frac{V_1(T)-\beta(\alpha,\theta)}{\theta})$ is exponentially equivalent to $a(\theta)(\frac{V_1(T)-\beta(\alpha,\theta)}{\theta})$ by Lemma 2.1 and Corollary 3.1 in [\[11\]](#page-17-8), it follows that $a(\theta)(P_1(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta})$ $\frac{\partial(\alpha,\theta)}{\partial(\theta)}$ and $a(\theta)$ ($\frac{V_1(T)-\beta(\alpha,\theta)}{\theta}$) are exponentially equivalent. Thus we have the following result.

Lemma 3.2 The MDP holds for $P_1(\alpha, \theta)$ with speed $\frac{a(\theta)}{\theta}$ and rate function

$$
J_1(x) = \begin{cases} x, & x \ge 0 \\ \infty, & otherwise. \end{cases}
$$

For each $n \geq 2$, we have

Lemma 3.3 The family $\Big\{ P\Big(a(\theta)\Big(P_1(\alpha,\theta)-\frac{\beta(\alpha,\theta)}{\theta}$ $\frac{\partial \alpha_i(\theta)}{\partial}$ \cdots , $P_n(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta}$ $\left(\frac{\alpha,\theta)}{\theta}\right) \in \cdot\bigg) : \theta > 0 \Big\}$ satisfies a LDP on \mathbb{R}^n with speed $\frac{a(\theta)}{\theta}$ and rate function

$$
I_n(x_1, \dots, x_n) = \begin{cases} \sum_{i=1}^n x_i, & \text{if } 0 \le x_n \le \dots \le x_1. \\ +\infty, & \text{otherwise.} \end{cases} \tag{3.15}
$$

Proof It follows from [\(3.3\)](#page-5-3) that for $x_1 \ge x_2 \cdots \ge x_n$ and $\frac{\theta}{a(\theta)} x_n + \beta(\alpha, \theta) > 0$, the density function $g_{\alpha,\theta,n}(x_1,\ldots,x_n)$ of $a(\theta)\Big(P_1(\alpha,\theta)-\frac{\beta(\alpha,\theta)}{\theta}$ $\frac{\partial \alpha}{\partial \theta} \cdots$, $P_n(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta}$ $\frac{\alpha,\theta)}{\theta}$) is

$$
g_{\alpha,\theta,n}(x_1,\ldots,x_n) = \left(\frac{1}{a(\theta)}\right)^n C_{n,\alpha,\theta} \left(\prod_{i=1}^n \left(\frac{\theta}{\frac{\theta}{a(\theta)}x_i + \beta(\alpha,\theta)}\right)^{\alpha+1}\right) \times \left(1 - \left(\frac{\theta}{a(\theta)}\sum_{i=1}^n x_i + n\beta(\alpha,\theta)\right)/\theta\right)^{\theta+n\alpha-1} g_{\alpha,\theta+n\alpha} \left(\frac{\frac{\theta}{a(\theta)}x_n + \beta(\alpha,\theta)}{\theta - \left(\frac{\theta}{a(\theta)}\sum_{i=1}^n x_i + n\beta(\alpha,\theta)\right)}\right).
$$
\n(3.16)

By Theorem [2.4](#page-4-1) and direct calculation, for $x_n > 0$

$$
\frac{a(\theta)}{\theta} \log g_{\alpha,\theta+n\alpha} \left(\frac{\frac{\theta}{a(\theta)} x_n + \beta(\alpha,\theta)}{\theta - \left(\frac{\theta}{a(\theta)} \sum_{i=1}^n x_i + n\beta(\alpha,\theta)\right)} \right) \to 0.
$$

For $x_n < 0$, set

$$
\psi(n, x, \theta, \alpha) = a(\theta) \left(\frac{\frac{\theta}{a(\theta)} x_n + \beta(\alpha, \theta)}{\theta - \left(\frac{\theta}{a(\theta)} \sum_{i=1}^n x_i + n\beta(\alpha, \theta) \right)} - \frac{\beta(\alpha, \theta + n\alpha)}{\theta + n\alpha} \right).
$$

Then

$$
g_{\alpha,\theta+n\alpha} \left(\frac{\frac{\theta}{a(\theta)} x_n + \beta(\alpha, \theta)}{\theta - \left(\frac{\theta}{a(\theta)} \sum_{i=1}^n x_i + n\beta(\alpha, \theta)\right)} \right)
$$

= $P\left(a(\theta) \left(P_1(\alpha, \theta + n\alpha) - \frac{\beta(\alpha, \theta + n\alpha)}{\theta + n\alpha} \right) < \psi(n, x, \theta, \alpha) \right)$

and

$$
\lim_{\theta \to \infty} \psi(n, x, \theta, \alpha) = x_n < 0
$$

which implies that

$$
\lim_{\theta \to \infty} \frac{a(\theta)}{\theta} \log g_{\alpha,\theta+n\alpha} \left(\frac{\frac{\theta}{a(\theta)} x_n + \beta(\alpha,\theta)}{\theta - \left(\frac{\theta}{a(\theta)} \sum_{i=1}^n x_i + n\beta(\alpha,\theta)\right)} \right) = -\infty.
$$

Therefore

$$
\frac{a(\theta)}{\theta} \log g_{\alpha,\theta,n}(x_1,\ldots,x_n) \to -\sum_{i=1}^n x_i, \ x_n > 0,
$$
\n(3.17)

$$
\frac{a(\theta)}{\theta} \log g_{\alpha,\theta,n}(x_1,\ldots,x_n) \to -\infty, \ x_n < 0,\tag{3.18}
$$

For $x_1 \ge x_2 \cdots \ge x_n$, let $B((x_1, \ldots, x_n), \delta)$ denote the closed ball centered at (x_1, \ldots, x_n) with radius δ , and $B^{\circ}((x_1,\ldots,x_n),\delta)$ be the corresponding open ball. Then for $x_n > 0$,

$$
\lim_{\delta \to 0} \limsup_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left(a(\theta) \Big(P_1(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta} \cdots, P_n(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta} \Big) \in B((x_1, \dots, x_n), \delta) \right)
$$
\n
$$
= \lim_{\delta \to 0} \liminf_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left(a(\theta) \Big(P_1(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta} \cdots, P_n(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta} \Big) \in B^{\circ}((x_1, \dots, x_n), \delta) \right)
$$
\n
$$
= - \sum_{i=1}^n x_i,
$$

and for any $x_n < 0$,

$$
\lim_{\delta \to 0} \limsup_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left(a(\theta) \Big(P_1(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta} \cdots, P_n(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta} \Big) \in B((x_1, \dots, x_n), \delta) \right)
$$
\n
$$
= \lim_{\delta \to 0} \liminf_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left(a(\theta) \Big(P_1(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta} \cdots, P_n(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta} \Big) \in B^{\circ}((x_1, \dots, x_n), \delta) \right)
$$
\n
$$
= -\infty,
$$

If $x_{r-1} > 0, x_r = 0$ for some $1 \leq r \leq n$, then the upper estimate is obtained from that of $a(\theta)\Big(P_1(\alpha,\theta)-\frac{\beta(\alpha,\theta)}{\theta}$ $\frac{\partial \alpha}{\partial \theta} \cdots$, $P_{r-1}(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta}$ $\left(\frac{\alpha,\theta}{\theta}\right)$. The lower estimates when $x_r = 0$ for some $1 \le r \le n$ are obtained by approximating the boundary with open subsets away from the boundary.

Noting that $\bigcup_{i=1}^n \{a(\theta)\left(P_i(\alpha,\theta) - \frac{\beta(\alpha,\theta)}{\theta}\right)$ $\left(\frac{\alpha,\theta)}{\theta}\right)$ > L} = { $a(\theta)\left(P_1(\alpha,\theta)-\frac{\beta(\alpha,\theta)}{\theta}\right)$ $\left(\frac{\alpha,\theta)}{\theta}\right) > L$, it follows that

$$
\lim_{L \to \infty} \limsup_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left\{ \bigcup_{i=1}^{n} \left\{ a(\theta) \left(P_i(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta} \right) > L \right\} \right\} = -\infty. \tag{3.19}
$$

On the other hand,

$$
\limsup_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left\{ \bigcup_{i=1}^{n} \left\{ a(\theta) \left(P_i(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta} \right) < -L \right\} \right\} \tag{3.20}
$$
\n
$$
\leq \limsup_{\theta \to \infty} \frac{a(\theta)}{\theta} \log P \left\{ a(\theta) \left(P_i(\alpha, \theta) - \frac{\beta(\alpha, \theta)}{\theta} \right) \leq -L \right\} = -\infty.
$$

These lead to the exponential tightness and the lemma.

Now we are ready to establish the MDP for $(P_1(\alpha, \theta), P_2(\alpha, \theta), \ldots)$.

Theorem 3.4 For each $n \geq 1$, the family $\left\{ P\left(a(\theta)\left(P_1(\theta) - \frac{\beta(\theta)}{\theta}\right)\right) \right\}$ $\frac{\theta}{\theta}$, \cdots , $P_n(\theta) - \frac{\beta(\theta)}{\theta}$ $\left(\frac{\theta}{\theta},\cdots\right)\in \cdot\bigg)\,:\,\, \theta>$ $\{0\}$ satisfies a LDP on \mathbb{R}^{∞} with speed $\frac{a(\theta)}{\theta}$ and rate function

$$
I(x_1, x_2, \dots) = \begin{cases} \sum_{i=1}^{\infty} x_i, & x_1 \geq \dots \geq 0 \\ \infty, & otherwise. \end{cases}
$$
 (3.21)

Proof Identify \mathbb{R}^{∞} with the projective limit of \mathbb{R}^{n} , $n = 1, \ldots$. Then the theorem follows from Theorem 3.3 in [\[4\]](#page-17-12) and Lemma [3.3.](#page-7-3)

 \Box

 \Box

4 Moderate Deviations for the Homozygosity

For each $m \geq 2$, it was shown in [\[12\]](#page-17-11) that the scaled homozygosity

$$
\sqrt{\theta} \left[\frac{\theta^{m-1} \Gamma(1-\alpha)}{\Gamma(m-\alpha)} H_m(\mathbf{P}(\alpha,\theta)) - 1 \right] \Rightarrow Z_{\alpha,m}
$$

where $Z_{\alpha,m}$ is a normal random variable with mean zero and variance

$$
\sigma_{\alpha,m}^2 = \frac{\Gamma(2m-\alpha)\Gamma(1-\alpha)}{\Gamma(m-\alpha)^2} + \alpha - m^2.
$$

It is thus natural to consider the MDP for $\frac{\theta^{m-1}}{\Gamma(m)}H_m(\mathbf{P}(\alpha,\theta))$ or equivalently the LDP for the family $\{a(\theta) \left[\frac{\theta^{m-1} \Gamma(1-\alpha)}{\Gamma(m-\alpha)} H_m(\mathbf{P}(\alpha, \theta)) - 1 \right] : \theta > 0 \}$ for a scale $a(\theta)$ satisfying

$$
\lim_{\theta \to \infty} a(\theta) = \infty, \lim_{\theta \to \infty} \frac{a(\theta)}{\sqrt{\theta}} = 0,
$$
\n(4.1)

which is different from (3.1) .

The MDP in the case of $\alpha = 0$ has been established in [\[11\]](#page-17-8) where that the following additional restriction on $a(\theta)$ is used: for some $0 < \epsilon < 1/(2m-1)$,

$$
\liminf_{\theta \to \infty} \frac{a^{1-\varepsilon}(\theta)}{\theta^{(m-1)/(2m-1)}} > 0. \tag{4.2}
$$

This condition is also needed for the two-parameter model. As shown in [\[11\]](#page-17-8), the conditions [\(4.1\)](#page-10-0) and [\(4.2\)](#page-10-1) guarantee that there exist $\tau > 0$ positive integer $l \geq 3 \vee \frac{2}{(2m-1)!}$ $\frac{2}{(2m-1)\varepsilon}$, and $r(\theta)$ that grows faster than a positive power of θ such that

$$
\lim_{\theta \to \infty} \frac{a(\theta)}{\theta^{\tau}} = +\infty
$$

and

$$
\lim_{\theta \to \infty} \frac{r(\theta)^{m-1}}{a^{(l-2)/l}(\theta)} = 0, \quad \lim_{\theta \to \infty} \frac{a^2(\theta)r(\theta)}{\theta} = \infty.
$$

For any $n \geq 1$, set

$$
G_{\alpha,\theta,r}^{(n)} = \sum_{i=1}^{\infty} V_i^n(T) I_{\{V_i(T) \le r(\theta)\}},
$$

$$
G_{\alpha,\theta}^{(n)} = \sum_{i=1}^{\infty} V_i^n(T),
$$

and

$$
G_{\alpha,\theta,r} = \left(G_{\alpha,\theta,r}^{(1)} - E(G_{\alpha,\theta,r}^{(1)}), G_{\alpha,\theta,r}^{(m)} - E(G_{\alpha,\theta}^{(m)}) \right).
$$

For any s, t in \mathbb{R} , define

$$
\Lambda(s,t)=\frac{1}{2}\left(s^2+\frac{2\Gamma(m-\alpha)\Gamma(m+1)}{\Gamma(m)\Gamma(1-\alpha)}st+\big(\frac{\Gamma(2m-\alpha)}{\Gamma(1-\alpha)}+\alpha\big(\frac{\Gamma(m-\alpha)}{\Gamma(1-\alpha)}\big)^2\big)t^2\right).
$$

It follows by direct calculation that the Fenchel-Legendre transform of $\Lambda(s,t)$ is given by

$$
\Lambda^*(x, y) = \sup_{s,t} \{sx + ty - \Lambda(s, t)\}
$$

=
$$
\frac{\Gamma(1 - \alpha)}{2(\Gamma(1 - \alpha)\Gamma(2m - \alpha) + (\alpha - m^2)\Gamma^2(m - \alpha))}
$$

$$
\times \left((\Gamma(2m - \alpha) + \alpha \frac{\Gamma^2(m - \alpha)}{\Gamma(1 - \alpha)})x^2 - 2m\Gamma(m - \alpha)xy + \Gamma(1 - \alpha)y^2 \right),
$$
 (4.3)

for x, y in \mathbb{R} .

Lemma 4.1 The family $\{\frac{a(\theta)}{\theta}G_{\alpha,\theta,r}:\theta>0\}$ satisfies a LDP on space \mathbb{R}^2 with speed $\frac{a^2(\theta)}{\theta}$ $\frac{\partial}{\partial \theta}$ and rate function $\Lambda^*(\cdot,\cdot)$.

Proof For any $s, t \in \mathbb{R}$, let

$$
g(x) = sx + tx^m
$$

and

$$
\varphi_r(x) = \frac{g(x)I_{\{x \le r(\theta)\}}}{a(\theta)}.
$$

It follows by direct calculation that

$$
\int_{0}^{r(\theta)} (e^{\varphi_r(x)} - 1)x^{-(1+\alpha)}e^{-x}dx \n= \int_{0}^{r(\theta)} \frac{g(x)}{a(\theta)}x^{-(1+\alpha)}e^{-x}dx + \frac{1}{2}\int_{0}^{r(\theta)} \frac{g^2(x)}{a^2(\theta)}x^{-(1+\alpha)}e^{-x}dx \n+ \sum_{k=3}^{l} \frac{1}{k!} \frac{1}{a^k(\theta)} \int_{0}^{\gamma(\theta)} |sx + tx^m|^k x^{-(1+\alpha)}e^{-x}dx \n+ O\left(\sum_{k=l+1}^{\infty} \frac{1}{k!} \frac{1}{a^k(\theta)} (|s| + |t|\gamma(\theta)^{m-1})^k \Gamma(k - \alpha)\right) \n= \int_{0}^{r(\theta)} \frac{g(x)}{a(\theta)}x^{-(1+\alpha)}e^{-x}dx + \frac{1}{2}\int_{0}^{r(\theta)} \frac{g^2(x)}{a^2(\theta)}x^{-(1+\alpha)}e^{-x}dx + o\left(\frac{1}{a^2(\theta)}\right),
$$
\n(4.4)

which implies that for θ large enough,

$$
|\int_0^{r(\theta)} (e^{\varphi_r(x)} - 1)x^{-(1+\alpha)}e^{-x} dx| < c_\alpha^{-1}.
$$

By the Campbell's theorem we get that

$$
E\left(\exp\left\{\frac{1}{a(\theta)}(sG_{\alpha,\theta,r}^{(1)} + tG_{\alpha,\theta,r}^{(m)})\right\}\right)
$$

\n
$$
= E\left(\exp\left\{\sum_{i=1}^{\infty} \varphi_r(V_i(T))\right\}\right)
$$

\n
$$
= E\left(E\left(\exp\left\{\sum_{i=1}^{\infty} \varphi_r(V_i(T))\right\}|\zeta(\alpha,\theta)\right)\right)
$$

\n
$$
= E\left(\exp\left\{c_{\alpha}\gamma\left(\frac{\theta}{\alpha}\right)\int_0^{r(\theta)} (e^{\varphi_r(x)} - 1)x^{-(1+\alpha)}e^{-x}dx\right\}\right)
$$

\n
$$
= \exp\left\{-\frac{\theta}{\alpha}\log\left(1 - c_{\alpha}\int_0^{r(\theta)} (e^{\varphi_r(x)} - 1)x^{-(1+\alpha)}e^{-x}dx\right)\right\}.
$$
\n(4.5)

Putting [\(4.4\)](#page-12-0) and [\(4.5\)](#page-12-1) together, we get that

$$
E\left(\exp\left\{\frac{1}{a(\theta)}(s(G_{\alpha,\theta,r}^{(1)}-E(G_{\alpha,\theta,r}^{(1)}))+t(G_{\alpha,\theta,r}^{(m)}-E(G_{\alpha,\theta,r}^{(m)})))\right\}\right)
$$

= $\exp\left\{\frac{\theta c_{\alpha}}{2\alpha a^2(\theta)}\left(c_{\alpha}\left(\int_0^{\infty}g(x)x^{-(1+\alpha)}e^{-x}dx\right)^2+\int_0^{\infty}g^2(x)x^{-(1+\alpha)}e^{-x}dx+o(\frac{1}{a^2(\theta)})\right)\right\}$
= $\exp\left(\frac{\theta}{a^2(\theta)}(\Lambda(s,t)+o(\frac{1}{a^2(\theta)})))\right),$

which leads to

$$
\lim_{\theta \to \infty} \frac{a^2(\theta)}{\theta} \log E \left(\exp \left\{ \frac{1}{a(\theta)} [s(G_{\alpha,\theta,r}^{(1)} - E(G_{\alpha,\theta,r}^{(1)})) + t(G_{\alpha,\theta,r}^{(m)} - E(G_{\alpha,\theta,r}^{(m)}))] \right\} \right) = \Lambda(s,t). \tag{4.6}
$$

The lemma now follows from (4.3) and the Gärtner-Ellis theorem.

Lemma 4.2 Set

$$
G_{\alpha,\theta} = \left(T - \theta, G_{\alpha,\theta}^{(m)} - E(G_{\alpha,\theta}^{(m)})\right).
$$

Then the family $\{\frac{a(\theta)}{\theta}G_{\alpha,\theta} : \theta > 0\}$ satisfies a LDP with speed $\frac{a^2(\theta)}{\theta}$ $\frac{\partial^{(0)}}{\partial^{(0)}}$ and the rate function $\Lambda^*(x, y)$.

Proof By definition for any $n \ge 1$ and any $\delta > 0$,

$$
\limsup_{\theta \to \infty} \frac{a^2(\theta)}{\theta} \log P \left(\left| G_{\alpha,\theta,r}^{(m)} - G_{\alpha,\theta}^{(m)} \right| \ge \delta \frac{\theta}{a(\theta)} \right)
$$
\n
$$
\leq \limsup_{\theta \to \infty} \frac{a^2(\theta)}{\theta} \log P \left(V_1(T) \ge r(\theta) \right)
$$
\n
$$
= \limsup_{\theta \to \infty} \frac{a^2(\theta)}{\theta} \log \left(1 - \left(1 + \frac{c_{\alpha}}{r^{\alpha}(\theta)} \int_1^{\infty} z^{-(1+\alpha)} e^{-r(\theta)z} dz \right)^{-\theta/\alpha} \right)
$$
\n
$$
\leq \limsup_{\theta \to \infty} \frac{a^2(\theta)}{\theta} \log \left(1 - \left(1 + \frac{c_{\alpha}}{r^{(1+\alpha)}(\theta) e^{r(\theta)}} \right)^{-\theta/\alpha} \right)
$$
\n
$$
= \limsup_{\theta \to \infty} \frac{a^2(\theta)}{\theta} \log \left(\frac{\theta}{\alpha} \log \left(1 + \frac{c_{\alpha}}{r^{(1+\alpha)}(\theta) e^{r(\theta)}} \right) \right)
$$
\n
$$
\leq - \limsup_{\theta \to \infty} \frac{a^2(\theta) r(\theta)}{\theta} \left(1 - \frac{\log \theta}{r(\theta)} \right)
$$
\n
$$
= -\infty.
$$

which implies that $\frac{a(\theta)}{\theta}G_{\alpha,\theta,r}$ and $\frac{a(\theta)}{\theta}G_{\alpha,\theta}$ are exponentially equivalent. Therefore $\left(\frac{a(\theta)}{\theta}G_{\alpha,\theta},\frac{a^2(\theta)}{\theta}\right)$ $\left(\frac{\theta)}{\theta},\Lambda^*\right)$ satisfies LDP.

 \Box

 \Box

Now we are ready to prove the main result of this section.

Theorem 4.3 The family $a(\theta) \left(\frac{\theta^{m-1} \Gamma(1-\alpha)}{\Gamma(m-\alpha)} H_m(\mathbf{P}(\alpha,\theta)) - 1 \right)$ satisfies a LDP with speed $\frac{a^2(\theta)}{\theta}$ $\frac{\overline{v}}{\theta}$ and rate function $\frac{z^2}{2\sigma^2}$ $\frac{z^2}{2\sigma_{\alpha,m}^2}$.

Proof By direct calculation,

$$
a(\theta) \left(\frac{\theta^{m-1} \Gamma(1-\alpha)}{\Gamma(m-\alpha)} H_m(\mathbf{P}(\alpha, \theta)) - 1 \right)
$$

= $a(\theta) \left(\frac{\theta^{m-1} G_{\alpha, \theta}^{(m)}}{T^m \Gamma(m-\alpha)/\Gamma(1-\alpha)} - 1 \right)$
= $a(\theta) \left(\left(\frac{\theta}{T} \right)^m - 1 \right) + \left(\frac{\theta}{T} \right)^m \frac{a(\theta) (G_{\alpha, \theta}^{(m)} - E(G_{\alpha, \theta}^{(m)})}{\theta \Gamma(m-\alpha)/\Gamma(1-\alpha)} - \frac{a(\theta)}{\theta} (\theta - T) \sum_{k=1}^m \left(\frac{\theta}{T} \right)^k + \left(\frac{\theta}{T} \right)^m \frac{a(\theta) (G_{\alpha, \theta}^{(m)} - E(G_{\alpha, \theta}^{(m)})}{\theta \Gamma(m-\alpha)/\Gamma(1-\alpha)} .$

Noting that for any $i \ge 1$ and for any $\delta > 0$,

$$
\lim_{\theta \to \infty} \frac{a^2(\theta)}{\theta} \log P\left(\left|\left(\frac{\theta}{T}\right)^i - 1\right| \ge \delta\right) = -\infty.
$$

It then follows that

$$
a(\theta)\left(\frac{\theta^{m-1}\Gamma(1-\alpha)}{\Gamma(m-\alpha)}H_m(\mathbf{P}(\alpha,\theta))-1\right)
$$

and

$$
\frac{a(\theta)m(\theta-T)}{\theta} + \frac{a(\theta)(G_{\alpha,\theta}^{(m)} - E(G_{\alpha,\theta}^{(m)}))}{\theta \Gamma(m-\alpha)/\Gamma(1-\alpha)}
$$

are exponentially equivalent, and so they have the same LDP.

The fact that

$$
\inf_{\frac{y\Gamma(1-\alpha)}{\Gamma(m-\alpha)}-mx=z} \Lambda^*(x,y) = \frac{z^2}{2\sigma_{\alpha,m}^2},
$$

combined with Lemma [4.2](#page-13-0) and the contraction principle implies the theorem.

 \Box

5 LDP for Small Parameters

Let

$$
\nabla = \left\{ \mathbf{p} = (p_1, p_2, \ldots) : p_1 \ge p_2 \ge \cdots \ge 0, \sum_{i=1}^{\infty} p_i \le 1 \right\}
$$

be equipped with the subspace topology of $[0,1]^\infty$, and $M_1(\nabla)$ be the space of all probability measures on ∇ equipped with the weak topology. Then $\Pi_{\alpha,\theta}$ belongs to $M_1(\nabla)$.

For any $\delta > 0$, it follows from the GEM representation [\(1.1\)](#page-0-0) that

$$
P\left(X_1^{\alpha,\theta}>1-\delta\right)\le P\left(P_1(\alpha,\theta)>1-\delta\right).
$$

By direct calculation, we have

$$
\lim_{\alpha+\theta\to 0} P\left(X_1^{\alpha,\theta} > 1 - \delta\right) = 1.
$$

Therefore, $\Pi_{\alpha,\theta}$ converges in $M_1(\nabla)$ to $\delta_{(1,0,...)}$ as $\alpha+\theta$ converges to zero. In this section, we establish the LDP associated with this limit. This is a two-parameter generalization to the result in [\[10\]](#page-17-9).

For any $n \geq 1$, set

$$
\nabla_n = \left\{ (p_1, ..., p_n, 0, 0, ...) \in \nabla : \sum_{i=1}^n p_i = 1 \right\},\,
$$

$$
\nabla_{\infty} = \bigcup_{i=1}^{\infty} \nabla_i,
$$

and

$$
a(\alpha, \theta) = \alpha \vee |\theta|, b(\alpha, \theta) = (-\log(a(\alpha, \theta)))^{-1}.
$$

Then we have

Lemma 5.1 The family of laws of $\{P_1(\alpha, \theta) : \alpha + \theta > 0, 0 < \alpha < 1\}$ satisfies a LDP on [0,1] as $a(\alpha, \theta)$ goes to zero with speed $b(\alpha, \theta)$ and rate function

$$
S_1(p) = \begin{cases} 0, & p = 1 \\ k, & p \in [\frac{1}{k+1}, \frac{1}{k}), k = 1, 2, \dots \\ \infty, & p = 0. \end{cases}
$$
 (5.1)

Proof Let $\{X_i^{\alpha,\theta}$ $\alpha_i^{\alpha,\sigma}: i = 1, 2, \ldots$ be defined in [\(1.1\)](#page-0-0). For any $n \geq 1$, set

$$
\tilde{P}_1^n(\alpha, \theta) = \max\{X_i^{\alpha, \theta} : 1 \le i \le n\}.
$$

Then it follows from direct calculation that for any $\delta > 0$

$$
P\{P_1(\alpha,\theta) - \tilde{P}_1^n(\alpha,\theta) > \delta\} \le P\{(1 - U_1) \cdots (1 - U_n) \ge \delta\}
$$

$$
\le \delta^{-1} \prod_{i=1}^n \frac{\theta + i\alpha}{\theta + i\alpha + 1 - \alpha},
$$

which leads to

$$
\limsup_{a(\alpha,\theta)\to 0} b(\alpha,\theta) \log P\{P_1(\alpha,\theta) - \tilde{P}_1^n(\alpha,\theta) > \delta\} \le -n.
$$

Thus the families $\{\tilde{P}_1^n(\alpha,\theta): 0 < \alpha < 1, \theta + \alpha > 0\}_{n=1,2,...}$ are exponential good approximations to the family $\{P_1(\alpha,\theta): 0 < \alpha < 1, \theta + \alpha > 0\}$. By the contraction principle, the family $\{\tilde{P}_1^n(\alpha,\theta):$ $0 < \alpha < 1, \theta + \alpha > 0$ } satisfies a LDP on [0,1] as $a(\alpha, \theta)$ goes to zero with speed $b(\alpha, \theta)$ and rate function

$$
I_n(p) = \begin{cases} 0, & p = 1 \\ k, & p \in [\frac{1}{k+1}, \frac{1}{k}), k = 1, 2, \dots, n-1 \\ n, & \text{else.} \end{cases}
$$

The lemma now follows from the fact that

$$
S_1(p) = \sup_{\delta > 0} \liminf_{n \to \infty} \inf_{|q-p| < \delta} I_n(q).
$$

Theorem 5.1 The family $\{\Pi_{\alpha,\theta} : \alpha + \theta > 0, 0 < \alpha < 1\}$ satisfies a LDP on ∇ as $a(\alpha,\theta)$ goes to zero with speed $b(\alpha, \theta)$ and rate function

$$
S(\mathbf{p}) = \begin{cases} n - 1, & \mathbf{p} \in \nabla_n, p_n > 0, n \ge 1 \\ \infty, & \mathbf{p} \notin \nabla_{\infty}. \end{cases}
$$
 (5.2)

Proof It suffices to establish the LDP for finite dimensional marginal distributions since the infinite dimensional LDP can be derived from the finite dimensional LDP through approximation. For any $n \geq 2$, $(P_1(\alpha, \theta), P_2(\alpha, \theta), \ldots, P_n(\alpha, \theta))$ and $(P_1(0, \alpha + \theta), P_2(0, \alpha + \theta), \ldots, P_n(0, \alpha + \theta))$ have respective joint density functions

$$
h_{\alpha,\theta,n}(p_1,\dots,p_n)=C_{\alpha,\theta,n}\frac{\left(1-\sum_{i=1}^n p_i\right)^{\theta+n\alpha-1}}{\prod_{i=1}^n p_i}P\left(P_1(\alpha,n\alpha+\theta)\leq \frac{p_n}{1-\sum_{i=1}^n p_i}\right),
$$

and

$$
g_{\alpha+\theta,n} = (\alpha+\theta)^n \frac{(1 - \sum_{i=1}^n p_i)^{\theta+\alpha-1}}{\prod_{i=1}^n p_i} P\left(P_1(0, \alpha+\theta) \le \frac{p_n}{1 - \sum_{i=1}^n p_i}\right)
$$

Since $\lim_{a(\alpha,\theta)\to 0} b(\alpha,\theta) \log(\alpha+\theta) = -1$ and $\lim_{a(\alpha,\theta)\to 0} b(\alpha,\theta)C_{\alpha,\theta,n} = -n$, it follows from Lemma 2.4 in [\[10\]](#page-17-9) and Lemma [5.1](#page-15-0) that the family of laws of $(P_1(\alpha, \theta), P_2(\alpha, \theta), \ldots, P_n(\alpha, \theta))$ satisfies a LDP as $a(\alpha, \theta)$ goes to zero with speed $b(\alpha, \theta)$ and rate function

$$
S_n(p_1, ..., p_n) = \begin{cases} 0, & (p_1, p_2, ..., p_n) = (1, 0..., 0) \\ l - 1, & 2 \le l \le n, \sum_{k=1}^l p_k = 1, p_l > 0 \\ n + S_1 \left(\frac{p_n}{1 - \sum_{i=1}^n p_i} \wedge 1 \right), & \sum_{k=1}^n p_k < 1, p_n > 0 \\ \infty, & \text{else.} \end{cases} \tag{5.3}
$$

 \Box

Acknowledgement

Fuqing Gao would like to thank the Department of Mathematics and Statistics at McMaster University for their hospitality during his visit.

References

- [1] Aoki, M. (2006). Thermodynamic limit of macroeconomic or financial models: one- and twoparameter Poisson-Dirichlet models. Preprint.
- [2] Carlton, M. A. (1999). Applications of the two-parameter Poisson-Dirichlet distribution. Unpublished Ph.D. thesis, Dept. of Statistics, University of California, Los Angeles.
- [3] Dawson, D.and Feng, S. (2006). Asymptotic behavior of Poisson-Dirichlet distribution for large mutation rate. Ann. Appl. Probab., 16, 562-582.
- [4] Dawson, D. and Gärtner, J. (1987). Large deviations from the McKean-Vlasov limit for weakly interacting diffusions *Stochastics*, **20**, 247-308.
- [5] Dembo, A. and Zeitouni, O. (1998). Large deviations techniques and applications. Second edition. Applications of Mathematics, 38. Springer-Verlag, New York.
- [6] Derrida, B. (1997). From random walks to spin glasses. Physica D, 107, 186-198.
- [7] Diaconis,P.W. and Kemperman, J. H. B. (1996). Some new tools for Dirichlet priors. Bayesian statistics, 5(Alicante, 1994), pp. 97-106, Oxford Univ. Press, New York.
- [8] Ewens, W. J. (1972). The sampling theory of selectively nuetral alleles. Theor. Pop. Biol., 3, 87–112.
- [9] Feng, S. (2007). Large deviations for Dirichlet processes and Poisson-Dirichlet distribution with two parameters. *Electro. J. Probab.*, **12**, 787-807.
- [10] Feng, S. (2009). Poisson-Dirichlet distribution with small mutation rate. Stoch. Proc. Appl., 116, 2082-2094.
- [11] Feng, S. and Gao, F. Q. (2008). Moderate deviations for Poisson-Dirichlet distribution. Ann. Appl. Probab., 18, 1794-1824.
- [12] Handa, K. (2007). The two-parameter Poisson-Dirichlet point process. arXiv: math/0705.3496v2.
- [13] Kingman, J. F. C. (1975). Random discrete distributions. J. Roy. Stat. Soc. Ser. B, 37, 1-22.
- [14] Perman, M. (1993). Order statistics for jumps of normalised subordinators. Stoch. Proc. Appl., 46, 267-281.
- [15] Perman, M., Pitman, J. and Yor,M. (1992). Size-biased sampling of Poisson point processes and excursions. Prob. Th. Rel. Fields, 92, 21-39.
- [16] Pitman, J. (1992). The two-parameter generalization of Ewen's random partition structure. Technical Report 345, Dept. Statistics, University of California,Berkeley.
- [17] Pitman, J. and Yor, M. (1997). The two-parameter Poisson-Dirichlet distribution derived from a stable subordinator. Ann. Probab., 25, 855-900.
- [18] Tsilevich, N. V. (1997). Distribution of mean values for some random measures. Zap. Nauchn. Sem., POMI 240, 268-279; translation in J. Math. Sci. (New York) 96 (1999), no. 5, 3616- 3623.