

Sequential edge-coloring on the subset of vertices of almost regular graphs

Petros A. Petrosyan

Department of Informatics and Applied Mathematics,
Yerevan State University, 0025, Armenia
Institute for Informatics and Automation Problems,
National Academy of Sciences, 0014, Armenia
E-mail: pet_petros@ipia.sci.am

September 27, 2018

Abstract

Let G be a graph and $R \subseteq V(G)$. A proper edge-coloring of a graph G with colors $1, \dots, t$ is called an R -sequential t -coloring if the edges incident to each vertex $v \in R$ are colored by the colors $1, \dots, d_G(v)$, where $d_G(v)$ is the degree of the vertex v in G . In this note, we show that if G is a graph with $\Delta(G) - \delta(G) \leq 1$ and $\chi'(G) = \Delta(G) = r$ ($r \geq 3$), then G has an R -sequential r -coloring with $|R| \geq \left\lceil \frac{(r-1)n_r+n}{r} \right\rceil$, where $n = |V(G)|$ and $n_r = |\{v \in V(G) : d_G(v) = r\}|$. As a corollary, we obtain the following result: if G is a graph with $\Delta(G) - \delta(G) \leq 1$ and $\chi'(G) = \Delta(G) = r$ ($r \geq 3$), then $\Sigma'(G) \leq \left\lfloor \frac{2n_r(2r-1)+n(r-1)(r^2+2r-2)}{4r} \right\rfloor$, where $\Sigma'(G)$ is the edge-chromatic sum of G .

1 Introduction

In this note we consider graphs which are finite, undirected, and have no loops or multiple edges. Let $V(G)$ and $E(G)$ denote the sets of vertices and edges of a graph G , respectively. The degree of a vertex $v \in V(G)$ is denoted by $d_G(v)$ and the chromatic index of G by $\chi'(G)$. For a graph G , let $\Delta(G)$ and $\delta(G)$ denote the maximum and minimum degrees of vertices in G , respectively. An (a, b) -biregular bipartite graph G is a bipartite graph G with the vertices in one part all having degree a and the vertices in the other part all having degree b . The terms and concepts that we do not define can be found in [5].

A proper edge-coloring of a graph G is a mapping $\alpha : E(G) \rightarrow \mathbf{N}$ such that $\alpha(e) \neq \alpha(e')$ for every pair of adjacent edges $e, e' \in E(G)$. If α is a proper

edge-coloring of a graph G , then $\Sigma'(G, \alpha)$ denotes the sum of the colors of the edges of G . For a graph G , define the edge-chromatic sum $\Sigma'(G)$ as follows: $\Sigma'(G) = \min_{\alpha} \Sigma'(G, \alpha)$, where minimum is taken among all possible proper edge-colorings of G . A proper t -coloring is a proper edge-coloring which makes use of t different colors. If α is a proper t -coloring of G and $v \in V(G)$, then $S(v, \alpha)$ denotes set of colors appearing on edges incident to v . Let G be a graph and $R \subseteq V(G)$. A proper edge-coloring of a graph G with colors $1, \dots, t$ is called an R -sequential t -coloring if the edges incident to each vertex $v \in R$ are colored by the colors $1, \dots, d_G(v)$.

The concept of sequential edge-coloring of graphs was introduced by Asratian [1]. In [1, 2], he proved the following result.

Theorem 1. If $G = (X \cup Y, E)$ is a bipartite graph with $d_G(x) \geq d_G(y)$ for every $xy \in E(G)$, where $x \in X$ and $y \in Y$, then G has an X -sequential $\Delta(G)$ -coloring.

On the other hand, in [2] Asratian and Kamalian showed that the problem of deciding whether a bipartite graph $G = (X \cup Y, E)$ with $\Delta(G) = 3$ has an X -sequential 3-coloring is NP -complete. Some other results on sequential edge-colorings of graphs were obtained in [3, 4]. In particular, in [4] Kamalian proved the following result.

Theorem 2. If G is an $(r-1, r)$ -biregular ($r \geq 3$) bipartite graph with n vertices, then G has an R -sequential r -coloring with $|R| \geq \left\lceil \frac{rn}{2r-1} \right\rceil$.

In this note we generalize last theorem. As a corollary, we also obtain the following result: if G is a graph with $\Delta(G) - \delta(G) \leq 1$ and $\chi'(G) = \Delta(G) = r$ ($r \geq 3$), then $\Sigma'(G) \leq \left\lfloor \frac{2n_r(2r-1) + n(r-1)(r^2+2r-2)}{4r} \right\rfloor$.

2 The Result

Theorem 3. If G a graph with $\Delta(G) - \delta(G) \leq 1$ and $\chi'(G) = \Delta(G) = r$ ($r \geq 3$), then G has an R -sequential r -coloring with $|R| \geq \left\lceil \frac{(r-1)n_r + n}{r} \right\rceil$, where $n = |V(G)|$ and $n_r = |\{v \in V(G) : d_G(v) = r\}|$.

Proof. Since $\chi'(G) = \Delta(G) = r$, there exists a proper r -coloring α of the graph G with colors $1, 2, \dots, r$. For $i = 1, 2, \dots, r$, define the set $V_{\alpha}(i)$ as follows:

$$V_{\alpha}(i) = \{v \in V(G) : i \notin S(v, \alpha)\}.$$

Clearly, for any $i', i'', 1 \leq i' < i'' \leq r$, we have

$$V_{\alpha}(i') \cap V_{\alpha}(i'') = \emptyset \quad \text{and} \quad \bigcup_{i=1}^r V_{\alpha}(i) = V(G) \setminus V_r,$$

where $V_r = \{v \in V(G) : d_G(v) = r\}$.

Hence,

$$n - n_r = |V(G)| - |V_r| = \left| \bigcup_{i=1}^r V_\alpha(i) \right| = \sum_{i=1}^r |V_\alpha(i)|.$$

This implies that there exists i_0 , $1 \leq i_0 \leq r$, for which $|V_\alpha(i_0)| \geq \lceil \frac{n-n_r}{r} \rceil$. Let $R = V_r \cup V_\alpha(i_0)$. Clearly, $|R| \geq n_r + \lceil \frac{n-n_r}{r} \rceil$.

If $i_0 = r$, then α is an R -sequential r -coloring of G ; otherwise define an edge-coloring β as follows: for any $e \in E(G)$, let

$$\beta(e) = \begin{cases} \alpha(e), & \text{if } \alpha(e) \neq i_0, r, \\ i_0, & \text{if } \alpha(e) = r, \\ r, & \text{if } \alpha(e) = i_0. \end{cases}$$

It is easy to see that β is an R -sequential r -coloring of G with $|R| \geq \lceil \frac{(r-1)n_r+n}{r} \rceil$. \square

Corollary 1. If G is an $(r-1, r)$ -biregular ($r \geq 3$) bipartite graph with n vertices, then G has an R -sequential r -coloring with $|R| \geq \lceil \frac{rn}{2r-1} \rceil$.

Corollary 2. If G is a graph with $\Delta(G) - \delta(G) \leq 1$ and $\chi'(G) = \Delta(G) = r$ ($r \geq 3$), then

$$\Sigma'(G) \leq \left\lfloor \frac{2n_r(2r-1) + n(r-1)(r^2+2r-2)}{4r} \right\rfloor$$

Proof. Let α be an R -sequential r -coloring of G with $|R| \geq \lceil \frac{(r-1)n_r+n}{r} \rceil$ described in the proof of Theorem 3. Now, we have

$$\begin{aligned} \Sigma'(G) \leq \Sigma'(G, \alpha) &\leq \frac{\frac{n_r \cdot r(r+1)}{2} + \lceil \frac{n-n_r}{r} \rceil \frac{r(r-1)}{2} + (n - n_r - \lceil \frac{n-n_r}{r} \rceil) \frac{(r+2)(r-1)}{2}}{2} \\ &\leq \frac{\frac{n_r \cdot r(r+1)}{2} + \frac{(n-n_r)r(r-1)}{2r} + (n - n_r - \frac{n-n_r}{r}) \frac{(r+2)(r-1)}{2}}{2} \\ &= \frac{\frac{n_r \cdot r(r+1)}{2} + \frac{(n-n_r)r(r-1)}{2r} + \frac{(n-n_r)(r+2)(r-1)^2}{2r}}{2} \\ &= \frac{n_r \cdot r(r+1)}{4} + \frac{(n-n_r)(r-1)(r^2+2r-2)}{4r} \\ &= \frac{2n_r(2r-1) + n(r-1)(r^2+2r-2)}{4r}. \end{aligned}$$

\square

References

- [1] A.S. Asratian, Investigation of some mathematical model of scheduling theory, Doctoral Thesis, Moscow, 1980.
- [2] A.S. Asratian, R.R. Kamalian, Interval colorings of edges of a multigraph, *Appl. Math.* 5 (1987) 25-34 (in Russian).
- [3] R.R. Kamalian, Interval edge-colorings of graphs, Doctoral Thesis, Novosibirsk, 1990.
- [4] R.R. Kamalian, On a number of vertices with an interval spectrum in proper edge colorings of some graphs, LiTH-MAT-R-2011/03-SE, Linköping University, 2011.
- [5] D.B. West, *Introduction to Graph Theory*, Prentice-Hall, New Jersey, 2001.