

Computation of gcd chain over the power of an irreducible polynomial

Xavier Dahan*

Ochanomizu university, Faculty of General Educational Research & Dept. of Mathematics
xdahan@gmail.com

Abstract. A notion of gcd chain has been introduced by the author at ISSAC 2017 for two univariate monic polynomials with coefficients in a ring $R = k[x_1, \dots, x_n]/\langle T \rangle$ where T is a *primary* triangular set of dimension zero. A complete algorithm to compute such a gcd chain remains challenging. This work treats completely the case of a triangular set $T = (T_1(x))$ in one variable, namely a power of an irreducible polynomial. This seemingly “easy” case reveals the main steps necessary for treating the general case, and it allows to isolate the particular one step that does not directly extend and requires more care.

1 Introduction

Computing gcd is without a doubt one of the most fundamental algorithm in computer algebra and computational aspects have been studied extensively, till today. In [4] is introduced the concept of *gcd chain* to bring a similar notion of the classical gcd of polynomials of one variable over a field, to the case of over a ring $R := k[x_1, \dots, x_n]/\langle T \rangle$ where T is a primary triangular set of dimension zero. Such a ring has nilpotent elements, and non nilpotent elements are invertible. Some attempts to treat this case in prior [4] have concluded in somewhat unsatisfactory solutions. Indeed, a desirable fundamental property of gcd is an ideal equality $\langle a, b \rangle = \langle g \rangle$. While if a and b have coefficients in such a ring of type R , it is well-known that a polynomial g does not exist in general, the outcome of [4] being to present a strategy to circumvent this impediment by “iterating” somehow a Pseudo Remainder Sequence when a nilpotent remainder is met. On the algorithmic side, this raises several challenging questions, even in the seemingly “easy” case of a primary triangular set $T = (T_1(x_1))$ of *one* variable. As this article shows, this case is already not simple. And it is important since it builds the framework to tackle the case of several variables. In particular we identify that all steps, except one that requires more work, extend to more than one variable.

1.1 Motivation

An early motivation in computing gcds over triangular sets come from the *triangular-decomposition* algorithm to solve polynomial (commutative or differential) systems [18,2]. This set of computational methods traces back to the early work of Ritt [16], and the major computational advances realized later by Wu-Wen Tsu [19]. This has lead to several new directions of researches, followed by many researchers. In term of algorithms, only *pseudo-divisions* were initially used. In 1993, in [7] Kalkbrenner introduced a “gcd”-point of view to realize the decomposition, and the elimination (See also the notes [6]). This point of view has later been significantly developed by M. Moreno-Maza *et al.* in particular with the implementation of the library `RegularChains` [5] in the software Maple.

However, such a gcd does not handle “faithfully” polynomials having multiplicities; this question was raised as early as 1995 [13] and later studied furthermore in [10], but without a satisfactory general answer. In this regard, the present work situates in the realm of triangular decomposition as initiated by Wu-Wen Tsu.

The gcd chain has the following *geometric* interpretation. The underlying triangular set can be thought as some algebraic constraints, over which one may want to compute with further polynomials, that is over

* extended version of a presentation made at the conference ADG’2018 (September 12-14, Nanning, China)

the solutions of the constraints *only*. It may happen that the solution (a constraint), is *multiple*. One may think of $t_1(x) = x^3$ for example, in the case where constraints are modeled by a polynomial of one variable like in this work. When computing *over* t_1 , this allows to consider Taylor expansions at order 2 (for example to control the first and second order derivative of the solution=constraint being modeled).

In Example 3 below, we want to compute the constraints defined by both polynomials a and b with coefficients in $\mathbb{R}[x]/\langle t_1 \rangle = \mathbb{R}[x]/\langle x^3 \rangle$. We obtain three cases, displayed in Figure 1.1, as computed by the algorithm of this paper.




Fig. 1. Precision x (left): Three points intersection of $y = 0, -1, 1$ with $x = 0$. Precision x^2 (middle): Two lines, expanded from $y = 0, 1$. Precision x^3 (right): One parabola, expanded from $y = 1$

1.2 Definitions

A primary triangular set in one variable is just a power of an irreducible polynomial p : $T = (T_1(x)) = (p(x)^e)$. The ring $R = k[x]/\langle T \rangle$ is local of maximal ideal $\mathfrak{m} = \langle p \rangle$. It is therefore *Henselian*: monic polynomials admit a unique factorization into coprime factors (not into irreducibles: see [4, § 3.1] for more details). Before giving the technical definitions, a connection with the classical gcd over unique factorization domains will enlighten the differences with the new notion.

Since R is Henselian we can write the unique factorization of the input polynomials a and b into coprime factors as follows:

$$a = a_1 \cdots a_\gamma \cdot a_{\gamma+1} \cdots a_\alpha, \quad b = b_1 \cdots b_\gamma \cdot b_{\gamma+1} \cdots b_\beta. \quad (1)$$

We have ordered the factors so that $\gcd(a_i \bmod \langle p \rangle, b_j \bmod \langle p \rangle) = \rho_i^{\nu_i}$ where $\rho_i \in (k[x]/\langle p \rangle)[y]$ is an irreducible polynomial; And $\nu_i = \min(\lambda_i, \delta_i)$ where $a_i = \rho_i^{\delta_i} + \cdots$ and $b_i = \rho_i^{\lambda_i} + \cdots$ (so that $a_i \equiv \rho_i^{\delta_i} \bmod \langle p \rangle$ and $b_i \equiv \rho_i^{\lambda_i} \bmod \langle p \rangle$). If a and b have some factors a_ℓ and b_m for $\ell, m > \gamma$ then we assume that $\gcd(a_\ell \bmod \langle p \rangle, b_m \bmod \langle p \rangle) = 1$.

Example 1 We reproduce the example of [4, Ex. 3.5, Ex. 5.2]. Consider the polynomials a and b along with their unique factorizations into coprimes modulo $T = x^3$.

$$\begin{aligned} a &= a_1 a_2 a_3 = ((y+1)^2 + x(2y+1) + x^2(y+1))(y+2x+3x^2)(y-1-x-2x^2) \\ b &= b_1 b_2 b_3 = (y+1+2x+x^2)(y+2x+4x^2)(y-1-x-2x^2) \end{aligned}$$

We have $\gamma = 3$ and in each case $\gcd(a_i \bmod \langle x \rangle, b_i \bmod \langle x \rangle)$ is non trivial.

The informal discussion that follows is rigorously detailed in Sections 4.1-4.2 of [4]. Under this point of view, it is convenient to refer to the more common terminology of unique factorization domains, but there is a caveat: “precision”. Modulo p , the gcd of a and b (over the field $k[x]/\langle p \rangle$) is well-defined and is as expected:

$g = \gcd(a_i \bmod \langle p \rangle, b_j \bmod \langle p \rangle) = \prod_{i=1}^{\gamma} \rho_i^{\nu_i}$. But there is no isomorphism $\langle a, b, T \rangle = \langle g, T \rangle$, only the more coarse $\langle a, b, p \rangle = \langle g, p \rangle$ holds. A more refined notion of ‘‘common factors at certain precision’’ allows to obtain an ideal equality $\langle a, b, T \rangle$.

Definition 1. A monic polynomial c is said to be a common factor of a and b at precision ℓ iff the Euclidean divisions of a and b by c have both zero remainder modulo p^ℓ , but at least one non-zero modulo $p^{\ell+1}$.

$$a = ca' + r_a, \quad b = cb' + r_b \quad \Rightarrow \quad r_a, r_b \equiv 0 \bmod \langle p^\ell \rangle, \quad r_a \text{ or } r_b \not\equiv 0 \bmod \langle p^{\ell+1} \rangle.$$

Let \mathcal{I}_1 be the set of indices of common factors of a and b at the smallest precision e_1 . Let \mathcal{I}_2 be the set of indices of common factors a and b at the next to smallest precision e_2 , etc. we obtain a partition $(\mathcal{I}_1, \mathcal{I}_2, \dots, \mathcal{I}_s)$ of the set of common factors of a and b and the associated precision exponents $[e_1, \dots, e_s]$.

Example 2 (Example 1 continued) We notice that $e_1 = 1$ and $\mathcal{I}_1 = \{1\}$. Indeed $y+1$ is the largest common divisor of a_1 and b_1 , and it is modulo $x^{e_1} = x$; There is no common divisor of a_1 and b_1 modulo $x^{e_1+1} = x^2$. Next we observe that $e_2 = 2$ and $\mathcal{I}_2 = \{2\}$: the largest common divisor of a_2 and b_2 is $y+2x$ and it is modulo $x^{e_2} = x^2$ (there is no common divisor of a_2 and b_2 modulo $x^{e_2+1} = x^3$). Last modulo $x^e = x^3$ the factors a_3 and b_3 have a common divisor implying that $\mathcal{I}_3 = \{e_3\}$.

Sections 4.1-4.2 of [4] prove the *existence and uniqueness*, in the case of a triangular set of one variable, of the tuple of indices $(\mathcal{I}_1, \mathcal{I}_2, \dots, \mathcal{I}_s)$ and of the sequence of increasing precision powers $[e_1, e_2, \dots, e_s]$. Before that, with the notations of Eq. (1), denote $G_j^{(a)} := \prod_{i \in \mathcal{I}_j} a_i$ and $G_j^{(b)} := \prod_{i \in \mathcal{I}_j} b_i$, so that:

$$a = \left(\prod_{i=1}^s G_i^{(a)} \right) \cdot a_{\gamma+1} \cdots a_\alpha, \quad b = \left(\prod_{i=1}^s G_i^{(b)} \right) \cdot b_{\gamma+1} \cdots b_\beta. \quad (2)$$

And from the discussion above both $G_i^{(a)}$ and $G_i^{(b)}$ have one maximal common factor which is at precision p^{e_i} . Let us write it G_i (since here it comes alone, it makes no harm to think of G_i as ‘‘the’’ gcd and to write $G_i \equiv \gcd(G_i^{(a)}, G_i^{(b)}) \bmod p^{e_i}$. There exist monic polynomials $c_i^{(a)}$ and $c_i^{(b)}$ both relatively prime modulo p , such that:

$$G_i^{(a)} \equiv c_i^{(a)} G_i \bmod \langle p^{e_i} \rangle, \quad \text{and} \quad G_i^{(b)} \equiv c_i^{(b)} G_i \bmod \langle p^{e_i} \rangle, \quad (3)$$

We obtain an equality of ideals: $\langle a, b, T \rangle = \langle G_1, p^{e_1} \rangle \cdots \langle G_s, p^{e_s} \rangle$. The formal definition of gcd chain modulo a triangular set of one variable is as follows:

Definition 2. Given two monic polynomials $a, b \in R[y]$, a gcd chain of a, b is a sequence $(g_i, p^{e_i})_{i=1, \dots, s}$ such that:

- $e_1 < \dots < e_s \leq e$ and $\deg_y(g_1) > \dots > \deg_y(g_s)$.
- g_i is the product of all common factors of a and b at precision $\geq e_i$.
- g_{i+1} divides g_i modulo p^{e_i+1} . Defining $G_i := g_i/g_{i+1} \bmod p^{e_i+1}$ for $i = 1, \dots, s-1$ and $G_s := g_s$, the following isomorphism holds:

$$R[y]/\langle a, b \rangle \simeq (k[x]/\langle p^{e_1} \rangle)[y]/\langle G_1 \rangle \times \cdots \times (k[x]/\langle p^{e_s} \rangle)[y]/\langle G_s \rangle \quad (4)$$

where the r.h.s is a direct product of rings. We have moreover for all $i = 1, \dots, s$:

$$(k[x]/\langle p^{e_i} \rangle)[y]/\langle a, b \rangle \simeq (k[x]/\langle p^{e_1} \rangle)[y]/\langle G_1 \rangle \times \cdots \times (k[x]/\langle p^{e_{i-1}} \rangle)[y]/\langle G_{i-1} \rangle \times (k[x]/\langle p^{e_i} \rangle)[y]/\langle g_i \rangle \quad (5)$$

Example 3 (Examples 1, 2 continued) Let $p(x) = x$, $T = (T_1(x)) = (x^3) = (p^3)$. The two monic polynomials a and b when expanded are written:

$$\begin{aligned} a &= y^4 + (2x^2 + 3x + 1)y^3 + (-x^2 - x - 1)y^2(-13x^2 - 4x - 1)y - 7x^2 - 2x \\ b &= y^3 + (3x^2 + 3x)y^2 + (-3x^2 - 3x - 1)y - 10x^2 - 2x \end{aligned}$$

According to the discussion made in Example 2, $e_1 = 1$, $e_2 = 2$, $e_3 = 3$. The gcd chain is given by:

$$[(y-1)y(y+1), x], ((y-1-x)(y+2x), x^2), (y-1-x-2x^2, x^3),$$

and yields the following isomorphism according to (4).

$$(k[x]/\langle x^3 \rangle)[y]/\langle a, b \rangle \simeq (k[x]/\langle x \rangle)[y]/\langle y+1 \rangle \times (k[x]/\langle x^2 \rangle)[y]/\langle y+2x \rangle \\ \times (k[x]/\langle x^3 \rangle)[y]/\langle y-1-x-2x^2 \rangle \quad (6)$$

However, Section 5 of [4] dealing with algorithms is more an indication of directions for future work, than a complete and definitive exposition. This is what the present work does, treating the case of one variable completely.

1.3 Related works

First of all, let us clarify what may appear as an elephant in the room:

why not computing the squarefree part of $T = p^e$?

The first reason is that sticking with $T = p^e$ really allows to discover the gcd-chain. Otherwise starting from the gcd known at precision p one could perform Hensel lifting and, roughly speaking, see at each step which parts still divide a and b etc. But then quadratic convergent lifting may not be convenient enough to recover each precision e_i , in that it may miss some. Some refinements should then be devised... with additional costs. The linearly convergent Hensel lifting is not as efficient. The second reason is for considerations of generalization to a primary triangular set of *several* variables. Computing the radical is still possible (see e.g. [15,11]) at a reasonable cost, but this time recovering the factors at the required “precision” becomes far more complicated, assuming it is possible. Consequently, for convenience and generalization in mind,

the squarefree part operation is not considered.

Gcds over non-radical triangular sets have been addressed in [13,10]. But none provided a structural isomorphism that mimics the case of gcd of polynomials over a field. In a different direction, some works have focused on representing not only solution points, but their *multiplicity* as well [3,9,12,1]. The multiplicity is a much coarser information than what provides an ideal isomorphism; Moreover the methods proposed in these works are not simpler than the one proposed in [4] and here. See Section “Previous Work” of [4] for more details. Compared to the algorithmic Section 5 of [4] the present article gives a complete treatment, with Hensel lifting; in particular the over-optimistic over-simplistic Assumption (C) made therein is elucidated with Weierstrass preparation theorem.

When the input irreducible polynomial is $p(x) = x$, the article [14] which computes a “truncated” resultant of bivariate polynomials a and b shares some common features with the present work. However, the gcd-chain carries more information than the classical resultant, and it is not clear whether the quite intricate algorithm that is devised in [14] can be adapted. Therein, a generalized version of the “half-gcd” is proposed whose purpose is a better asymptotic complexity. Another significant difference is the use of Weierstrass factorization to cope with nilpotent leading coefficients. This notion is more natural and efficient than the “normalization” lemma 1 of [14]; It does not require Hensel lifting and to know the squarefree part p of the input $T = p^e$.

Besides, the “clarity” of the classical subresultant based study presented here allows to detect that all tasks used in the algorithm extend quite straightforwardly to primary triangular sets of more than one variable, except one: Algo. 1 “nilpotentFactor” which “removes” nilpotent part. See Section “Concluding remarks” for more about this.

1.4 Organization of the paper

Section 2 introduces the core routine “largestFactor” that executes one iteration of the subresultant algorithm. It finds the “last non-nilpotent subresultant”, uses the Weierstrass factorization to make it, together with the first nilpotent subresultant, monic before another iteration. The complete algorithm “gcdChain” that calls “largestFactor” is presented and analyzed in Section 3. The last section 4 details the variation of the Weierstrass preparation theorem that we have used. Some concluding remarks end the article.

Acknowledgement

The idea of Weierstrass preparation theorem was transmitted to me by M. Moreno-Maza. I am grateful for this precious advice and other discussions we had in March 2018.

2 A subresultant based main algorithm

The outcome of this section is Algorithm 4 “largestFactor” presented in Subsection 2.2. It introduces Weierstrass factorizations at Lines 29 and 35. In the first subsection several subroutines used in this algorithm are introduced first.

2.1 Preliminary routines

Algorithm 1 computes the largest power of p that divides all the coefficients of a polynomial in $(k[x]/\langle p^e \rangle)[y]$. It is a key routine in the present work.

Input: $T = p^e$, power an irreducible polynomial. Nilpotent polynomial $F = F_0 + F_1y + \dots + F_r y^r$ modulo T
Output: $P = p^\ell$, where $\ell \leq e$, and $p^\ell \mid F$ but $p^{\ell+1} \nmid F$
Largest degree i such that F_i is invertible if exists; -1 otherwise.

```

1  $P \leftarrow T$ ,  $i \leftarrow r$ 
2 while  $i > -1$  and  $\deg(P) > 0$  do
3    $P \leftarrow \gcd(F_i, P)$  ;  $i \leftarrow i - 1$ 
4 return  $P, i$ 

```

Algo 1: nilpotentFactor: find the largest power of p that divides F , and T

Proof (Correctness of Algo. 1). For any value of i , one has $P = \gcd(F_r, F_{r-1}, \dots, F_i, T)$. Since $T = p^e$, we can write $p^{\ell_i} := \gcd(F_r, \dots, F_i, T)$. The sequence $\{\ell_i\}_i$ decreases. Let ℓ be the minimal value. By definition $p^\ell \mid F$ since $p^\ell \mid F_j$ for all j , and ℓ is the largest integer having this property. Note that the output P is precisely p^ℓ . On the other hand, if $\deg(P) = 0$, then $\gcd(F_i, p^{\ell_{i-1}}) = 1$ and $\ell_{i-1} > 0$ since otherwise the while loop would have been exited before. We see that F_i is the coefficient of F of largest degree that is invertible — when it exists. \square

Let P, i be the output of Algorithm 1. The first statement in the remark hereunder follows directly from the algorithm. The second one from the classical fact that a polynomial is nilpotent iff all its coefficients are nilpotent.

Remark 1 *If $\deg(P) > 0$ then $i = -1$ and reciprocally. The polynomial F is nilpotent iff $i = -1$.*

A key subroutine Algo. 2 “WeierstrassMonic”, is used to “make monic” a non-nilpotent polynomial. It is based on Corollary 2, which details are postponed to Section 4.

Last, the routine Algo. 3 below computes the subresultant pseudo remainder sequence modulo T and proceeds to the necessary modifications. All specifications are described within the algorithm.

```

Input: Primary triangular set  $T = (p(x)^e)$ 
Univariate polynomial  $f = \dots + f_k y^k + \dots$  in  $R[y]$  where  $R = k[x]/\langle p^e \rangle$ , index  $k$  of the non nilpotent coefficient of  $f$ 
of largest degree
Output: Monic polynomial  $h = y^k + h_{k-1} y^{k-1} + \dots + h_0$  in the Weierstrass factorization theorem (Corollary 2)
In particular  $\langle f, T \rangle = \langle h, T \rangle$  (ideal equality)
5  $(up^e + vf_k = 1) \leftarrow$  Extended Euclidean Algorithm of  $p^e$  and  $f_k$  in  $k[x]$ 
6  $f_k^{-1} \leftarrow v$  // inverse of  $f_k$  modulo  $p^e$ 
7  $y^k = fg + r \leftarrow$  Division of  $y^k$  by  $f$  following Corollary 2 (using  $v$ ), up to precision  $O(y^{k+1})$ 
8 return  $y^k - r$ 

```

Algo 2: WeierstrassMonic

```

Input: Monic polynomials  $a, b$  with  $\deg_y(a) \geq \deg_y(b) > 0$ .
Power of an irreducible polynomial  $T = (p(x)^e)$ .
Output: Subresultant pseudo-remainder sequence modulo  $T$ :  $\mathbf{S} = [S_{r_0} = a, S_{r_1} = b, S_{r_2}, \dots, S_{r_t}, \dots]$ 
Index  $j$  such that  $S_{r_j}$  is not nilpotent and  $S_{r_{j+1}}$  is nilpotent or zero
Integer  $i$  such that  $\text{coeff}(S_{r_j}, i)$  is the non-nilpotent coefficient of largest degree
If  $S_{r_{j+1}}$  is not zero, largest degree polynomial  $P(x) = p(x)^\ell \in k[x]$  such that  $P|S_{r_{j+1}}$ .
9 try
10 | Compute a subresultant sequence modulo  $T$ :  $\mathbf{S} = [S_{r_0} = a, S_{r_1} = b, S_{r_2}, \dots, S_{r_t}, \dots]$ 
11 |  $j + 1 \leftarrow$  index of last non-zero subresultant
12 |  $i_{new} \leftarrow -1$ ;  $P_{new} \leftarrow 0$ 
13 catch (“Error: Leading coefficient of  $S_{r_{t+2}}$  is not invertible”) // division failed at the  $r_{t+2}$ -th subresultant
14 |  $j \leftarrow t + 1$ ;  $P_{new}, i_{new} \leftarrow$  nilpotentFactor( $S_{r_j}, T$ )
15 | if  $i_{new} > -1$  then //  $S_{r_j}$  is not nilpotent
16 | |  $A \leftarrow$  WeierstrassMonic( $S_{r_j}, T, i_{new}$ )
17 | |  $-, i \leftarrow$  nilpotentFactor( $S_{r_{j-1}}, T$ ) //  $\text{coeff}(S_{r_{j-1}}, i)$  is invertible
18 | |  $B \leftarrow$  WeierstrassMonic( $S_{r_{j-1}}, T, i$ )
19 | | return modifiedSubres( $A, B, T$ )
20 while  $i_{new} = -1$  do //  $S_{r_j}$  is nilpotent
21 | |  $i_{old} \leftarrow i_{new}$ ;  $P_{old} \leftarrow P_{new}$ ;  $j \leftarrow j - 1$ 
22 | |  $P_{new}, i_{new} \leftarrow$  nilpotentFactor( $S_{r_j}, T$ )
23 return  $\mathbf{S}, j, i_{new}, P_{old}$ 

```

Algo 3: modifiedSubres: Modified subresultant pseudo-remainder sequence

Lemma 1 (Correctness of Algo. 3) *The output \mathbf{S}, j, i, P satisfies the specifications mentioned in “output”.*

Proof. Step 1: The computation of the subresultant p.r.s. modulo T raises no exception at Line10. This means that the full chain has been successfully computed, in particular the last subresultant is zero. By definition of the index j at Line 11, its index is r_{j+2} and $S_{r_{j+1}}$ is the last non zero (mod T).

It goes next to Line 20. The “return” occurs exactly when $i_{new} > -1$ or equivalently that S_{r_j} is not nilpotent; Additionally that $S_{r_{j+1}}$ is nilpotent or zero. The specification of Algorithm “nilpotentFactor” tell that i_{new} is the largest degree of S_{r_j} whose coefficient $c_{i_{new}}$ is not nilpotent. Since a polynomial is nilpotent iff all its coefficients are nilpotent, this implies that S_{r_j} is not nilpotent. Since the while loop (Line 20) is bottom-up on the subresultant chain, j is the *first* index for which the corresponding subresultant is not nilpotent. Therefore $S_{r_{j+1}}$ is nilpotent, and the specifications of Algorithm “nilpotentFactor” imply that $i_{old} = -1$ and that $P_{old} | S_{r_{j+1}}$. This shows that all the requirements sateted in the “output” of Algo. 3 are satisfied.

Step 2: The computation of the subresultant p.r.s. failed, an exception is caught. Lines 10-11 are skept to go directly to Line 14. The leading coefficient of S_{r_t} is not invertible therefore the subresultant $S_{r_{t+2}}$ *cannot*

be computed (mod T) following the (classical subresultant p.r.s.) formula:

$$S_{r_{t+2}}(a, b) = \pm \frac{\text{prem}(S_{r_t}, S_{r_{t+1}})}{c_t^{r_t - r_{t+1} + 1} \text{lc}(S_{r_t})}, \quad \begin{cases} c_1 = 1, & r_i = \deg(S_{r_i}) \\ c_i = \text{lc}(S_{r_i})^{r_{i-1} - r_i} c_{i-1}^{r_i - r_{i-1} + 1}. \end{cases}$$

Nonetheless, S_{r_t} and $S_{r_{t+1}}$ have been both computed successfully. Index j is set to $t + 1$. If $i_{new} > -1$ then $S_{r_{t+1}}$ is not nilpotent, therefore neither is S_{r_t} . Thus the call to Algo. “WeierstrassMonic” is valid at Lines 16 and 18. And the recursive call to “modifiedSubres” makes sense. Eventually, no exception is raised, or S_{r_j} is nilpotent (at Line 14). Moreover the recursive call is consistent since $\langle a, b, T \rangle = \langle S_{r_t}, S_{r_{t+1}}, T \rangle = \langle A, B, T \rangle$.

If $i_{new} = -1$ then S_{r_j} is nilpotent, but it may not be the first nilpotent subresultant. It goes next to the while loop (line 20) and the proof follows that of Step 1. \square

2.2 Algorithm “largestFactor”

Now that the subroutines are defined, we describe the main algorithm of this section. It builds upon the subresultant algorithm, where instead of computing the “last non-zero” one, it computes the “last non-nilpotent” subresultant. Besides the comments and specifications, the proof of correctness discusses in details of the several steps.

<p>Input: Monic polynomials f, A with $\deg_y(f) \geq \deg_y(A)$. Power of an irreducible polynomial $T = (p(x))^e$.</p> <p>Output: g, p^{e_1}, B, p^ℓ or $-$ where g is monic and $\langle g, p^{e_1} \rangle = \langle f, A, p^{e_1} \rangle$; $B = \text{“end”}$; or $B \in k[y]$ monic, $\deg_y(g) \geq \deg_y(B)$; $p^\ell = p^{e - e_1}$ is used for iterative calls in Algorithm 5</p> <pre style="font-family: monospace; font-size: 0.9em;"> 24 if A = 0 then // Finished: No iteration necessary 25 return f, T, “end”, - 26 if deg(A) = 0 then // Finished: No iteration necessary 27 return A, T, “end”, - 28 S, j, i, p^{e₁} ← modifiedSubres(A, f, T) // S_{r_j} is not nilpotent, S_{r_{j+1}} is. p^{e₁} S_{r_{j+1}} 29 g ← WeierstrassMonic(S_{r_j}, T, i) // Put S_{r_j} in monic form 30 if p^{e₁} = T then // case where e₁ = e 31 return g, T, “end”, - 32 S ← S_{r_{j+1}}/p^{e₁} // Precision loss of deg(p^{e₁}); S is no more nilpotent 33 p^ℓ ← T/p^{e₁} = p^{e - e₁} 34 -, i ← nilpotentFactor(S, p^ℓ) // coeff(S, i) is invertible 35 B ← WeierstrassMonic(S, p^ℓ, i) // Put S in monic form 36 return g, p^{e₁}, B, p^ℓ </pre>

Algo 4: The Largest Common Factor (largestFactor)

Lemma 2 (Correctness of Algo. 4) *The output of Algorithm 4 verifies: $\langle f, A, p^{e_1} \rangle = \langle g, p^{e_1} \rangle$.*

Proof. If the algorithm exits at Line 25 then $\langle f, A, T \rangle = \langle f, 0, T \rangle = \langle f, T \rangle$. It then returns $f, T, \text{“end”}$, hence $g = f$ and $p^{e_1} = T$, and $\langle f, T \rangle = \langle g, p^{e_1} \rangle$. The gcd chain has one block.

If the algorithm exits at Line 27 then A is monic constant, hence equal to 1; therefore $\langle f, A, T \rangle = \langle A, T \rangle = \langle T, 1 \rangle = \langle 1 \rangle$. It returns $A, T, \text{“end”}$, hence $g = A$ and $p^{e_1} = T$, thereby $\langle g, p^{e_1} \rangle = \langle A, T \rangle$. Here too the gcd chain has one block.

Assume now that the algorithm ends at Line 36 or at Line 31. At Line 28, according to the specifications of Algo. 3, S_{r_j} is not nilpotent and $S_{r_{j+1}}$ is either nilpotent or zero (mod T). Moreover p^{e_1} is the largest power of $p(x)$ that divides $S_{r_{j+1}}$, and $\text{coeff}(S_{r_j}, i)$ is the coefficient of largest degree that is invertible. We

can apply Algorithm 2 “WeierstrassMonic” at Line 29; It outputs a monic polynomial g such that $S_{r_j} = u \cdot g$ where u is a unit in $(k[x]/\langle p^{e_1} \rangle)[y]$. In particular $\langle g, p^{e_1} \rangle = \langle S_{r_j}, p^{e_1} \rangle$ (§).

If the test of Line 30 is satisfied, then S_{r_j} is not only the last non-nilpotent subresultant but also the last non-zero one: classical subresultant theory insures that $\langle S_{r_j} \rangle = \langle f, A \rangle$ modulo T (that is $e_1 = e$). From the specifications of the “WeierstrassMonic” algorithm we have $\langle g \rangle = \langle S_{r_j} \rangle$ and thus $\langle g, T \rangle = \langle f, A, T \rangle$ (with $T = p^{e_1}$ since $e = e_1$). The output Line 36 is then in accordance with the specifications.

If this test is not satisfied, more data needs to be output since additional works must be performed, like Hensel lifting. We compute then $S = S_{r_{j+1}}/p^{e_1}$ at Line 32 which is not nilpotent. By the “last non-nilpotent criterion” of subresultant [4, Thm 5.1] $\langle S_{r_j}, p^{e_1} \rangle = \langle f, A, p^{e_1} \rangle$. Hence $\langle g, p^{e_1} \rangle = \langle f, A, p^{e_1} \rangle$, by (§), as required.

Similarly, Algorithm 2 at Line 35 returns a monic polynomial B such that $S = v \cdot B$, v being a unit modulo p^ℓ . Note that B is monic and satisfies $\deg_y(B) \leq \deg_y(g)$, as required. \square

Lemma 3 *If a monic polynomial $c \neq 1$ is a common factor of f and A at precision $r > e_1$ then c is a monic divisor of B at precision $r - e_1$. A monic divisor $c \neq 1$ of B is not a common factor of f and A at precision $< e_2$.*

Proof. Let c be a factor of f and A at precision $r > e_1$. We can write: $f \equiv cf' \pmod{p^r}$ and $A \equiv cA' \pmod{p^r}$ (by Definition 1, both equalities do not hold together modulo p^{r+1}). Consider the Bézout coefficients (a.k.a cofactors) u_ν, v_ν of the subresultant S_{r_ν} . From the equality $S_{r_{j+1}} = u_{j+1}f + v_{j+1}A$, one deduce that $S_{r_{j+1}} \equiv c(u_{j+1}f' + v_{j+1}A') \pmod{p^r}$. We also have $p^{e_1}Bv \equiv c(u_{j+1}f' + v_{j+1}A') \pmod{p^r}$. Now c being monic p^{e_1} does not divide c , and we have: $B \equiv c \frac{v^{-1}u_{j+1}f' + v^{-1}v_{j+1}A'}{p^{e_1}} \pmod{p^{r-e_1}}$. Since $r > e_1$ by assumption, $B \equiv 0 \pmod{\langle p^{r-e_1}, c \rangle}$ and c is a factor of B at precision at least $r - e_1 \geq 1$. Moreover, since p^{e_1} is the largest power of p that divides $S_{r_{j+1}}$, the precision is exactly $r - e_1$. This proves the first assertion.

Consider a monic divisor C of B at precision $< e_2$. This means that the remainder of the Euclidean division of B by C is in $\langle p \rangle$ but not in $\langle p^{e_2} \rangle$. If C were a common factor of A and f at precision $\ell < e_2$, then it is at precision $\ell \leq e_1$ since there is no common factor of f and A at precision $e_1 + 1, \dots, e_2 - 1$, by definition. We would obtain $p^{e_1}B \equiv v^{-1}(u_{j+1}Cf' + v_{j+1}CA' + p^\ell W) \pmod{\langle p^\ell \rangle}$ for a non-zero polynomial W , and $B \equiv C \frac{v^{-1}u_{j+1}f' + v^{-1}v_{j+1}A'}{p^{e_1}} + p^{\ell-e_1}W \pmod{\langle p^{e-e_1} \rangle}$. For C to divide B modulo p^ℓ , the term $p^{\ell-e_1}W$ shall be zero at least modulo p . This happens only if $\ell > e_1$ since $W \neq 0$. Contradiction with $\ell \leq e_1$. \square

3 The Gcd-Chain Algorithm

Now that all sub-routines used in Algorithm 4 “largestFactor” are defined, we introduce in this section main Algorithm 5 below. It is made of a while loop that has two main components:

1. a call to Algorithm 4 “largestFactor” (line 40) . A loss of precision is entailed in the division at Line 32.
2. recovering this precision loss at each iteration by Hensel lifting (Lines 43-51).

Each iteration computes one “block” of the gcd-chain. The correctness is shown in Theorem 1, with several preliminaries made of Lemmas 4-6, Proposition 1 and its corollary 1. These preliminaries contain a detailed description of each step of the algorithm.

Remark 2 *In the sequel the triangular sets T_i, T'_i are all power of the irreducible polynomial p . These exponents denoted e_i or e'_i are introduced only for the analysis and are not needed in the algorithms. This is consistent with the principle made in § 1.3 to not compute squarefree parts.*

Lemma 4 *The Hensel lifting at Line 48 of Algorithm 5 returns G_i^*, g_{i+1}^* verifying: $G_i^* \equiv G_i \pmod{\langle T'_{i+1} \rangle}$, $g_{i+1}^* \equiv g_{i+1} \pmod{\langle T'_{i+1} \rangle}$, $g_i \equiv G_i^* g_{i+1}^* \pmod{\langle T_{i+1} \rangle}$.*

Proof. This is classical: the algorithm follows exactly the steps presented in Algorithm 15.10 of [17]. \square

The next lemma clarifies the status of the input/output when calling Algorithm “largestFactor” at Line 40, within the several iterations. The notations introduced will be used thereafter.


```

Input: Power of an irreducible polynomial  $T = (p(x))^e$ 
Univariate polynomials  $a$  and  $b$  in  $R[y]$  where  $R = k[x]/\langle T \rangle$ 
Output: Lists of polynomials  $\mathcal{C} = [g_1, \dots, g_s]$ ,  $\mathcal{D} = [G_1, \dots, G_{s-1}]$ 
List of polynomials equal to powers of the irreducible polynomial  $p$ :  $\mathcal{T} = [p^{e_1}, p^{e_2}, \dots, p^{e_s}]$  such that
 $[(g_1, p^{e_1}), \dots, (g_s, p^{e_s})]$  is the gcd-chain of  $(a, b, T)$ 
37  $g_0 \leftarrow a$ ;  $B_0 \leftarrow b$ ;  $S_0 \leftarrow T$ ;  $i \leftarrow 0$ ;  $e_0 \leftarrow 0$ ;  $T_0 \leftarrow T$ 
38  $\mathcal{C} \leftarrow []$ ;  $\mathcal{D} \leftarrow []$ ;  $\mathcal{T} \leftarrow []$ 
39 while  $B_i \neq \text{"end"}$  and  $B_i \neq 1$  do // Algo. 4 ended Line 25 or 27 or 31
40    $g_{i+1}, T'_{i+1}, B_{i+1}, S_{i+1} \leftarrow \text{largestFactor}(g_i, B_i, S_i)$  // Iteration of Algo. 4
41   if  $i = 0$  then // First iteration treated apart
42      $\mathcal{T} \leftarrow \mathcal{T} \text{ cat } [T'_{i+1}]$ 
43   else // Trigger Hensel Lifting from second iteration
44      $G_i \leftarrow g_i / g_{i+1} \bmod \langle T'_{i+1} \rangle$  //  $g_i = G_i g_{i+1} \bmod \langle T'_{i+1} \rangle$ 
45      $\alpha, \beta \leftarrow \text{xgcd}(G_i, g_{i+1}) \bmod \langle T'_{i+1} \rangle$  //  $\alpha G_i + \beta g_{i+1} \equiv 1 \bmod \langle T'_{i+1} \rangle$ 
46      $T_{i+1} \leftarrow T_i \cdot T'_{i+1}$  //  $T_{i+1} = p^{e_{i+1}}$ 
47      $\text{precision} \leftarrow \text{deg}(T_{i+1})$ 
48      $G_i^*, g_{i+1}^* \leftarrow \text{HenselLift}(g_i, G_i, g_{i+1}, \alpha, \beta, T'_{i+1}, \text{precision})$ 
//  $G_i^* \equiv G_i \bmod \langle T'_{i+1} \rangle$ ,  $g_{i+1}^* \equiv g_{i+1} \bmod \langle T'_{i+1} \rangle$ ,  $g_i \equiv G_i^* g_{i+1}^* \bmod \langle T_{i+1} \rangle$ 
49      $g_{i+1} \leftarrow g_{i+1}^* \bmod \langle T_{i+1} \rangle$ 
50      $\mathcal{D} \leftarrow \mathcal{D} \text{ cat } [G_i^* \bmod \langle T_i \rangle]$ 
51      $\mathcal{T} \leftarrow \mathcal{T} \text{ cat } [T_{i+1}]$  // Next block computed
52    $\mathcal{C} \leftarrow \mathcal{C} \text{ cat } [g_{i+1}]$ 
53    $i \leftarrow i + 1$ 
54 return  $\mathcal{C}, \mathcal{D}, \mathcal{T}$ 

```

Algo 5: The gcd-chain algorithm

Lemma 5 Write $\mathcal{T} = [T_1, \dots, T_s]$ with $T_i = p^{e_i}$, $\mathcal{C} = [g_1, \dots, g_s]$ and $\mathcal{D} = [G_1, \dots, G_{s-1}]$ the three outputs of Algorithm 5. For all $i = 0, \dots, s-1$, consider the output $g_{i+1}, T'_{i+1}, B_{i+1}, S_{i+1}$ of the call to “largestFactor” at Line 40, and T_{i+1} as defined at Line 46. They are related as follows:

$$T'_{i+1} = p^{e_{i+1}-e_i}, \quad S_{i+1} = p^{e-e_{i+1}}, \quad T_{i+1} = p^{e_{i+1}} \quad (\text{with } e_0 := 0).$$

Proof. The proof goes by induction on $i = 0, \dots, s-1$. According to the specifications of Algorithm 4 “largestFactor” (Line 40) proved in Lemma 2. we have when $i = 0$, $T'_1 = p^{e_1}$, $S_1 = p^{e-e_1}$. Note that the case $i = 0$ (Line 41) does not require Hensel lifting, and thus $T_1 = T'_1 = p^{e_1-e_0} = p^{e_1}$, as required.

By induction hypothesis we can assume that the result holds up to $i < s-1$. There is an $i+1$ -th call to Algorithm 4 “largestFactor” (at Line 40), with input $S_i = p^{e-e_i}$, by induction hypothesis. From the Specification of Algorithm 4, the output is $T'_{i+1} = p^{e_{i+1}-e_i}$ because the input S_i is only at precision $e-e_i$; And $S_{i+1} = p^{e-e_i-(e_{i+1}-e_i)} = p^{e-e_{i+1}}$ as required. The definition of T_{i+1} at Line 46 gives $T_{i+1} = T'_{i+1} \cdot T_i = p^{e_{i+1}-e_i} p^{e_i} = p^{e_{i+1}}$ as required. \square

The proposition below connects the different outputs obtained after each iteration, to the initial input a, b, T . It is crucial in the proof of Theorem 1

Proposition 1. *With the notation of Lemma 5, we have*

$$\text{for } i \geq 1, \quad p^{e_{i-1}} g_i \in \langle a, b, p^e \rangle, \quad p^{e_i} B_i \in \langle a, b, p^e \rangle.$$

Proof. We proceed by induction on i , starting with the base case $i = 1$.

By definition g_1 is the first output polynomial of the first call to Algorithm 4 “largestFactor” in Algorithm 5 “gcdChain”. From the definition of the Weierstrass factorization at Line 29 of Algorithm 4, there is a unit $\nu_1 \in (k[x]/\langle p^e \rangle)[y]$ such that $\nu_1 g_1 = S_{r_j}$. Write the Bézout coefficients $u_j^{(1)} a + v_j^{(1)} b = S_{r_j}$, so that $g_1 = \nu_1^{-1} (u_j^{(1)} a + v_j^{(1)} b)$; this proves that $g_1 \in \langle a, b, T \rangle$ (and that $g_1 \in \langle a, b, p^e \rangle$: indeed $p^e = T$).

B_1 is the third output polynomial of the first call to Algorithm 4 “largestFactor” in Algorithm 5 “gcdChain”. From Line 35 of Algorithm 4 and by definition of the Weierstrass factorization, there is a unit $\epsilon_1 \in (k[x]/\langle p^{e-e_1} \rangle)[y]$ such that $p^{e_1} \epsilon_1 B_1 = S_{r_{j+1}}$. With the Bézout coefficients written $S_{r_{j+1}} = u_{j+1}^{(1)} a + v_{j+1}^{(1)} b$, we obtain $p^{e_1} B_1 = \epsilon_1^{-1} (u_{j+1}^{(1)} a + v_{j+1}^{(1)} b) \in \langle a, b, p^e \rangle$.

Next assume that the result holds up to a value $1 \leq i < s$ and let us prove it for $i+1$. By definition g_i (resp. B_i) is the first (resp. the third) output polynomial of the i -th call to Algorithm 4 “largestFactor” in Algorithm 5 “gcdChain”. From the definition of the Weierstrass factorization at Line 29 (resp. at Line 35) of Algorithm 4, there is a unit $\nu_{i+1} \in (k[x]/\langle p^{e-e_i} \rangle)[y]$ (resp. ϵ_{i+1}) such that $g_{i+1} \nu_{i+1} = S_{r_j}$ (resp. $p^{e_{i+1}-e_i} \epsilon_{i+1} B_{i+1} = S_{r_{j+1}}$). Write the Bézout coefficients as follows:

$$S_{r_j} = u_j^{(i+1)} g_i + v_j^{(i+1)} B_i, \quad S_{r_{j+1}} = u_{j+1}^{(i+1)} g_i + v_{j+1}^{(i+1)} B_i.$$

It follows that:

$$p^{e_i} g_{i+1} \stackrel{(\bullet)}{=} \nu_{i+1}^{-1} (u_j^{(i+1)} p^{e_i} g_i + v_j^{(i+1)} p^{e_i} B_i), \quad p^{e_{i+1}-e_i} B_{i+1} = \epsilon_{i+1}^{-1} (u_{j+1}^{(i+1)} g_i + v_{j+1}^{(i+1)} B_i).$$

The latter equality implies $p^{e_{i+1}} B_{i+1} \stackrel{(\star)}{=} \epsilon_{i+1}^{-1} (u_{j+1}^{(i+1)} p^{e_i} g_i + v_{j+1}^{(i+1)} p^{e_i} B_i)$. By induction hypothesis $p^{e_{i-1}} g_i \in \langle a, b, p^e \rangle$ and $p^{e_i} B_i \in \langle a, b, p^e \rangle$. Thus $p^{e_i} g_i = p^{e_i-e_{i-1}} p^{e_{i-1}} g_i \in \langle a, b, p^e \rangle$. Consequently, both the r.h.s of Eqs. (\star) and (\bullet) are in $\langle a, b, p^e \rangle$, therefore so are $p^{e_i} g_{i+1}$ and $p^{e_{i+1}} B_{i+1}$. \square

The first iteration of the loop of Line 39 is special since it doesn’t require Hensel Lifting (case $i = 0$). The lemma hereunder treats this base case apart, and used in the induction proof of Theorem 1.

Lemma 6 *The first iteration of the while loop (Line 39, $i = 0$) of Algorithm 5 fills \mathcal{C} with g_1 and \mathcal{T} with $T'_1 = p^{e_1}$. We have $\langle g_1, p^{e_1} \rangle = \langle a, b, p^{e_1} \rangle$.*

Moreover, if there is only one iteration the stronger equality $\langle g_1, p^{e_1} \rangle = \langle a, b, p^e \rangle$ holds.

Proof. From the proof of correctness of Algorithm 4 one has at Line 40 of Algorithm 5 $\langle g_1, T'_1 \rangle = \langle g_0, B_0, T'_1 \rangle$. But $T'_1 = p^{e_1}$, $g_0 = a$ and $B_0 = b$ yielding the first equality.

For the second one, by Proposition 1 we know that $p^{e_0} g_1 = g_1 \in \langle a, b, p^e \rangle$ and that $p^{e_1} B_1 \in \langle a, b, p^e \rangle$. If there is only one iteration of the while loop Line 39, namely for $i = 0$, then there are two sub-cases. First $B_1 \neq \text{“end”}$ and $B_1 = 1$. Then Proposition 1 gives $p^{e_1} B_1 = p^{e_1} \in \langle a, b, p^e \rangle$ which proves $\langle p^{e_1}, g_1 \rangle = \langle a, b, p^e \rangle$. Second, $B_1 = \text{“end”}$. This happens when the call to Algorithm “largestFactor” (Line 40) with input a, b, T outputs $B_{i+1} = B_1 = \text{“end”}$. According to Lines 25, 27, 31 the output denoted p^{e_1} (in this Lemma) is equal to T (as denoted in Algorithm 4) which is equal to p^e . This implies $\langle g_1, p^{e_1} \rangle = \langle g_1, p^e \rangle = \langle a, b, p^e \rangle$. \square

To prove correctness of the “gcdChain” Algorithm 5, one must consider the isomorphisms (4) and (5). This amounts to prove that a product of ideals of type $\langle G_1, p^{e_1} \rangle \langle G_2, p^{e_2} \rangle \cdots \langle g_i, p^{e_i} \rangle$ is equal to ideals of type $\langle a, b, p^{e_i} \rangle$. Thus we must examine the generators of the product of ideals, which is the purpose of the following corollary (of Proposition 1).

Corollary 1. *The same notations as in Lemma 5 are used. Let \mathcal{I} be a subset of indices in $\llbracket 1, s-1 \rrbracket$ and let $\bar{\mathcal{I}}$ its complement in $\llbracket 1, s \rrbracket$ (so that $\mathcal{I}, \bar{\mathcal{I}}$ is a partition $\llbracket 1, s \rrbracket$).*

Then $g_s \prod_{i \in \mathcal{I}} p^{e_i} \prod_{j \in \bar{\mathcal{I}}} G_j \in \langle a, b, p^e \rangle$.

Proof. Assume that $\mathcal{I} \neq \emptyset$ first and let $\varsigma := \max \mathcal{I}$. Then $\{\varsigma + 1, \dots, s-1\} \subset \bar{\mathcal{I}}$ (possibly empty). We claim that $p^{e_\varsigma} g_s \prod_{j=\varsigma+1}^{s-1} G_j \in \langle a, b, p^e \rangle$, from which the conclusion follows since $\langle g_s \prod_{i \in \mathcal{I}} p^{e_i} \prod_{j \in \bar{\mathcal{I}}} G_j \rangle \subset \langle p^{e_\varsigma} g_s \prod_{j=\varsigma+1}^{s-1} G_j \rangle$.

From the definition of G_i and g_i after Hensel Lifting at Lines 50 and 49 of Algorithm 5 one has: $g_{i-1} = G_{i-1} g_i \bmod p^{e_{i-1}}$ (see Lemma 4). It implies the following equality where (\cdots) denotes a polynomial:

$$\begin{aligned} \left(\prod_{j=\varsigma+1}^{s-1} G_j \right) g_s &= \left(\prod_{j=\varsigma+1}^{s-2} G_j \right) (g_{s-1} + p^{e_{s-1}} g_s (\cdots)) = \left(\prod_{j=\varsigma+1}^{s-3} G_j \right) (g_{s-2} + p^{e_{s-1}} g_{s-1} (\cdots) + p^{e_{s-1}} g_s (\cdots)) \\ &= g_{\varsigma+1} + p^{e_{\varsigma+1}} g_{\varsigma+2} (\cdots) + \cdots + p^{e_{s-3}} g_{s-2} (\cdots) + p^{e_{s-2}} g_{s-1} (\cdots) + p^{e_{s-1}} g_s (\cdots) \end{aligned}$$

If we multiply the above equation by p^{e_ζ} then each term is a multiple of $p^{e_\ell} g_{\ell+1}$ for some $\ell = \zeta, \dots, s-1$. Hence, by Proposition 1 each term is in $\langle a, b, p^e \rangle$ yielding the conclusion in the case $\mathcal{I} \neq \emptyset$.

Otherwise when $\mathcal{I} = \emptyset$, the product is equal to $G_1 \cdots G_{s-1} g_s$ and similarly to the above, we get $g_1 + p^{e_1} g_2(\cdots) + \cdots + p^{e_{s-1}} g_s(\cdots)$, which is in $\langle a, b, p^e \rangle$ according to Proposition 1. \square

The proof of Theorem 1 requires a last intermediate result.

Lemma 7 *With the notations of Lemma 5 assume that $s \geq 2$. Fix an $1 \leq i \leq s-1$ and refer to the notations of Eqs (2)-(3), to define $a' := G_{i+1}^{(a)} \cdots G_s^{(a)} \cdot a_{\gamma+1} \cdots a_\alpha$ and $b' := G_{i+1}^{(b)} \cdots G_s^{(b)} \cdot b_{\gamma+1} \cdots b_\beta$. We have:*

1. $\langle a, b, p^{e_{i+1}} \rangle \subset \langle G_1, p^{e_1} \rangle \cdots \langle G_i, p^{e_i} \rangle \langle a', b', p^{e_{i+1}} \rangle$.
2. $\langle a', b', p^{e_{i+1}} \rangle = \langle g_{i+1}, p^{e_{i+1}} \rangle$.

Proof. By Eqs.(2)-(3), we have $a = a' \prod_{j=1}^i G_j^{(a)}$ (similarly $b = b' \prod_{j=1}^i G_j^{(b)}$) with $G_j^{(a)} \equiv c_j^{(a)} G_j \pmod{\langle p^{e_j} \rangle}$ (respectively $G_j^{(b)} \equiv c_j^{(b)} G_j \pmod{\langle p^{e_j} \rangle}$). Therefore

$$a = a' \cdot \prod_{j=1}^i (c_j^{(a)} G_j + p^{e_j}(\cdots)), \quad b = b' \cdot \prod_{j=1}^i (c_j^{(b)} G_j + p^{e_j}(\cdots)),$$

where (\cdots) denotes some polynomials. This proves the inclusion 1.

To prove 2, let us treat first the case $i = 1$. Consider the polynomials g_1, B_1 , which are the output of the first call to “largestFactor” at Line 40 of Algorithm 5. Note that if we apply the lemma 6 with g_1, B_1, p^{e-e_1} instead of a, b, T , we obtain the ideal equality: $\langle g_1, B_1, p^{e_2-e_1} \rangle = \langle g_2, p^{e_2-e_1} \rangle$; Note the precision loss of e_1 entailed at the previous iteration $i = 0$. Let us prove that $\langle a', b', p^{e_2-e_1} \rangle = \langle g_1, B_1, p^{e_2-e_1} \rangle$. According to Definition 1 and to the definitions of the polynomials a' and b' , the common factors of a' and b' are exactly those of a and b at precision $\geq e_2$. By Lemma 3, such common factors are divisors of B_1 at precision $\geq e_2 - e_1$. On the other hand, by Lemma 2 the divisors of g_1 (at precision p) are *exactly* the common factors of a and b . Therefore, *up to precision*, the common factors of g_1 and B_1 are *exactly* those of a and b at precision $\geq e_2$, hence are the same as those of a' and b' . As for *precision*, the common factors at precision $\geq e_2$ of a, b , and those of g_1, B_1 (and they are at precision $\geq e_2 - e_1$ due to precision loss). We deduce that $\langle g_2, p^{e_2-e_1} \rangle = \langle a', b', p^{e_2-e_1} \rangle$. After Hensel lifting at Line 48, we obtain $\langle g_2, p^{e_2} \rangle = \langle a', b', p^{e_2} \rangle$.

The general case $i > 1$ reduces to the case $i = 1$ by considering instead of a and b , $E := a/G_1^{(a)} \cdots G_{i-1}^{(a)} = a_{\gamma+1} \cdots a_\alpha \cdot \prod_{j=i}^s G_j^{(a)}$ and respectively $F := b/G_1^{(b)} \cdots G_{i-1}^{(b)} = b_{\gamma+1} \cdots b_\beta \cdot \prod_{j=i}^s G_j^{(b)}$; And instead of a' and b' to take $E' := a'/G_1^{(a)} \cdots G_{i-1}^{(a)} = a_{\gamma+1} \cdots a_\alpha \cdot \prod_{j=i+1}^s G_j^{(a)}$ and respectively $F' := b'/G_1^{(b)} \cdots G_{i-1}^{(b)} = b_{\gamma+1} \cdots b_\beta \cdot \prod_{j=i+1}^s G_j^{(b)}$. \square

Theorem 1. *At the end of the i -th iteration of the while loop (Line 39) of Algorithm 5, the output lists $\mathcal{C}, \mathcal{T}, \mathcal{D}$ are currently filled with:*

$$\mathcal{C} = [g_1, \dots, g_i], \quad \mathcal{T} = [p^{e_1}, \dots, p^{e_i}], \quad \mathcal{D} = [G_1, \dots, G_{i-1}],$$

and at the end of the algorithm, after say s iterations, the sequence $[(g_1, p^{e_1}), \dots, (g_s, p^{e_s})]$ is a gcd chain.

Proof. We must show that Isomorphisms (4) and (5) hold. By the Chinese remaindering theorem it suffices to show that

$$\langle a, b, p^e \rangle = \langle G_1, p^{e_1} \rangle \cdots \langle G_{s-1}, p^{e_{s-1}} \rangle \langle g_s, p^{e_s} \rangle. \quad (7)$$

holds, in order to prove (4); and to prove that (5) holds, it suffices to show that:

$$\text{for all } s \geq i \geq 1, \quad \langle a, b, p^{e_i} \rangle = \langle G_1, p^{e_1} \rangle \cdots \langle G_{i-1}, p^{e_{i-1}} \rangle \langle g_i, p^{e_i} \rangle, \quad (8)$$

Assume first that $s = 1$. Then Eq. (8) and Eq. (7) corresponds respectively to the first and to the second equality of Lemma 6.

Assume now that $s > 1$. Let us prove Eq. (8) first. The case $i = 1$ amounts to $\langle g_1, p^{e_1} \rangle = \langle a, b, p^{e_1} \rangle$ and is provided by Lemma 6. Assuming now $i > 1$. Statements 1.-2 of Lemma 7 together insure that $\langle a, b, p^{e_i} \rangle \subset \langle G_1, p^{e_1} \rangle \cdots \langle G_{i-1}, p^{e_{i-1}} \rangle \langle g_i, p^{e_i} \rangle$ (†).

On the other hand, Corollary 1 insures of the inclusion:

$$\langle G_1, p^{e_1} \rangle \cdots \langle G_{i-1}, p^{e_{i-1}} \rangle \langle g_i \rangle \subset \langle a, b, p^e \rangle.$$

Indeed the generators of the product ideals on the l.h.s. are proved in that corollary to belong to the r.h.s. It follows that:

$$\langle G_1, p^{e_1} \rangle \cdots \langle G_{i-1}, p^{e_{i-1}} \rangle \langle g_i, p^{e_i} \rangle \subset \langle a, b, p^{e_i} \rangle.$$

Together with (†), this proves Eq. (8). Let us prove Eq. (7). Eq. (8) with $i = s$ yields:

$$\langle G_1, p^{e_1} \rangle \cdots \langle G_{s-1}, p^{e_{s-1}} \rangle \langle g_s, p^{e_s} \rangle = \langle a, b, p^{e_s} \rangle.$$

Corollary 1 provides the inclusion:

$$\langle a, b, p^e \rangle \supset \langle G_1, p^{e_1} \rangle \cdots \langle G_{s-1}, p^{e_{s-1}} \rangle \langle g_s \rangle.$$

Moreover, since by assumption there are s iterations, the while loop at Line 39 of Algorithm 5 stops at the $s + 1$ -th iteration. Either $B_s \neq \text{“end”}$ and then $B_s = 1$. Proposition 1 then guarantees that $p^{e_s} B_s = p^{e_s} \in \langle a, b, p^e \rangle$. We obtain the equality (7) required. Either $B_s = \text{“end”}$: the $s - 1$ -th call to “largestFactor” at Line 40 with input $g_{s-1}, B_{s-1}, p^{e - e_{s-1}}$ exits at one of the Lines 25, 27, 31. If it is at Line 25, then $B_{s-1} = 0$ which happens eventually only if $b = 0$, that is if there is only one iteration; Since $s > 1$ here, this cannot happen. If it is at Line 27, then $B_{s-1} = 1$, and there is no such s -th iteration since the test at Line 39 is not passed. It remains the case of Line 31. Then the output T is equal here to p^{e_s} and we have $e = e_s$. Then Eq. (7) and Eq. (8) coincide. □

4 A variant of Weierstrass preparation’s theorem

The Weierstrass preparation theorem states that a formal power series $f = \sum_i a_i X^i \in \mathfrak{a}[[X]]$ with coefficients in a local complete ring $(\mathfrak{a}, \mathfrak{m})$, not all of them lying in \mathfrak{m} , has a unique factorization $f = qu$ where $q = q_0 + \cdots + q_{n-1} X^{n-1} + X^n$ is monic and $q_i \in \mathfrak{m}$, and where $u \in \mathfrak{a}[[x]]^*$ is an invertible power series.

In our context the local complete ring is $\mathfrak{a} = k[x]/\langle p^e \rangle$, $\mathfrak{m} = \langle p \rangle$ (indeed it is equal to $k[[x]]/\langle p^e \rangle$, which is a finite quotient of the local complete ring $k[[x]]$). But the factorization supplied by the classical version (*e.g.* [8, Theorem 9.1]) does not fit the needs of this work. The following variant does:

Proposition 2. *Let $(\mathfrak{a}, \mathfrak{m})$ be a complete local ring and let $f \in \mathfrak{a}[X]$ a polynomial, say $f = f_d X^d + \cdots + f_0$ which has not all of its coefficients f_i lying in \mathfrak{m} . Write f_k the coefficient of highest degree that is not in \mathfrak{m} . There are unique polynomials q and u in $\mathfrak{a}[X]$ such that $f = uq$ where:*

- q is monic of degree k .
- $u = u_0 + u_1 X + \cdots + u_{d-k} X^{d-k}$ where $u_0 \notin \mathfrak{m}$ and $u_i \in \mathfrak{m}$ for $i \geq 1$ (note that u is a unit of $\mathfrak{a}[[X]]$).

The proof is adapted from [8, Theorem 9.1-9.2].

Proof. Write $rev_d(f) := X^d f(\frac{1}{X})$ the reversal polynomial of f (this is why we need f to be a polynomial and not a general power series). The term of smallest degree not in \mathfrak{m} is $f_k X^{d-k}$. By the standard Weierstrass preparation Theorem 9.2 of [8] $rev_d(f) = gs$ where s is an invertible power series in $\mathfrak{a}[[X]]$ and $g = X^{d-k} + g_{d-k-1} X^{d-k-1} + \cdots + g_0$ is a monic polynomial which satisfies $g_i \in \mathfrak{m}$. Since $\deg(rev_d(f)) = d$ and that g is monic, necessarily $s \in \mathfrak{a}[X]$ and is of degree k . We write $s = s_0 + \cdots + s_k X^k$. Thus:

$$rev_d(rev_d(f)) = f = rev_{d-k}(g) rev_k(s) = (1 + g_{d-k-1} X + \cdots + g_0 X^{d-k})(s_k + \cdots + s_0 X^k)$$

Now since s is invertible, $s_0 \in \mathfrak{a}^*$. Letting $u = s_0 + \cdots + g_0 s_0 X^{d-k}$ and $q = s_0^{-1} s_k + \cdots + s_0^{-1} s_1 X^{k-1} + X^k$ provides polynomials satisfying $f = uq$ as well as the requirement of the proposition. □

The polynomial q can be computed by an Euclidean division. This possibility is a minor adaptation of [8, Theorem 9.1], that we have reproduced in Appendix for sake of completeness.

Corollary 2. *With the same notations and hypotheses of Proposition 2, given the inverse coefficient f_k^{-1} , the monic polynomial q can be computed by mimicking the Euclidean division of X^k by $f_k X^k + f_{k-1} X^{k-1} + \dots$, but with f instead of $f_k X^k + f_{k-1} X^{k-1} + \dots$ to obtain $X^k = gf + r$. The power series g needs to be computed only up to modulo X^{d-k+1} . Followed by the inversion of the truncated power series g to get: $q = (X^k - r)g^{-1}$.*

Proof. By Lemma 8, there exists a unique invertible power series $g \in \mathfrak{a}[[X]]$ and a unique polynomial $r \in \mathfrak{a}[X]$ such that $X^k = gf + r$. Therefore $(X^k - r)g^{-1} = f$ is the factorization of Proposition 2, by uniqueness. We deduce that g^{-1} is a polynomial of degree $d-k$ that verifies the conditions of the aforementioned proposition. In consequence, the power series g needs to be known only modulo X^{d-k+1} . \square

Example 4 *Let $\mathfrak{a} = k[[x]]/\langle x^2 \rangle \simeq k[x]/\langle x^2 \rangle$ be the complete local ring of maximal ideal $\mathfrak{m} = \langle x \rangle$, $f = xy^2 + y + 1$. The Weierstrass factorization of Proposition 2 insures the existence of a polynomial $q = y + \dots \in \mathfrak{a}[y]$ and $u = u_0 + u_1 y$ with $u_0 \in \mathfrak{a}^*$ and $u_1 \in \mathfrak{m}$, such that $f = uq$. To compute it, Euclidean division as explained in Corollary 2 works as follows (boxed terms are parts of the remainder).*

$$\begin{array}{r|l} y & xy^2 + y + 1 \\ -xy^2 & \boxed{-1} \quad 1 - xy + x \\ \hline xy & \boxed{-x} \end{array}$$

We obtain $y = (xy^2 + y + 1)(1 - xy + x) - x - 1$, hence $qu = (y + x + 1)(1 - xy + x)^{-1} = xy^2 + y + 1$ is the factorization of Proposition 2. Now $u = (1 - xy + x)^{-1} = 1 + \sum_{i \geq 1} x^i (y - 1)^i = 1 + x(y - 1)$ in $(k[x]/\langle x^2 \rangle)[y]$. We check that $u_0 = 1 - x \in \mathfrak{a}^*$ and $u_1 = x \in \mathfrak{m}$.

5 Concluding remarks

A running example Let $p(x) = x$, $e = 7$, that is the input primary triangular set in one variable is $T(x) = x^7$. The two input monic polynomials in $(\mathbb{Q}[x]/\langle T(x) \rangle)[y]$ are defined through their unique factorization into (two) coprime factors as:

$$\begin{aligned} a &:= (y + x + x^2 + x^3 + x^4 + x^5 + x^6) * (y + 1 + x^3 + x^6) \\ b &:= (y + x + x^2 + 2 * x^3 + x^4 + x^5 + x^6) * (y + 1 + x^3 + x^5) \end{aligned}$$

`gcdChain(a, b, x7) ==`

– First call to “largestFactor” Line 39 ($i = 0$): `largestFactor(a, b, x7) ==`

• Line 28 `modifiedSubres(a, b, x7) ==`

* Line 10: Subresultant mod x^7 does not fail; We obtain $\mathbf{S} = [a, b, \dots]$, precisely:

$$\begin{aligned} \mathbf{S} &= [y^2 + (2 * x^6 + x^5 + x^4 + 2 * x^3 + x^2 + x + 1) * y + 2 * x^6 + 2 * x^5 + 2 * x^4 + x^3 + x^2 + x, \\ & y^2 + (x^6 + 2 * x^5 + x^4 + 3 * x^3 + x^2 + x + 1) * y + 4 * x^6 + 2 * x^5 + 2 * x^4 + 2 * x^3 + x^2 + x, \\ & (-x^6 + x^5 + x^3) * y + 2 * x^6 + x^3, \\ & 0] \end{aligned}$$

* The last non nilpotent subresultant is $b = \mathbf{S}[2]$ and the first nilpotent is $\mathbf{S}[3] = (-x^6 + x^5 + x^3) * y + 2 * x^6 + x^3$.

* While loop Line 20 ($j = 3$) $S_{r_j} = \mathbf{S}[3]$

$$\text{nilpotentFactor}(S_{r_j}, x^7) == x^3, -1 \quad (P_{new} = x^3, x^3 \mid \mathbf{S}[3], i_{new} = -1)$$

* While loop Line 20 ($j = 2$) $S_{r_j} = \mathbf{S}[2] = b$

$$\text{nilpotentFactor}(S_{r_j}, x^7) == 1, 2 \quad (P_{new} = 1, \text{coeff}(b, i_{new}) = \text{coeff}(b, y^2) = 1)$$

- * exit while loop and return $\text{modifiedSubres}(a, b, T) == \mathbf{S}, 2, 2, x^3$
- Line 29: $\text{WeierstrassMonic}(b, x^7, 2) == b$
- Line 32: $S = \mathbf{S}[3]/x^3 = (-x^3 + x^2 + 1) * y + 2 * x^3 + 1$
- Lines 33-34: $p^\ell = T/x^3 = x^4$, $\text{nilpotentFactor}(S, x^4) == 1, 1$ ($i = 1$)
- Line 35: $B_1 = y + 3 * x^3 - x^2 + 1$
- return b, x^3, B_1, x^4
- Line 42, Line 52: $\mathcal{T} = [x^3]$, $\mathcal{C} = [g_1]$ ($g_1 = b \bmod \langle x^3 \rangle$)
- Line 39 ($i = 1$): $\text{largestFactor}(g_1, B_1, x^4) ==$
 - Line 28: $\text{modifiedSubres}(g_1, B_1, x^4) ==$
 - * Line 10 Subresultant mod x^4 does not fail; We obtain

$$\mathbf{S} = [y^2 + (3 * x^3 + x^2 + x + 1) * y + 2 * x^3 + x^2 + x, \\ y + 3 * x^3 - x^2 + 1, \\ 3 * x^3 - x^2, \\ 0]$$

- * The last non nilpotent is actually $B_1 = \mathbf{S}[2]$ and the first one is $\mathbf{S}[3] = 3 * x^3 - x^2$.
- * While loop Line 20 ($j = 3$) $S_{r_j} = \mathbf{S}[3]$
 $\text{nilpotentFactor}(S_{r_j}, x^4) == x^2, -1$ ($P_{new} = x^2$, $x^2 \mid \mathbf{S}[3]$, $i_{new} = -1$)
- * While loop Line 20 ($j = 2$) $S_{r_j} = \mathbf{S}[2] = B_1$
 $\text{nilpotentFactor}(S_{r_j}, x^4) == 1, 1$ ($P_{new} = 1$, $\text{coeff}(B_1, i_{new}) = \text{coeff}(B_1, y) = 1$)
- * exit while loop and return $\text{modifiedSubres}(g_1, B_1, x^4) == \mathbf{S}, 2, 1, x^2$
- Line 29: $\text{WeierstrassMonic}(B_1, x^4, 1) == B_1$
- Lines 32 $S = \mathbf{S}[3]/x^2 = 3 * x - 1$
- Lines 33-34: $p^\ell = x^4/x^2 = x^2$, $\text{nilpotentFactor}(S, x^2) == 1, 0$ ($i = 0$)
- Line 35: $B_2 = 3 * x - 1$
- return $B_1 (= g_2), x^2 (= T'_2), 3 * x - 1 (= B_2), x^2 (= S_2)$
- (Hensel Lifting, preparation) Line 44: $G_1 = g_1/g_2 = b/B = y + x \bmod \langle x^2 \rangle$
- Lines 45-47 $\alpha = -(1 + x)$, $\beta = (1 + x)$, $T_2 = x^3 \cdot x^2 = x^5$, $\text{precision} = 5$.
- (Hensel Lifting): Step 1: from x^2 to x^4 :

$$G_1^* = y + 2 * x^3 + x^2 + x, \quad g_2^* = y + x^3 + 1$$

Step2: from x^4 to x^8 :

$$G_1^* = y - 2 * x^7 + x^6 + x^5 + x^4 + 2 * x^3 + x^2 + x, \quad g_2^* = y + 2 * x^7 + x^5 + x^3 + 1$$

- Line 49: $g_2 = g_2^* \bmod \langle x^5 \rangle = y + x^3 + 1$
- Lines 50-52: $\mathcal{D} = [G_1^* \bmod \langle x^3 \rangle] = [y + x^2 + x]$, $\mathcal{T} = [x^3, x^5]$, $\mathcal{C} = [b, y + x^3 + 1]$
- Line 39 ($i = 2$): $\text{largestFactor}(g_2, 3 * x - 1, x^5) ==$
 - Line 27 $\text{deg}_y(3 * x - 1) = 0$, return $3 * x - 1, x^6$, “end”.
- return $\mathcal{C} = [b, y + x^3 + 1]$, $\mathcal{D} = [y + x^2 + x]$, $\mathcal{T} = [x^3, x^5]$.

According to Definition 2, we have here $s = 2$ blocks. And moreover $G_1 = \mathcal{D}[1]$ and $G_2 = g_2 = \mathcal{C}[2]$, yielding the isomorphism:

$$(\mathbb{Q}[x]/\langle x^7 \rangle)[y]/\langle a, b \rangle \simeq (\mathbb{Q}[x]/\langle x^3 \rangle)[y]/\langle y + x^2 + x \rangle \times (\mathbb{Q}[x]/\langle x^5 \rangle)[y]/\langle y + x^3 + 1 \rangle$$

In this example, this decomposition coincides with the primary decomposition.

Generalization All steps of Algorithm 4 “largestFactor” extends to more than one variable, except the management of the first nilpotent subresultant. To illustrate this difficulty, let us consider a primary triangular set $T = (x^2, y^2)$ of radical (x, y) , and some input polynomials a and b in $(k[x, y]/\langle T \rangle)[z]$. It may happen that a subresultant is equal to say $xz + y$ (for example $a = z^2 + 2xz - y$ and $b = z^2 + xz - 2y$). It is nilpotent and the iterated resultant criterion allows to detect it. But there is no way to “remove” the nilpotent part as in the case of a polynomial of one variable. To apply Weierstrass preparation theorem, the polynomial must not be nilpotent. A solution consists of “adding” this polynomial to the coefficient ring. How this integration shall be done requires more work and will be investigated in the future.

Complexity The running-time of the algorithm is dominated by that of the subresultant calls. Indeed, all other subroutines are indeed based on classical algorithms which have a lower cost: Hensel lifting, inversion of a truncated power series are endowed of fast algorithms; And Weierstrass factorization is reduced to an Euclidean division. The standard subresultant p.r.s. have a quadratic (operations in $k[x]/\langle p^e \rangle$) cost in the degrees of the input polynomials. Therefore a running-time of $O((e \deg(p))^2 \cdot \deg(a)^2)$ can be estimated using naive but realistic algorithms..

We speculate that a fast version of our algorithm has a quasi-linear cost $\tilde{O}(e \deg(p) \cdot \deg(a))$, by using fast “divide and conquer” (a.k.a half-gcd). This is quite challenging, and as one knows, is not reflected in any practical algorithm. For now, we care on *feasibility* with a view toward extensions to several variables.

References

1. Parisa Alvandi, Marc Moreno Maza, Éric Schost, and Paul Vrbik. *A Standard Basis Free Algorithm for Computing the Tangent Cones of a Space Curve*, pages 45–60. Springer International Publishing, Cham, 2015.
2. C. Chen and M. Moreno-Maza. Algorithms for computing triangular decomposition of polynomial systems. *Journal of Symbolic Computation*, 47(6):610–642, 2012.
3. Jin-San Cheng and Xiao-Shan Gao. Multiplicity-preserving triangular set decomposition of two polynomials. *Journal of Systems Science and Complexity*, 27(6):1320–1344, 2014.
4. Xavier Dahan. Gcd modulo a primary triangular set of dimension zero. In *Proceedings of the 2017 ACM on International Symposium on Symbolic and Algebraic Computation*, ISSAC '17, pages 109–116, New York, NY, USA, 2017. ACM.
5. M. Moreno Maza F. Lemaire and Y. Xie. The `RegularChains` library.
6. É. Hubert. Notes on triangular sets and triangulation-decomposition algorithms. I. Polynomial systems. volume 2630 of *Lecture Notes in Comput. Sci.*, pages 1–39. Springer, Berlin, 2003.
7. M. Kalkbrenner. A generalized Euclidean algorithm for computing triangular representations of algebraic varieties. *J. Symbolic Comput.*, 15(2):143–167, 1993.
8. Serge Lang. *Algebra (revised third edition)*, volume 211 of *Graduate Texts in Mathematics*. Springer Science and Media, 2002.
9. B.-H Li. A method to solve algebraic equations up to multiplicities via ritt-wu’s characteristic sets. *Acta Analysis Functionalis Applicata*, 2:97–109, 2003.
10. X. Li, M. Moreno-Maza, and W. Pan. Computations modulo regular chains. In *Proceedings of the 2009 international symposium on Symbolic and algebraic computation*, pages 239–246. ACM, 2009.
11. X. Li, C. Mou, and D. Wang. Decomposing polynomial sets into simple sets over finite fields: The zero-dimensional case. *Computers & Mathematics with Applications*, 60(11):2983 – 2997, 2010.
12. Steffen Marcus, Marc Moreno Maza, and Paul Vrbik. *On Fulton’s Algorithm for Computing Intersection Multiplicities*, pages 198–211. Springer Berlin Heidelberg, Berlin, Heidelberg, 2012.
13. M. Moreno Maza and R. Rioboo. Polynomial gcd computations over towers of algebraic extensions. In *Proc. AAECC-11*, pages 365–382. Springer, 1995.
14. Guillaume Moroz and Éric Schost. A fast algorithm for computing the truncated resultant. In *Proceedings of the ACM on International Symposium on Symbolic and Algebraic Computation*, pages 341–348. ACM, 2016.
15. Chenqi Mou, Dongming Wang, and Xiaoliang Li. Decomposing polynomial sets into simple sets over finite fields: The positive-dimensional case. *Theoretical Computer Science*, 468:102–113, 2013.
16. J. F. Ritt. Differential equations from an algebraic standpoint. *Colloquium publications of the AMS*, 14, 1932.
17. J. von zur Gathen and J. Gerhard. *Modern computer algebra*. Cambridge University Press, New York, NY, USA, 2003. Second Edition.
18. D. Wang. *Elimination Methods*. Texts & Monographs in Symbolic Computation. Springer Vienna, 2012.
19. W. T. Wu. On zeros of algebraic equations - an application of Ritt principle. *Kexue Tongbao*, 5:1–5, 1986.

Appendix

This is a modification of Lang’s proof [8, Theorem 9.1] up to a minor point indicated below, of the Weierstrass division, which is used in Corollary 2.

Lemma 8 *With the same notations and assumptions of Proposition 2, given $g \in \mathfrak{a}[[X]]$, we can solve the equation $g = qf + r$ uniquely with $q \in \mathfrak{a}[[X]]$ and $r \in \mathfrak{a}[X]$ and $\deg r < k$.*

Proof. Define the linear maps $\alpha : \mathfrak{a}[[X]] \rightarrow \mathfrak{a}[X]$, $\sum_i a_i X^i \mapsto \sum_{i=0}^{k-1} a_i X^i$ and $\tau : \mathfrak{a}[[X]] \rightarrow \mathfrak{a}[[X]]$, $\sum_i a_i X^i \mapsto \sum_i a_{k+i} X^i$. It is clear that $b \in \mathfrak{a}[X]$ has degree $< k$ if and only if $\tau(b) = 0$ or $\alpha(b) = b$. Therefore if $g = qf + r$, $\deg(r) < k$ then $\tau(g) = \tau(qf)$. Moreover $\tau(X^k h) = h$ and $h = \tau(h)X^k + \alpha(h)$ for any $h \in \mathfrak{a}[[X]]$. Thus

$$\tau(g) = \tau(q(\alpha(f) + X^k \tau(f))) = \tau(q\alpha(f)) + \tau(f)q.$$

Let $Z = \tau(f)q$. Notice that $\tau(f) = f_k + X m'$ with $f_k \in \mathfrak{a}^*$ and $m' \in \mathfrak{ma}[[X]]$ by assumption, hence $\tau(f)$ is invertible in $\mathfrak{a}[[X]]$. The equation above can be rewritten: $\tau(g) = \tau(Z \frac{\alpha(f)}{\tau(f)}) + Z$. Being able to solve this equation in Z uniquely gives $\tau(f)q$, hence q , hence $r = g - f q$ in a unique way.

To do it, the proof differs slightly from that of [8, Theorem 9.2] to the following point. The reason why the image of the map below

$$\tau \circ \frac{\alpha(f)}{\tau(f)} : \mathfrak{a}[[X]] \rightarrow \mathfrak{ma}[[X]]$$

is $\mathfrak{ma}[[X]]$ is because $\tau(f)^{-1} = f_k^{-1} \sum_{\ell \geq 0} (-1)^\ell (mX)^\ell$ for an $m \in \mathfrak{ma}[[X]]$ and hence $\alpha(f)\tau(f)^{-1} = \alpha(f)f_k^{-1} + \alpha(f)M$ where $M \in \mathfrak{ma}[[X]]$. Therefore $\tau(\alpha(f)\tau(f)^{-1}) \in \mathfrak{ma}[[X]]$ since $\deg(\alpha(f)f_{k-1}) = \deg(\alpha(f)) \leq k-1$. Thus for any power series $h \in \mathfrak{a}[[X]]$,

$$\left(\tau \circ \frac{\alpha(f)}{\tau(f)} \right) (h) \in \mathfrak{ma}[[X]].$$

Now $\mathfrak{a}[[X]]$ being a complete ring, $I + \tau \circ \frac{\alpha(f)}{\tau(f)}$ is invertible and $Z = (I + \tau \circ \frac{\alpha(f)}{\tau(f)})^{-1}(\tau(g))$ is determined in a unique way. \square