

The Harmony of Spheres

Otfried Cheong*

Let $\mathcal{F} = \{X_1, \dots, X_n\}$ be a family of disjoint compact convex sets in \mathbb{R}^d . An oriented straight line ℓ that intersects every X_i is called a (line) *transversal* to \mathcal{F} . A transversal naturally induces an ordering of the elements of \mathcal{F} , this ordering, together with its reverse (which is induced by the same line with opposite orientation), is called a *geometric permutation*.

Geometric transversal theory studies properties of line transversals, and of families of objects admitting line transversals—see the handbook chapter by Wenger for a survey of this area [6]. Topics of interest are a characterization of the geometric permutations that a given family of objects can admit, and Helly-type theorems of the form: If every subfamily of \mathcal{F} of size at most k admits a line transversal, then \mathcal{F} must admit a line transversal.

In this talk we discuss recent results on transversals to disjoint *spheres* (balls) in three and more dimensions.

Katchalski et al. [5] had proven in 2003 that a family of sufficiently many disjoint unit spheres admits at most four different geometric permutations (for arbitrary disjoint spheres, the bound is $\Theta(n^{d-1})$). By showing that it is impossible for four spheres to realize the two geometric permutations ABCD and BADC at the same time, we were able to improve this, showing that $n \geq 9$ disjoint unit spheres admit at most two geometric permutations, which differ in the swapping of two adjacent spheres only [3]. (For $4 \leq n \leq 8$, it is still open whether the correct answer is two or three.)

The first Helly-type theorem for line transversals to spheres was proven by Holmsen et al. [4]. Their result is based on the following fact: Given a sequence of disjoint spheres in \mathbb{R}^d , and consider all oriented lines intersecting all spheres in the given order. Then the set of orientations of these lines is a convex set (on the sphere of directions).

Holmsen et al. proved this for disjoint unit spheres in \mathbb{R}^3 . We were able to generalize the result to unit spheres in arbitrary dimensions [2], and very recently Borcea et al. could generalize it to families of arbitrary disjoint spheres using algebraic techniques [1].

The convexity of the cone of directions appears to be a distinguishing feature of spheres—it does not even generalize to convex objects that are perfectly smooth and arbitrarily fat. It even appears that the convexity

fails to hold if the spheres are allowed to intersect ever so slightly.

References

- [1] C. Borcea, X. Goaoc, and S. Petitjean. Line transversals to disjoint balls. In *Proc. 23rd Annual Symposium on Computational Geometry 2007*, pages 245–254.
- [2] O. Cheong, X. Goaoc, A. Holmsen, and S. Petitjean. Helly-type theorems for line transversals to disjoint unit balls. To appear in *Discrete Comput. Geom.*
- [3] O. Cheong, X. Goaoc, and H.-S. Na. Geometric permutations of disjoint unit spheres. *Computational Geometry: Theory and Applications* **30** (2005) 253–270.
- [4] A. Holmsen, M. Katchalski, and T. Lewis. A Helly-type theorem for line transversals to disjoint unit balls. *Discrete Comput. Geom.* **29** (2003), 197–206.
- [5] M. Katchalski, S. Suri, and Y. Zhou. A constant bound for geometric permutations of disjoint unit balls. *Discrete Comput. Geom.* **29** (2003) 161–173.
- [6] R. Wenger. Helly-type theorems and geometric transversals. In J. E. Goodman and J. O’Rourke, editors, *Handbook of Discrete and Computational Geometry*, chapter 4, pages 73–96. CRC Press LLC, Boca Raton, FL, 2nd edition, 2004.

*Department of Computer Science, Korea Advanced Institute of Science and Technology, otfried@kaist.edu