

UCSF

UC San Francisco Previously Published Works

Title

Crowdsourced estimation of cognitive decline and resilience in Alzheimer's disease

Permalink

<https://escholarship.org/uc/item/7x89j34x>

Journal

Alzheimer's & Dementia, 12(6)

ISSN

1552-5260

Authors

Allen, Genevera I

Amoroso, Nicola

Anghel, Catalina

et al.

Publication Date

2016-06-01

DOI

10.1016/j.jalz.2016.02.006

Peer reviewed



Published in final edited form as:

Alzheimers Dement. 2016 June ; 12(6): 645–653. doi:10.1016/j.jalz.2016.02.006.

Crowdsourced estimation of cognitive decline and resilience in Alzheimer's disease

A full list of authors and affiliations appears at the end of the article.

Abstract

Identifying accurate biomarkers of cognitive decline is essential for advancing early diagnosis and prevention therapies in Alzheimer's Disease. The Alzheimer's Disease DREAM Challenge was designed as a computational crowdsourced project to benchmark the current state-of-the-art in predicting cognitive outcomes in Alzheimer's Disease based on high-dimensional, publicly available genetic and structural imaging data. This meta-analysis failed to identify a meaningful predictor developed from either data modality, suggesting that alternate approaches should be considered for prediction of cognitive performance.

2. Background

The Alzheimer's Disease DREAM Challenge (<http://dx.doi.org/10.7303/syn2290704>) was designed to provide an unbiased assessment of current capabilities for estimation of cognition and prediction of cognitive decline using genetic and imaging data from public data resources using a crowd-sourced approach. The ability to predict rate of cognitive decline – both prior to and following diagnosis – is essential to effective trial design for the development of therapies for Alzheimer's Disease (AD) prevention and treatment. Major collaborative efforts in the field are assessing the association of genetic loci with AD diagnosis and the application of structural imaging for development of early biomarkers of diagnosis, but the utility of these approaches to estimate cognition or predict cognitive decline is not well established. This project was designed under the advisement of a panel of experts in the field to evaluate whether these questions could be meaningfully addressed

Correspondence to: John SK Kauwe; Arno Klein; Lara M Mangravite; Mette A Peters.

*Data used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI contributed to the design and implementation of ADNI and/or provided data but did not participate in analysis or writing of this report. A complete listing of ADNI investigators can be found at: http://adni.loni.usc.edu/wp-content/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf

AUTHOR CONTRIBUTIONS: CJB, ECN, DWF, SHF, SSG, AK, JSKK, YK, BAL, LMM, TJM, TCN, MAP, GS, GV and NJT contributed to the challenge organization

VB, DD, PSH, RCG contributed with compute resources and scientific advice

GIA, NA, CA, DB, RB, KB, PCB, LC, CC, FC, YCC, BC, CYC, TYC, TC, SD, CD, JD, QD, JD, DD, RE, GE, EE, HF, JG, EG, YG, MYH, CH, JH, JI, PI, QJ, DK, RK, EL, ML, EL, XL, ZL, JL, AL, MM, AM, EM, JN, AN, MN, MN, RNN, YJO, YP, SP, SRP, PP, CP, VYS, PS, XS, AS, ST, AT, YAT, EV, GX, LX, YX, JX, HY, XZ, YZ, FZ, HZ and SZ participated in the challenge community phase

DAB, RJBD, SL, SJN, AS and MWW contributed with data used in the challenge

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

with current methodologies given existing public data sources. To ensure that these questions were tested across a broad spectrum of the latest analytical approaches, the study was designed as a crowdsourced, community-based challenge in which participants were invited to address one or more of the following three problems: (1) The prediction of cognitive decline over time based on genetic data. (2) The prediction of resilience to cognitive decline in individuals with elevated amyloid burden based on genetic data. (3) The estimation of cognitive state based on structural magnetic resonance (MR) imaging data.

3. Results

3.1 Study design and data harmonization

To ensure that predictors were detecting true biological variation rather than study-specific technical variation, this project required inclusion of data from multiple study sources. While genetic and imaging data have been generated within many rich longitudinal cohorts across the field, the procurement and harmonization of these data sets was a non-trivial problem that required solutions to overcome political, ethical, and technical barriers. For example, the generation of whole genome sequencing data across multiple AD cohorts within the NIH-funded AD sequencing project has resulted in a powerful resource for genetic analysis in the field but longitudinal information on cognitive traits is not readily available in those datasets. Despite limitations on data accessibility, multiple relevant data sources were identified and used in this project including: the Alzheimer's Disease Neuroimaging Initiative (ADNI)(1), the Rush Alzheimer's Disease Center Religious Orders Study(2) and Memory and Aging Project (MAP)(3) and the European AddNeuroMed(4) study, which is part of InnoMed, a precursor to the Innovative Medicines Initiative. Data selection and processing was performed based on data availability across these three datasets. As such, cognition was defined using Mini Mental State Examination (MMSE) scores(5), genetic data was provided based on imputation across array-based genotype data, and structural MR imaging data was reprocessed in each cohort using a common processing pipeline. Genetic and imaging data was supplemented with a limited set of covariates including diagnosis, initial MMSE score, age at the initial examination, years of education, gender, and *APOE* haplotype. Participants were provided with data from ADNI to train algorithms over a four-month period and, to ensure that participation was not limited by access to compute resources, they were offered use of the IBM z-Enterprise cloud to perform analyses. The challenge generated significant interest with 527 individuals from around the world registered to participate. A leaderboard displayed accuracy of submissions throughout the duration of the challenge: 1,157 submissions were made for problem 1, 1,478 submissions for problem 2, and 434 submissions for problem 3. Thirty-two teams submitted final results that were scored based on prediction/estimation of blinded outcomes within ROS/MAP for genetic predictions and AddNeuroMed for imaging-based estimations (Figure 1).

3.2 Genetic prediction of cognitive decline

The first challenge question assessed the ability of current methods to predict change in cognitive examination performance based on genetic data. High prediction accuracy would signal the potential for noninvasive biomarkers of cognition to have a major clinical impact on early AD diagnosis and prevention. Previous efforts to develop predictors of change in

cognitive function have not succeeded in providing robust and replicable models(6–8). Genetic variation has been demonstrated to influence AD status: rare genetic mutations at several loci are implicated in familial forms of early-onset disease(9) while common variation contributes 33% to variance in sporadic AD and 22 loci have been implicated by large-scale genetic association analyses(10, 11). However, with the exception of the *APOE4* haplotype, there has been little success in transforming these genetic associations into meaningful clinical predictions of cognitive decline. For this purpose, participants were challenged to predict 2-year changes in MMSE scores based on genotypes imputed from SNP array data. Participants trained their algorithms with 767 ADNI samples and the algorithms' predictions were evaluated on a test set of 1,175 ROS/MAP samples with blinded outcome measures. The algorithm with the best predictive performance at the midpoint of the challenge did not contain any genetic features beyond *APOE* haplotype. Since the goal of this subchallenge was to assess genetic contribution to prediction of cognitive decline, this top-ranked algorithm was openly shared across teams as an interim baseline upon which to incorporate additional genetic predictors (<http://dx.doi.org/10.7303/syn2838779>). Eighteen teams submitted final predictions. The majority of methods performed significantly better than a permutation-based random model prediction (Figure 2a). A cluster of six methods performed significantly better than the others (including the interim baseline model) but were statistically indistinguishable amongst themselves (Figure 2d). Of these, the prediction with the best overall score (team GuanLab_umich from the University of Michigan) achieved a Pearson correlation of 0.382 and a Spearman correlation of 0.433 (the overall score was a rank-based combination of these two measures of performance; see online Supplement and Supplementary Methods: <http://dx.doi.org/10.7303/syn3383106>). However, no significant contribution of genetics beyond *APOE* haplotype to predictive performance was observed across any of the submissions. Given the small sample size, no conclusions can be inferred from this analysis regarding the existence of genetic loci associated with cognitive decline. Rather, these observations suggest that predictors of cognitive decline developed based on genetic data will not be useful within the clinical setting.

3.3 Genetic prediction of cognitive resilience

The second question challenged participants to identify genetic predictors that could distinguish individuals who exhibit resilience to AD pathology as defined by minimal change in cognitive function despite evidence of amyloid deposition(12, 13). Identification of genetic signatures predictive of cognitive resilience would aid in the elucidation of mechanisms that may confer resilience, providing a powerful tool to help advance AD prevention strategies and treatment development. Eleven teams submitted predictions of resilience based on a training set derived from 176 ADNI subjects. Evaluations were made using data derived from 257 individuals from the ROS/MAP data. Despite using the largest such public dataset assembled to date, participants were unable to develop algorithms with predictive performances significantly better than random (see Figure 2b, online Supplement and Supplementary Methods in Synapse: <http://dx.doi.org/10.7303/syn3383106>). While it is likely that the study was underpowered due to small sample size and trait heterogeneity, this result suggest that information about cognitive resilience is not easily discoverable from SNP analysis.

3.4 Structural imaging-based estimation of cognition

The third question challenged participants to estimate cognitive state using structural brain image data (Figure 1, lower panel). Brain imaging has emerged as a powerful method for monitoring neurodegeneration and there is great enthusiasm in the field to make use of images for diagnosis and prediction. There have been many attempts in the past to correlate changes in brain shape with disease progression and/or diagnosis, conventionally using measures of volume for a given brain region(14, 15). More detailed shape measures of image features including cortical thickness, curvature, and depth have also been found to be relevant to a variety of neurological conditions(16). Participants were challenged to estimate MMSE scores based on structural brain images, or shape information derived from these images. Participants trained algorithms using ADNI data (N=628) and were evaluated using AddNeuromed data (N=182) for which they were blind to outcome measures. To engage as many participants as possible from both within and beyond the neuroimaging community, the data were provided both as raw MR images and as tables containing shape measures (volume, thickness, area, curvature, depth, etc.) for every labeled brain region. Thirteen teams submitted estimates for final evaluation and all teams performed better than a random model (see online Supplement and Supplementary Methods in Synapse: <http://dx.doi.org/10.7303/syn3383106>). Three teams performed significantly better than the others (teams GuanLab_umich from the University of Michigan, ADDT from the Karolinska Institute and Pythia from the University of Pennsylvania) (Figure 2c) but were statistically indistinguishable from one another and tied for top average rank (Figure 2e). The algorithm that generated the best absolute mean combined rank (Team GuanLab_umich) achieved a concordance correlation coefficient of 0.569 and Pearson's correlation of 0.573 (the overall score was a rank-based combination of these two measures of performance). The most common features that contributed heavily to the MMSE estimates across the algorithms were hippocampal volume and entorhinal thickness, corroborating prior work(17–19). The top three teams also found that inclusion of shape measures of the entorhinal cortex (volume, curvature, surface area, travel and geodesic depth) improved overall estimation. Other features that contributed to predictions within the top three teams' results included volume of inferior lateral ventricle and amygdala (see online Supplement and Supplementary Methods in Synapse: <http://dx.doi.org/10.7303/syn3383106>). These results validate an established relationship between structural imaging data and cognition. However, the correlative performance of these estimators was low suggesting that their application in the clinical setting may not be sufficient to inform patient care.

4. Discussion

The AD DREAM Challenge provided a formalized assessment of the ability to develop meaningful predictions of cognitive performance from public genetic or imaging data using contemporary state-of-the-art predictive algorithms. Predictive performance across all three of the subchallenges was modest and most methods performed roughly equivalently. Given this uniform performance, we do not expect that the presented results are a failure of current modeling methodologies. A more likely explanation is that the data used to address these questions were inadequate to support these tasks. We also note that the majority of research teams that participated in this challenge did not have expertise in the field of AD. Although

the few teams that did possess this knowledge did not do better than the others, there remains the possibility that performance would have been improved by the inclusion of more domain experts.

4.1 Use of genetic information for cognitive prediction

The modest performance observed in the subchallenges focused on genetic analysis demonstrated that contemporary algorithms were not able to leverage genetic signal to make useful predictions for cognition. These results support the prevailing expectation that genetic variants of moderate to high frequency will not support viable biomarker development in AD (9–11). Although heritability estimates and linkage studies have demonstrated that there is a considerable estimated genetic contribution to AD onset and progression (11, 20, 21), evidence both within the AD field and across other complex disease (22) traits has indicated that this overall genetic contribution is the aggregated compilation of a large number of loci with small – independent or epistatic – effects. Historically, this type of signal is difficult to capture in predictive models and unlikely to be useful in a diagnostic setting (23). Furthermore, cognition is highly influenced by a host of non-genetic factors relating to lifestyle choices and accumulated exposures that were not represented across all of these datasets and, in fact, are not fully captured in most cohorts (24–27). Non-genetic contributions to cognitive performance may themselves provide an important base for successful predictions. Within the context of genetic analysis, the absence of these factors from models confounds the ability to detect real genetic signal and impacts the ability to accurately model state-specific genetic contributions. As such, future inquiry into the use of genetic testing for prediction of cognitive performance and AD risk assessment may be better served by focusing on the contribution of rare genetic variation. Recently discovered disease-associated rare variants have larger effect sizes than common variants and confer 2 to 5 fold greater risk or protection in carriers relative to the general population (28–30). Ongoing large-scale sequencing analyses will identify additional associated rare risk variants. In sufficient numbers, the aggregate prevalence would support the development of a genetic diagnostic containing a library of rare variants.

4.2 Use of structural imaging data for cognitive estimation

While the inexpensive and noninvasive nature of genetic testing makes this approach amenable to population-level disease screening, the resource-intensive nature of image-based testing is better positioned for careful evaluation of high-risk individuals. As such, these approaches are needed to provide a higher confidence estimate of cognitive performance. Although a variety of methods developed within the context of this challenge were able to successfully estimate cognition, none of these methods were sufficiently accurate to merit clinical consideration. These observations support previous work in the field (17, 19) and highlight the imperfect relationship between brain structure and function. Newer imaging modalities that focus on brain function and/or pathology – such as FDG-PET (31) or tau imaging (32) – may prove more successful for assessing cognitive dysfunction.

4.3 Effective performance of meta-analysis across diverse cohorts

A major consideration for any meta-analysis is the issue of appropriate harmonization of data across disparate sources. Despite leveraging several of the most deeply phenotyped cohorts in the field, this challenge limited analysis to those traits that were in common across cohorts. Although this approach to data harmonization is standard practice for meta-analyses (10), it greatly reduced the depth of the information available for modeling and influenced the selection of cognitive measures for use as prediction outcomes. Because each cohort had performed a battery of study-specific tests, this greatly limited the ability for finer grained assessment across cognitive processes. A more sensible approach for future analyses may be to focus effort on more sophisticated methods to calibrate disparate cognitive phenotypes across cohorts (33). Another undesirable consequence of the focus on traits measured in common was the inability to incorporate into model development the full spectrum of non-genetic and non-imaging factors that are known to influence cognitive performance (24–27). This suggests the need for development of alternate approaches for integrating heterogeneous data and/or assessing replication across cohorts. Alternatively, smaller scale analyses that prioritize phenotypic depth over sample size may afford a more refined view of disease.

In summary, this challenge demonstrated that predictions of cognitive performance developed from genetic or structural imaging data were modest across a diverse set of contemporary modeling methods. Future efforts to identify clinically relevant predictors of cognition will benefit from a focus on alternate data sources as well as methods that work to incorporate greater phenotypic complexity.

Authors

Genevera I Allen¹, Nicola Amoroso^{2,3}, Catalina Anghel⁴, Venkat Balagurusamy⁵, Christopher J Bare⁶, Derek Beaton⁷, Roberto Bellotti^{2,3}, David A Bennett⁸, Kevin Boehme⁹, Paul C Boutros^{4,53,54}, Laura Caberlotto¹⁰, Cristian Caloian⁴, Frederick Campbell¹, Elias Chaibub Neto⁶, Yu-Chuan Chang¹¹, Beibei Chen¹², Chien-Yu Chen¹³, Ting-Ying Chien¹⁴, Tim Clark^{15,16}, Sudeshna Das^{15,16}, Christos Davatzikos¹⁷, Jieyao Deng^{18,19}, Donna Dillenberger⁵, Richard JB Dobson^{20,21}, Qilin Dong^{18,19}, Jimit Doshi¹⁷, Denise Duma²², Rosangela Errico²³, Guray Erus¹⁷, Evan Everett¹, David W Fardo^{24,25}, Stephen H Friend⁶, Holger Fröhlich²⁶, Jessica Gan¹, Peter St George-Hyslop²⁷, Satrajit S Ghosh^{28,29}, Enrico Glaab³⁰, Robert C Green³¹, Yuanfang Guan^{32,33,34}, Ming-Yi Hong¹³, Chao Huang³⁵, Jinseub Hwang³⁷, Joseph Ibrahim³⁵, Paolo Inglese³⁸, Qijia Jiang¹, Yuriko Katsumata²⁵, John SK Kauwe⁹, Arno Klein⁶, Dehan Kong³⁵, Roland Krause³⁰, Emilie Lalonde⁴, Mario Lauria¹⁰, Eunjee Lee³⁵, Xihui Lin⁴, Zhandong Liu¹, Julie Livingstone⁴, Benjamin A Logsdon⁶, Simon Lovestone³⁹, Anandhi Lyappan^{40,26}, Michelle Ma¹², Ashutosh Malhotra^{40,26}, Lara M Mangravite⁶, Taylor J Maxwell⁴¹, Emily Merrill¹⁵, John Nagorski¹, Aishwarya Namasivayam³⁰, Manjari Narayan¹, Mufassra Naz^{40,26}, Stephen J Newhouse^{20,42}, Thea C Norman⁶, Ramil N Nurtdinov⁴³, Yen-Jen Oyang¹¹, Yudi Pawitan³⁶, Shengwen Peng^{18,19}, Mette A Peters⁶, Stephen R Piccolo⁴⁴, Paurush Praveen^{10,26}, Corrado Priami¹⁰, Veronica Y Sabelnykova⁴,

Philipp Senger⁴⁰, Xia Shen⁴⁵, Andrew Simmons²⁰, Aristeidis Sotiras¹⁷, Gustavo Stolovitzky^{46,5}, Sabina Tangaro³, Andrea Tateo², Yi-An Tung⁴⁷, Nicholas J Tustison⁴⁸, Erdem Varol¹⁷, George Vradsenburg⁴⁹, Michael W Weiner⁵⁰, Guanghua Xiao¹², Lei Xie⁵¹, Yang Xie¹², Jia Xu¹², Hojin Yang³⁵, Xiaowei Zhan¹², Yunyun Zhou¹², Fan Zhu³², Hongtu Zhu³⁵, Shanfeng Zhu^{18,19,52}, and Alzheimer's Disease Neuroimaging Initiative*

Affiliations

¹Department of Statistics and Electrical and Computer Engineering, Rice University, 6100 Main Street, Houston, TX, USA

²Dipartimento di Fisica "M. Merlin", Università degli studi di Bari "A. Moro", Via Orabona, 4, Bari, Italy

³Sezione di Bari, Istituto Nazionale di Fisica Nucleare, Via Orabona, 4, Bari, Italy

⁴Ontario Institute for Cancer Research, Informatics and Bio-computing Program, MaRS Centre, 661 University Avenue, Suite 510, Toronto, ON, Canada

⁵IBM Computational Biology Center, IBM Research, 1101 Kitchawan Road, Yorktown Heights, NY, USA

⁶Sage Bionetworks, 1100 Fairview Avenue North, M1-C108, Seattle, WA, USA

⁷School of Behavioral and Brain Sciences, The University of Texas at Dallas, 800 W. Campbell Road, Richardson, TX, USA

⁸Rush Alzheimer's Disease Center, Rush University Medical Center, 600 South Paulina, Suite 1028, Chicago, IL, USA

⁹Department of Biology, Brigham Young University, 4102 LSB, Provo, UT, USA

¹⁰The Microsoft Research- University of Trento Centre- COSBI, Piazza Manifattura 1, Rovereto, Italy

¹¹Graduate Institute of Biomedical Electronics and Bioinformatics, National Taiwan University, No. 1, Sec. 4, Roosevelt Rd. Taipei, Taiwan

¹²Quantitative Biomedical Research Center, The University of Texas Southwestern Medical Center, Suite NC8.512 5323 Harry Hines Blvd. Dallas, TX, USA

¹³Department of Bio-Industrial Mechatronics Engineering, National Taiwan University, No. 1, Sec. 4, Roosevelt Rd. Taipei, Taiwan

¹⁴Innovation Center for Big Data and Digital Convergence, Yuan Ze University, 135 Yuan-Tung Road, Taoyuan, Taiwan

¹⁵Department of Neurology, Massachusetts General Hospital, 65 Landsdowne St, Suite 200, Cambridge, MA, USA

¹⁶Department of Neurology, Harvard Medical School, 25 Shattuck St, Boston, MA USA

- ¹⁷Center for Biomedical Image Computing and Analytics, University of Pennsylvania, 3600 Market Street, Philadelphia, PA, USA
- ¹⁸School of Computer Science, Fudan University, 220 Handan Road, Shanghai, Shanghai, China
- ¹⁹Shanghai Key Lab of Intelligent Information Processing, Fudan University, 220 Handan Road, Shanghai, Shanghai, China
- ²⁰NIHR Biomedical Research Centre for Mental Health, Kings College London, Box P096, De Crespigny Park, London, UK
- ²¹Institute of Psychiatry, Psychology and Neuroscience, MRC Social, Genetic and Developmental Psychiatry Centre, Kings College London, Box P096, De Crespigny Park, London, UK
- ²²Department of Pediatrics-Neurology, Baylor College of Medicine, 1250 Moursund street, Houston, TX, USA
- ²³Università degli Studi di Genova, via Dodecaneso 33, Genova, Italy
- ²⁴Sanders-Brown Center on Aging, University of Kentucky, 800 S. Limestone Street, Lexington, KY, USA
- ²⁵Department of Biostatistics, University of Kentucky, 725 Rose Street, Lexington, KY, USA
- ²⁶Bonn-Aachen International Center for IT, University of Bonn, Dahlmannstr. 2, Bonn, Germany
- ²⁷Cambridge Institute for Medical Research, University of Cambridge and University of Toronto, Hills Road, Cambridge, CB2 UK
- ²⁸McGovern Institute for Brain Research, Massachusetts Institute of Technology, 43 Vassar St, 46-4033F, Cambridge, MA, USA
- ²⁹Department of Otology and Laryngology, Harvard Medical School, Boston, MA, USA
- ³⁰Luxembourg Centre for Systems Biomedicine, University of Luxembourg, 7, avenue des Hauts Fourneaux, Esch-sur-Alzette, Luxembourg
- ³¹Division of Genetics, Department of Medicine, Brigham and Women's Hospital, Broad Institute and Harvard Medical School, 41 Avenue Louis Pasteur, Suite 301, Boston, MA, USA
- ³²Department of Computational Medicine and Bioinformatics, University of Michigan, 500 South State Street, Ann Arbor, MI, USA
- ³³Department of Electrical Engineering and Computer Science, University of Michigan, 500 South State Street, Ann Arbor, MI, USA
- ³⁴Department of Internal Medicine, University of Michigan, 500 South State Street, Ann Arbor, MI, USA

- ³⁵Department of Biostatistics, The University of North Carolina at Chapel Hill, McGavran Greenberg Hall, CB#7422, Chapel Hill, NC, USA
- ³⁶Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Solnavägen 1, Stockholm, Sweden Paolo Inglese
- ³⁷Department of Computer science and Statistics, Daegu University, Gyeongsan-si, Gyeongsangbuk-do, Republic of Korea
- ³⁸Department of Surgery and Cancer, Faculty of Medicine, Imperial College London, London, UK
- ³⁹Department of Psychiatry, University of Oxford, Warneford Hospital, Oxford, UK
- ⁴⁰Fraunhofer Institute for Algorithms and Scientific Computing (SCAI), Department for Bioinformatics, Schloss Birlinghoven, Sankt Augustin, Germany
- ⁴¹Computational Biology Institute, The George Washington University, 45085 University Drive, Suite 305, Ashburn, VA, USA
- ⁴²Department of Biostatistics, Kings College London, Box P096, De Crespigny Park, SE5 8AF, London, UK
- ⁴³Department of Neuroimmunology, Foundation Institut de Recerca, Hospital Universitari Vall d'Hebron, Ps Vall Hebron 119-129, Barcelona, Spain
- ⁴⁴Department of Biology, Brigham Young University, 4102 Life Sciences Building, Brigham Young University, Provo, UT, USA
- ⁴⁵Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Solnavägen 1, Stockholm, Sweden
- ⁴⁶Genetics and Genomics Sciences Department, Icahn School of Medicine at Mount Sinai, 1425 Madison Ave, New York, NY, USA
- ⁴⁷Genome and systems biology degree program, National Taiwan University, No. 1, Sec. 4, Roosevelt Rd. Taipei, Taiwan
- ⁴⁸Department of Radiology and Medical Imaging, The University of Virginia, Charlottesville, VA, USA
- ⁴⁹Global CEO Initiative on Alzheimer's disease, 1101 K St., NW, #400, Washington, DC
- ⁵⁰Radiology, Medicine, Psychiatry, and Neurology, UCSF, SFVAMC, 4150 Clement Street (114M), San Francisco, CA, USA
- ⁵¹Department of Computer Science, Hunter College, The City University of New York, 695 Park Avenue, New York, NY, USA
- ⁵²Centre for Computational Systems Biology, Fudan University, 220 Handan Road, Shanghai, China
- ⁵³Department of Medical Biophysics, University of Toronto, Toronto, Canada
- ⁵⁴Department of Pharmacology & Toxicology, University of Toronto, Toronto, Canada

Acknowledgments

The following people provided final submissions, but did not participate in the community phase: Lorna Barron¹, Oliver Barron¹, Riccardo Bellazzi², Jungwoo Chang¹, Marianne H Cowherd¹, Grace Ganzel¹, Łukasz Grad³, Inhan Lee¹, Ivan Limongelli², Simone Marini², Szymon Migacz³, Ettore Rizzo², Witold R Rudnicki^{3,4}, Andrzej Sutecki³, Leo Tunkle¹, Francesca Vitali²

This study was supported by the following individuals and organizations: Alan Evans (McGill University), Gaurav Pandey (MSSM), Gil Rabinovici (UCSF), Kaj Blennow (Göteborg University), Kristine Yaffe (UCSF), Maria Isaac (EMA), Nolan Nichols (University of Washington), Paul Thompson (UCLA), Reisa Sperling (Harvard), Scott Small (Columbia), Guy Eakin (BrightFocus Foundation), Maria Carillo (Alzheimer's Association), Neil Buckholz (NIA), Alzheimer's Research UK, European Medicines Agency, Global CEO Initiative on Alzheimer's Disease, Pfizer, Inc, Ray and Dagmar Dolby Family Fund, Rosenberg Alzheimer's Project, Sanofi S.A, and Takeda Pharmaceutical Company Ltd, USAgainstAlzheimer's.

Study data were provided by the following groups:

The Alzheimer's Disease Neuroimaging Initiative (ADNI)

ADNI is funded by the National Institutes of Health (U01 AG024904), the National Institute on Aging, the National Institute of Biomedical Imaging and Bioengineering, and through generous contributions from the following: Alzheimer's Association; Alzheimer's Drug Discovery Foundation; BioClinica, Inc.; Biogen Idec Inc.; Bristol-Myers Squibb Company; Eisai Inc.; Elan Pharmaceuticals, Inc.; Eli Lilly and Company; F. Hoffmann-La Roche Ltd and its affiliated company Genentech, Inc.; GE Healthcare; Innogenetics, N.V.; IXICO Ltd.; Janssen Alzheimer Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research & Development LLC.; Medpace, Inc.; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.; NeuroRx Research; Novartis Pharmaceuticals Corporation; Pfizer Inc.; Piramal Imaging; Servier; Synarc Inc.; and Takeda Pharmaceutical Company. The Canadian Institutes of Health Research is providing funds to support ADNI clinical sites in Canada. Private sector contributions are facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee organization is the Northern California Institute for Research and Education, and the study is coordinated by the Alzheimer's Disease Cooperative Study at the University of California, San Diego. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of Southern California. This research was also supported by NIH grants P30 AG010129 and K01 AG030514.

The Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago

Data collection was supported through funding by NIA grants P30AG10161, R01AG15819, R01AG17917, R01AG30146, R01AG36836, U01AG32984, and U01AG46152, the Illinois Department of Public Health, and the Translational Genomics Research Institute.

European AddNeuroMed study

The AddNeuroMed data are from a public-private partnership supported by EFPIA companies, SMEs and the EU under the FP6 programme. Clinical leads responsible for data collection are Iwona Kłoszewska (Lodz), Simon Lovestone (London), Patrizia Mecocci (Perugia), Hilka Soininen (Kuopio), Magda Tsolaki (Thessaloniki), and Bruno Vellas (Toulouse).

References

1. Mueller SG, Weiner MW, Thal LJ, Petersen RC, Jack C, Jagust W, et al. The Alzheimer's disease neuroimaging initiative. *Neuroimaging clinics of North America*. 2005 Nov; 15(4):869–77. xi–xii. [PubMed: 16443497]
2. Bennett DA, Schneider JA, Arvanitakis Z, Wilson RS. Overview and findings from the religious orders study. *Current Alzheimer research*. 2012 Jul; 9(6):628–45. [PubMed: 22471860]

¹GIDAS, miRcore, 2929 Plymouth Rd. Suite 207, Ann Arbor, MI, USA

²Electrical, Computer and Biomedical Engineering Department, Via Ferrata, 1, Pavia, Italy

³Interdisciplinary Centre for Mathematical and Computational Modelling, Pawińskiego 5A, Warsaw, Poland

⁴Department of Bioinformatics, University of Białystok, Ciołkowskiego 1M, Białystok, Poland

3. Bennett DA, Schneider JA, Buchman AS, Barnes LL, Boyle PA, Wilson RS. Overview and findings from the rush Memory and Aging Project. *Current Alzheimer research*. 2012 Jul; 9(6):646–63. [PubMed: 22471867]
4. Lovestone S, Francis P, Kloszewska I, Mecocci P, Simmons A, Soininen H, et al. AddNeuroMed--the European collaboration for the discovery of novel biomarkers for Alzheimer's disease. *Annals of the New York Academy of Sciences*. 2009 Oct. 1180:36–46. [PubMed: 19906259]
5. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *Journal of psychiatric research*. 1975 Nov; 12(3):189–98. [PubMed: 1202204]
6. Ercoli LM, Siddarth P, Dunkin JJ, Bramen J, Small GW. MMSE items predict cognitive decline in persons with genetic risk for Alzheimer's disease. *Journal of geriatric psychiatry and neurology*. 2003 Jun; 16(2):67–73. [PubMed: 12801154]
7. Hsiung GY, Alipour S, Jacova C, Grand J, Gauthier S, Black SE, et al. Transition from cognitively impaired not demented to Alzheimer's disease: an analysis of changes in functional abilities in a dementia clinic cohort. *Dementia and geriatric cognitive disorders*. 2008; 25(6):483–90. [PubMed: 18417973]
8. Vemuri P, Wiste HJ, Weigand SD, Shaw LM, Trojanowski JQ, Weiner MW, et al. MRI and CSF biomarkers in normal, MCI, and AD subjects: predicting future clinical change. *Neurology*. 2009 Jul 28; 73(4):294–301. [PubMed: 19636049]
9. Ridge PG, Ebbert MT, Kauwe JS. Genetics of Alzheimer's disease. *BioMed research international*. 2013; 2013:254954. [PubMed: 23984328]
10. Lambert JC, Ibrahim-Verbaas CA, Harold D, Naj AC, Sims R, Bellenguez C, et al. Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nature genetics*. 2013 Dec; 45(12):1452–8. [PubMed: 24162737]
11. Ridge PG, Mukherjee S, Crane PK, Kauwe JS. Alzheimer's Disease Genetics C. Alzheimer's disease: analyzing the missing heritability. *PloS one*. 2013; 8(11):e79771. [PubMed: 24244562]
12. Bennett DA, Schneider JA, Arvanitakis Z, Kelly JF, Aggarwal NT, Shah RC, et al. Neuropathology of older persons without cognitive impairment from two community-based studies. *Neurology*. 2006 Jun 27; 66(12):1837–44. [PubMed: 16801647]
13. Price JL, Morris JC. Tangles and plaques in nondemented aging and "preclinical" Alzheimer's disease. *Annals of neurology*. 1999 Mar; 45(3):358–68. [PubMed: 10072051]
14. Davatzikos C, Xu F, An Y, Fan Y, Resnick SM. Longitudinal progression of Alzheimer's-like patterns of atrophy in normal older adults: the SPARE-AD index. *Brain: a journal of neurology*. 2009 Aug; 132(Pt 8):2026–35. [PubMed: 19416949]
15. Misra C, Fan Y, Davatzikos C. Baseline and longitudinal patterns of brain atrophy in MCI patients, and their use in prediction of short-term conversion to AD: results from ADNI. *NeuroImage*. 2009 Feb 15; 44(4):1415–22. [PubMed: 19027862]
16. Im K, Lee JM, Seo SW, Hyung Kim S, Kim SI, Na DL. Sulcal morphology changes and their relationship with cortical thickness and gyral white matter volume in mild cognitive impairment and Alzheimer's disease. *NeuroImage*. 2008 Oct 15; 43(1):103–13. [PubMed: 18691657]
17. Haight TJ, Jagust WJ. Alzheimer's Disease Neuroimaging I. Relative contributions of biomarkers in Alzheimer's disease. *Annals of epidemiology*. 2012 Dec; 22(12):868–75. [PubMed: 23102709]
18. Nho K, Risacher SL, Crane PK, DeCarli C, Glymour MM, Habeck C, et al. Voxel and surface-based topography of memory and executive deficits in mild cognitive impairment and Alzheimer's disease. *Brain imaging and behavior*. 2012 Dec; 6(4):551–67. [PubMed: 23070747]
19. Thung KH, Wee CY, Yap PT, Shen D. Alzheimer's Disease Neuroimaging I. Neurodegenerative disease diagnosis using incomplete multi-modality data via matrix shrinkage and completion. *NeuroImage*. 2014 May 1. 91:386–400. [PubMed: 24480301]
20. Escott-Price V, Sims R, Bannister C, Harold D, Vronskaya M, Majounie E, et al. Common polygenic variation enhances risk prediction for Alzheimer's disease. *Brain: a journal of neurology*. 2015 Dec; 138(Pt 12):3673–84. [PubMed: 26490334]
21. Lee SH, Harold D, Nyholt DR, Consortium AN, International Endogene C, Genetic, et al. Estimation and partitioning of polygenic variation captured by common SNPs for Alzheimer's

- disease, multiple sclerosis and endometriosis. *Hum Mol Genet.* 2013 Feb 15; 22(4):832–41. [PubMed: 23193196]
22. Chatterjee N, Wheeler B, Sampson J, Hartge P, Chanock SJ, Park JH. Projecting the performance of risk prediction based on polygenic analyses of genome-wide association studies. *Nature genetics.* 2013 Apr; 45(4):400–5. 5e1–3. [PubMed: 23455638]
 23. Manolio TA. Bringing genome-wide association findings into clinical use. *Nat Rev Genet.* 2013 Aug; 14(8):549–58. [PubMed: 23835440]
 24. Scarmeas N, Stern Y, Mayeux R, Luchsinger JA. Mediterranean diet, Alzheimer disease, and vascular mediation. *Arch Neurol.* 2006 Dec; 63(12):1709–17. [PubMed: 17030648]
 25. Podewils LJ, Guallar E, Kuller LH, Fried LP, Lopez OL, Carlson M, et al. Physical activity, APOE genotype, and dementia risk: findings from the Cardiovascular Health Cognition Study. *Am J Epidemiol.* 2005 Apr 1; 161(7):639–51. [PubMed: 15781953]
 26. Lindsay J, Laurin D, Verreault R, Hebert R, Helliwell B, Hill GB, et al. Risk factors for Alzheimer's disease: a prospective analysis from the Canadian Study of Health and Aging. *Am J Epidemiol.* 2002 Sep 1; 156(5):445–53. [PubMed: 12196314]
 27. Wang HX, Karp A, Winblad B, Fratiglioni L. Late-life engagement in social and leisure activities is associated with a decreased risk of dementia: a longitudinal study from the Kungsholmen project. *Am J Epidemiol.* 2002 Jun 15; 155(12):1081–7. [PubMed: 12048221]
 28. Guerreiro R, Wojtas A, Bras J, Carrasquillo M, Rogaeva E, Majounie E, et al. TREM2 variants in Alzheimer's disease. *N Engl J Med.* 2013 Jan 10; 368(2):117–27. [PubMed: 23150934]
 29. Jonsson T, Stefansson H, Steinberg S, Jonsdottir I, Jonsson PV, Snaedal J, et al. Variant of TREM2 associated with the risk of Alzheimer's disease. *N Engl J Med.* 2013 Jan 10; 368(2):107–16. [PubMed: 23150908]
 30. Jonsson T, Atwal JK, Steinberg S, Snaedal J, Jonsson PV, Bjornsson S, et al. A mutation in APP protects against Alzheimer's disease and age-related cognitive decline. *Nature.* 2012 Aug 2; 488(7409):96–9. [PubMed: 22801501]
 31. Gray KR, Wolz R, Heckemann RA, Aljabar P, Hammers A, Rueckert D, et al. Multi-region analysis of longitudinal FDG-PET for the classification of Alzheimer's disease. *NeuroImage.* 2012 Mar; 60(1):221–9. [PubMed: 22236449]
 32. James OG, Doraiswamy PM, Borges-Neto S. PET Imaging of Tau Pathology in Alzheimer's Disease and Tauopathies. *Front Neurol.* 2015; 6:38. [PubMed: 25806018]
 33. Gross AL, Sherva R, Mukherjee S, Newhouse S, Kauwe JS, Munsie LM, et al. Calibrating longitudinal cognition in Alzheimer's disease across diverse test batteries and datasets. *Neuroepidemiology.* 2014; 43(3–4):194–205. [PubMed: 25402421]

Systematic Review

Extensive literature searches using PubMed establish this as the largest study to date using demographic, clinical, imaging and genetic data to predict cognitive decline and the first major instance of crowdsourcing analysis in AD.

Interpretation

Over 500 scientists worldwide in the analytical portion of the Challenge, demonstrating the viability of crowdsourced approaches in AD research. Unfortunately, we were unable to detect meaningful predictors of either cognitive decline or resilience through this effort.

Future Directions

This experiment in crowdsourcing AD analyses is an invaluable first-of-its-kind contribution that provides a snapshot of both the strengths and limitations in big data analytics in AD research. The relative inaccessibility and heterogeneity across data sources severely limits formalized integration. Mandates on data sharing, considerations of standardized data collection, and mechanisms to integrate heterogeneous data are necessary to address these issues. We anticipate that this work will initiate those discussions across the community.




Fig. 1. Challenge overview

The top schematic summarizes the three challenge questions on the left column, the training data in the middle, and the test data on the right, including numbers of subjects. The symbols represent sources of data (demographic, ROS/MAP genetic, and ADNI or ANM brain images and shape information). The bottom panel provides example brain image labels and shape information provided to the participants for question 3. Anatomical labels for left cortical regions are shown on the left and just a couple of the cortical surface shape measures are shown on the right (travel depth on top and mean curvature below), for both uninflated and inflated surfaces (top and bottom rows, respectively).




Figure 2. Performance evaluation results

Panels a, b, and c report the p-values (in negative log10 scale) for intersection union tests investigating which teams performed better than random for questions 1, 2, and 3, respectively. Explicitly, for question 1 (panel a) we tested the null hypothesis that at least one of the four correlation coefficients (namely, Pearson/clinical, Pearson/clinical + genetics, Spearman/clinical, Spearman/clinical + genetics) is equal to zero, against the alternative that all four correlation coefficients are larger than zero. Adopting a 0.05 significance level, 26 out of the 32 submissions were statistically better than random, after Bonferroni multiple testing correction for 32 tests (submissions crossing the black vertical line). For question 2

(panel b), we tested the null hypothesis that balanced accuracy = 0.5 or AUC = 0.5, against the alternative that balanced accuracy > 0.5 and AUC > 0.5. In this case, no model performed significantly better than random and, therefore, no best performer was declared. For question 3 (panel c), we tested the null hypothesis that Pearson's correlation (COR) or Lin's concordance correlation coefficient (CCC) are equal to zero, against the alternative that both COR and CCC are larger than zero. Adopting a 0.05 significance level, all 23 submissions were statistically better than random, after Bonferroni correction. For all three questions, the p-values were computed from an empirical null distribution based on 10,000 permutations. Panels d and e report the bootstrapped assessment of ranks for questions 1 and 3, respectively. Samples were resampled with replacement from the original data (true outcome and team's predictions), and the ranks of the different teams were re-assessed in each of 100,000 re-samplings. Submissions were sorted according to the median of their bootstrapped average ranking distributions. The black horizontal line represents the posterior odds cutoff from the Bayesian analysis. Teams above the black line are statistically tied to the top ranked model, according to a posterior odds threshold of 3.