

## **UC Davis**

### **Endocrinology, Diabetes, and Metabolism**

#### **Title**

Does Diabetic Ketoacidosis Increase Risk of Diabetic Nephropathy?

#### **Permalink**

<https://escholarship.org/uc/item/8t719544>

#### **Authors**

Ramesh, Janani

Huang, Jia

Pitts, Casey

et al.

#### **Publication Date**

2021

#### **Data Availability**

The data associated with this publication are not available for this reason: N/A





# Does Diabetic Ketoacidosis Increase Risk of Diabetic Nephropathy?



Janani Ramesh B.A.<sup>1</sup> Jia Xin Huang M.D.<sup>1</sup> Casey Pitts M.D.<sup>2</sup> Lindsey Loomba-Albrecht M.D.<sup>1</sup> Sage Myers M.D.<sup>2</sup> Nicole Glaser M.D.<sup>1</sup>  
1 – UC Davis Medical Center 2 – Children’s Hospital of Philadelphia

## Introduction


Diabetic nephropathy is the most common cause of death in patients with type 1 diabetes (T1D).<sup>1</sup>

- T1D patients hospitalized with Diabetic Ketoacidosis (DKA) are more likely to experience acute kidney injury (AKI)<sup>2,3</sup>
- In other disease states, AKI increases risk of developing chronic kidney disease.<sup>2</sup>
- Proposed pathophysiology: AKI induces endothelial dysfunction, T cell activation, fibrosis, and tubular injury.<sup>4</sup>
- Mechanism suggests that DKA induced AKI may increase risk of diabetic nephropathy.
- To date, association between DKA and future diabetic nephropathy has not previously been investigated in children.
- Microalbuminuria is one of the earliest markers of diabetic nephropathy.<sup>1</sup> and can be used to track if there is an increased risk of diabetic nephropathy in T1D pediatric patients with ≥ 1 episode of DKA.

## Hypothesis

T1D patients with a history of DKA have higher urine microalbumin levels, suggesting an increased likelihood of developing diabetic nephropathy.

## Methods



## Demographics

<b>Total Participants</b>		<b>2394</b>
<b>Age</b>		
	At Diagnosis of Diabetes Mellitus*	9.5 ± 4.4 years
	At Study Enrollment*	16.8 ± 5.1 years
<b>Sex</b>		
	Female	1087 (45.4%)
	Male	1307 (54.6%)
<b>DKA at Diagnosis, Yes</b>		841 (35.1%)
	pH*	7.18 ± 0.1
<b>DKA after Diagnosis, Yes</b>		413 (17.3%)
	pH*	7.17 ± 0.11
	1 DKA episode	257 (10.7%)
	2 DKA episodes	66 (2.8%)
	3 DKA episodes	27 (1.1%)
	> 3 DKA episodes	63 (2.6%)
<b>HbA1C*</b>		8.4% ± 1.5
<b>Patients Diagnosed with Microalbuminuria</b>		62 (2.6%)
	Age at Diagnosis of Microalbuminuria*	15 ± 4.4 years
<b>Patients with Possible Microalbuminuria</b>		128 (5.3%)

\*Mean (SD)

Cr levels collected to calculate AKI stages. Data not yet available.

## Preliminary Results

	Developed Microalbuminuria	No History of Microalbuminuria	Total
<b>History of DKA</b>	121	964	1085
<b>No History of DKA</b>	69	1240	1309
<b>Total</b>	190	2204	2394

The evidence cannot reject the null hypothesis. This suggests that there is a significant association between a history of DKA and developing microalbuminuria.  $X^2 (1, N = 2394) = 28.08, p < 0.0001$ .

## Discussion

Preliminary analysis suggests a significant association between having an episode of DKA and developing microalbuminuria. Analyses involving multivariable models are needed to adjust for possible confounding variables: age, diabetes duration, and glycemic control (HbA1c). In addition, comparisons of patients with and without AKI during DKA are needed to determine whether effects of DKA on future risk of microalbuminuria are related to DKA per se or to the occurrence of AKI during DKA.

Other Limitations:

- Lapses in medical history of patients whose records are not fully in the UC Davis or CHOP system.
- “Possible history of microalbuminuria” are patients with one urinalysis showing elevated MA:Cr but were lost to follow up, resulting in lack of diagnosis.

Future studies should explore the mechanisms upon which we can act to either slow or prevent the progression of DKA induced AKI to diabetic nephropathy.

Understanding this risk would help better inform our current practices in the prevention of DKA and early prophylaxis against AKI. Doing so may help slow the progression to diabetic nephropathy following DKA episodes.

## References

1. Bjornstad P, Cherney D, Maahs D.M. Early Diabetic Nephropathy in Type 1 Diabetes – New Insights. *Current Opinion in Endocrinology & Diabetes and Obesity*. 2014; 21(4):279–286.
2. Hursh BE, Ronsley R, Islam N, Mammen C, Panagiotopoulos C. Acute Kidney Injury in Children With Type 1 Diabetes Hospitalized for Diabetic Ketoacidosis. *JAMA Pediatrics*. 2017; 171(5):e170020.
3. Orban JC, Maizière EM, Ghaddab A, Van Obberghen E, Ichai C. Incidence and characteristics of acute kidney injury in severe diabetic ketoacidosis. *PLoS One*. 2014; 9(10):e110925.
4. Guzzi F, Cirillo L, Roperto RM, Romagnani P, Lazzeri E. Molecular Mechanisms of the Acute Kidney Injury to Chronic Kidney Disease Transition: An Updated View. *International Journal of Molecular Sciences*. 2019; 20(19): 4941.