

Supplement of Atmos. Chem. Phys., 15, 10839–10856, 2015
<http://www.atmos-chem-phys.net/15/10839/2015/>
doi:10.5194/acp-15-10839-2015-supplement
© Author(s) 2015. CC Attribution 3.0 License.



Supplement of


**Springtime daily variations in lower-tropospheric ozone over east Asia:
the role of cyclonic activity and pollution as
observed from space with IASI**

G. Dufour et al.

Correspondence to: G. Dufour (gaelle.dufour@lisa.u-pec.fr)


The copyright of individual parts of the supplement might differ from the CC-BY 3.0 licence.

NOAA HYSPLIT MODEL
 Backward trajectories ending at 0000 UTC 06 May 08
 GDAS Meteorological Data




1
 2 Figure S1. 36-hours backward trajectories ending at 3km on 6 May 2008 in the region of
 3 Tokyo.

NOAA HYSPLIT MODEL
 Backward trajectories ending at 0000 UTC 13 May 08
 GDAS Meteorological Data



4
 5 Figure S2. 48-hours backward trajectories ending at 3km on 13 May 2008 in the region of
 6 South Korea.

NOAA HYSPLIT MODEL
 Backward trajectories ending at 0000 UTC 11 May 08
 GDAS Meteorological Data



1
 2 Figure S3. 24-hours backward trajectories ending at 3km on 11 May 2008 in the region of
 3 Beijing.

4