Construction of extremal Type II \mathbb{Z}_{2k} -codes

Masaaki Harada*

June 22, 2022

Abstract

We give methods for constructing many self-dual \mathbb{Z}_m -codes and Type II \mathbb{Z}_{2k} -codes of length 2n starting from a given self-dual \mathbb{Z}_m -code and Type II \mathbb{Z}_{2k} -code of length 2n, respectively. As an application, we construct extremal Type II \mathbb{Z}_{2k} -codes of length 24 for $k = 4, 5, \ldots, 20$ and extremal Type II \mathbb{Z}_{2k} -codes of length 32 for $k = 4, 5, \ldots, 10$. We also construct new extremal Type II \mathbb{Z}_4 -codes of lengths 56 and 64.

1 Introduction

A \mathbb{Z}_m -code of length n is a \mathbb{Z}_m -submodule of \mathbb{Z}_m^n , where m is a positive integer with $m \geq 2$ and \mathbb{Z}_m denotes the ring of integers modulo m. A \mathbb{Z}_m -code C of length n is self-dual if $C = C^{\perp}$, where C^{\perp} denotes the dual code of C. Self-dual codes are one of the most interesting classes of codes. This interest is justified by many combinatorial objects and algebraic objects related to self-dual codes (see e.g. [26] and the references given therein).

Many methods for constructing self-dual codes are known. For example, starting from a given self-dual code with generator matrix (I_n A), by transforming A, some methods for constructing many self-dual codes are known, where I_n denotes the identity matrix of order n (see e.g. [14], [19], [23] and [27]). In this paper, we give a new method for constructing many self-dual \mathbb{Z}_m -codes starting from a given self-dual \mathbb{Z}_m -code with generator matrix (I_n A) by transforming A. For self-dual codes over finite fields, the corresponding result can be found in [23].

^{*}Research Center for Pure and Applied Mathematics, Graduate School of Information Sciences, Tohoku University, Sendai 980–8579, Japan. email: mharada@tohoku.ac.jp.

Now let us consider the case m=2k. A binary doubly even self-dual code is often called Type II. Self-dual \mathbb{Z}_{2k} -codes with the property that all Euclidean weights are multiples of 4k are called Type II \mathbb{Z}_{2k} -codes (see [2] and [22] for k=2 and [1] for $k\geq 3$). Type II \mathbb{Z}_{2k} -codes are a remarkable class of self-dual codes related to even unimodular lattices. For example, by Construction A, Type II \mathbb{Z}_{2k} -codes give even unimodular lattices. There is a Type II \mathbb{Z}_{2k} -code of length n if and only if n is divisible by eight [1]. For Type II \mathbb{Z}_{2k} -codes C of length n, the following upper bound on the minimum Euclidean weight

$$d_E(C) \le 4k \left\lfloor \frac{n}{24} \right\rfloor + 4k \tag{1}$$

holds if k=1 [24], if k=2 [2] (see also [22]) and if $k \in \{3,4,5,6\}$ [21]. Also, the bound (1) holds under the assumption that $\lfloor n/24 \rfloor \leq k-2$ for arbitrary k [1]. When the bound (1) holds, we say that a Type II \mathbb{Z}_{2k} -code meeting (1) with equality is extremal. There have been significant researches on constructing extremal Type II \mathbb{Z}_{2k} -codes for k=1 and 2 (see e.g. [2], [14], [19], [22], [26], [27] and the references given therein). In this paper, we also give a new method for constructing many Type II \mathbb{Z}_{2k} -codes starting from a given Type II \mathbb{Z}_{2k} -code with generator matrix (I_n A) by transforming A. This generalizes the methods given in [14] and [23]. By four-negacirculant codes and by the new method, we construct extremal Type II \mathbb{Z}_{2k} -codes of length 24 for $k=4,5,\ldots,20$ and extremal Type II \mathbb{Z}_{2k} -codes of length 32 for $k=4,5,\ldots,10$. We also construct new extremal Type II \mathbb{Z}_4 -codes of lengths 56 and 64.

The paper is organized as follows. In Section 2, we give the definitions and basic facts on self-dual \mathbb{Z}_m -codes and Type II \mathbb{Z}_{2k} -codes used throughout this paper. In Section 3, we give a method for constructing self-dual \mathbb{Z}_m -codes (resp. Type II \mathbb{Z}_{2k} -codes) of length 2n starting from a given self-dual \mathbb{Z}_m -code (resp. Type II \mathbb{Z}_{2k} -code) of length 2n having generator matrix (I_n A), by transforming A (Theorem 3.1 (resp. Theorem 3.4)). In Section 4, we construct extremal Type II \mathbb{Z}_{2k} -codes of length 24, by four-negacirculant codes and by Theorem 3.4 for $k = 4, 5, \ldots, 20$. Consequently, it is shown that if $k \in \{2, 3, \ldots, 20\}$ then there is an extremal Type II \mathbb{Z}_{2k} -code C such that $C^{(2)} \cong B$ for every binary doubly even self-dual code B of length 24, where $C^{(2)}$ denotes the binary part of C and $C \cong D$ means that C and D are equivalent. In Section 5, we construct extremal Type II \mathbb{Z}_{2k} -codes of length 32, by four-negacirculant codes and by Theorem 3.4 for $k = 4, 5, \ldots, 10$. Consequently, it is shown that if $k \in \{2, 3, \ldots, 10\}$ then there is an extremal

Type II \mathbb{Z}_{2k} -code C such that $C^{(2)} \cong B$ for every binary extremal doubly even self-dual code B of length 32. In Section 6, by four-negacirculant codes and by Theorem 3.4, we construct new extremal Type II \mathbb{Z}_4 -codes of lengths 56 and 64.

All computer calculations in this paper were done with the help of MAGMA [3].

2 Preliminaries

In this section, we give the definitions and basic facts on self-dual \mathbb{Z}_m -codes and Type II \mathbb{Z}_{2k} -codes used throughout this paper.

Let \mathbb{Z}_m be the ring of integers modulo m, where m is a positive integer with $m \geq 2$. A \mathbb{Z}_m -code C of length n is a \mathbb{Z}_m -submodule of \mathbb{Z}_m^n . An element of C is called a *codeword*. A *generator matrix* of C is a matrix whose rows generate C. We define the inner product of $x = (x_1, x_2, \ldots, x_n)$ and $y = (y_1, y_2, \ldots, y_n) \in \mathbb{Z}_m^n$ by

$$\langle x, y \rangle = x_1 y_1 + x_2 y_2 + \dots + x_n y_n.$$

The dual code C^{\perp} of C is defined as

$$C^{\perp} = \{ x \in \mathbb{Z}_m^n \mid \langle x, y \rangle = 0 \text{ for all } y \in C \}.$$

A code C is self-dual if $C = C^{\perp}$. Two \mathbb{Z}_m -codes C and C' are equivalent, denoted $C \cong C'$, if one can be obtained from the other by permuting the coordinates and (if necessary) changing the signs of certain coordinates.

We now consider self-dual \mathbb{Z}_{2k} -codes. In this paper, we take the set \mathbb{Z}_{2k} to be either $\{0, 1, \ldots, 2k-1\}$ or $\{0, \pm 1, \ldots, \pm (k-1), k\}$, using whichever form is more convenient. We define a map ρ from \mathbb{Z}_{2k} to \mathbb{Z} as follows

$$\rho(i) = \begin{cases} 0, \dots, k & \text{if } i = 0, \dots, k, \text{ respectively,} \\ 1 - k, \dots, -1 & \text{if } i = k + 1, \dots, 2k - 1, \text{ respectively.} \end{cases}$$
 (2)

The Euclidean weight $\operatorname{wt}_E(v)$ of a vector $v = (v_1, v_2, \dots, v_n) \in \mathbb{Z}_{2k}^n$ is

$$\sum_{i=1}^{k} i^{2} n_{i}(v) + \sum_{i=k+1}^{2k-1} (2k-i)^{2} n_{i}(v),$$

where

$$n_i(v) = |\{j \in \{1, 2, \dots, n\} \mid \rho(v_j) \equiv i \pmod{2k}\}| \ (i = 0, 1, \dots, 2k - 1).$$

A binary doubly even self-dual code is often called Type II. Using Euclidean weights, the notion of binary Type II codes has been generalized as follows. Self-dual \mathbb{Z}_{2k} -codes with the property that all Euclidean weights are multiples of 4k are called Type II \mathbb{Z}_{2k} -codes (see [2] and [22] for k=2 and [1] for $k\geq 3$). It was shown in [1] that there is a Type II \mathbb{Z}_{2k} -code of length n if and only if n is divisible by eight.

Throughout this paper, I_n denotes the identity matrix of order n, and A^T denotes the transpose of a matrix A. The following gives a criteria for self-dual \mathbb{Z}_{2k} -codes and Type II \mathbb{Z}_{2k} -codes.

- **Lemma 2.1.** (i) Let C be a \mathbb{Z}_{2k} -code of length 2n having generator matrix of form $(I_n \ A)$, where A is an $n \times n$ matrix. If $AA^T = -I_n$, then C is self-dual.
 - (ii) Let C be a self-dual \mathbb{Z}_{2k} -code having generator matrix G. If each row of G has Euclidean weight divisible by 4k, then C is Type II.

Proof. (i) is trivial. (ii) follows from
$$[1, Lemma 2.2]$$
.

The minimum Euclidean weight $d_E(C)$ of a Type II \mathbb{Z}_{2k} -code C is the smallest Euclidean weight among all nonzero codewords of C. For Type II \mathbb{Z}_{2k} -codes of length n, the upper bound (1) on the minimum Euclidean weight $d_E(C)$ holds if k = 1 [24], if k = 2 [2] and [22] and if $k \in \{3, 4, 5, 6\}$ [21]. Also, the bound (1) holds under the assumption that $\lfloor n/24 \rfloor \leq k - 2$ for arbitrary k [1]. When the bound (1) holds, we say that a Type II \mathbb{Z}_{2k} -code meeting (1) with equality is extremal.

The binary part $C^{(2)}$ of a Type II \mathbb{Z}_{2k} -code C of length n is defined as

$$\{(\rho(c_1) \pmod{2}, \rho(c_2) \pmod{2}, \dots, \rho(c_n) \pmod{2}) \mid (c_1, c_2, \dots, c_n) \in C\},\$$

where ρ is the map given in (2). If a Type II \mathbb{Z}_{2k} -code has generator matrix of form $(I_n \ A)$, then the binary part is a binary doubly even self-dual code [8] and [9]. The following is trivial.

Lemma 2.2. Let C and C' be Type II \mathbb{Z}_{2k} -codes. If $C \cong C'$, then $C^{(2)} \cong C'^{(2)}$.

An $n \times n$ negacirculant matrix has the following form

$$\begin{pmatrix} r_0 & r_1 & r_2 & \cdots & r_{n-1} \\ -r_{n-1} & r_0 & r_1 & \cdots & r_{n-2} \\ -r_{n-2} & -r_{n-1} & r_0 & \cdots & r_{n-3} \\ \vdots & \vdots & \vdots & & \vdots \\ -r_1 & -r_2 & -r_3 & \cdots & r_0 \end{pmatrix}.$$

Let C be the \mathbb{Z}_m -code of length 4n having the following generator matrix

$$\begin{pmatrix}
& I_{2n} & A & B \\
& -B^T & A^T
\end{pmatrix},$$
(3)

where A and B are $n \times n$ negacirculant matrices. Such a code is called a four-negacirculant code [18]. By Lemma 2.1 (i), if $AA^T + BB^T = -I_n$, then C is self-dual. Let C be a four-negacirculant self-dual \mathbb{Z}_{2k} -code with generator matrix of form (3). Let r_A and r_B denote the first rows of A and B, respectively. By Lemma 2.1 (ii), if $\operatorname{wt}_E(r_A) + \operatorname{wt}_E(r_B) \equiv -1 \pmod{4k}$, then C is Type II.

We end this section by describing unimodular lattices and Construction A. A (Euclidean) integral lattice $L \subset \mathbb{R}^n$ in dimension n is unimodular if $L = L^*$, where L^* is the dual lattice under the standard inner product (x, y). A unimodular lattice L is even if the norm (x, x) of every vector x of L is even, and odd otherwise. There is an even unimodular lattice in dimension n if and only if n is divisible by eight. The $minimum\ norm\ min(L)$ of a unimodular lattice L is the smallest norm among all nonzero vectors of L. The $kissing\ number$ of L is the number of vectors of minimum norm in L. We now give a method to construct even unimodular lattices from Type II \mathbb{Z}_{2k} -codes, which is called Construction A. For a Type II \mathbb{Z}_{2k} -code C of length n, define the following lattice

$$A_{2k}(C) = \frac{1}{\sqrt{2k}} \{ \rho(C) + 2k\mathbb{Z}^n \}, \tag{4}$$

where

$$\rho(C) = \{ (\rho(c_1), \rho(c_2), \dots, \rho(c_n)) \mid (c_1, c_2, \dots, c_n) \in C \}.$$

It is known that $A_{2k}(C)$ is an even unimodular lattice with minimum norm $\min\{d_E(C)/2k, 2k\}$, where $d_E(C)$ denotes the minimum Euclidean weight of C [1]. The minimum norms of $A_{2k}(C)$ are used to determine the minimum Euclidean weights of C in Sections 4, 5 and 6.

3 Methods for constructing self-dual \mathbb{Z}_m -codes

In this section, starting from a given self-dual \mathbb{Z}_m -code (resp. Type II \mathbb{Z}_{2k} -code) with generator matrix (I_n A), by transforming A, we give a new method for constructing many self-dual \mathbb{Z}_m -codes (resp. Type II \mathbb{Z}_{2k} -codes).

Theorem 3.1. Let C be a self-dual \mathbb{Z}_m -code of length 2n having generator matrix $(I_n \ A)$. Let r_i be the i-th row of A. Let x and y be vectors of \mathbb{Z}_m^n . Define an $n \times n$ matrix A(x, y), where the i-th row r'_i is given by

$$r_i' = r_i + \langle r_i, y \rangle x - \langle r_i, x \rangle y.$$

Let C(A, x, y) be the \mathbb{Z}_m -code having the following generator matrix

$$(I_n A(x,y)).$$

If $\langle x, x \rangle = \langle y, y \rangle = \langle x, y \rangle = 0$, then C(A, x, y) is a self-dual \mathbb{Z}_m -code of length 2n.

Proof. Since $\langle x, x \rangle = \langle y, y \rangle = \langle x, y \rangle = 0$, we have

$$\langle r'_{i}, r'_{j} \rangle = \langle r_{i}, r_{j} \rangle + \langle r_{j}, y \rangle \langle r_{i}, x \rangle - \langle r_{j}, x \rangle \langle r_{i}, y \rangle + \langle r_{i}, y \rangle \langle x, r_{j} \rangle + \langle r_{i}, y \rangle \langle r_{j}, y \rangle \langle x, x \rangle - \langle r_{i}, y \rangle \langle r_{j}, x \rangle \langle x, y \rangle - \langle r_{i}, x \rangle \langle y, r_{j} \rangle - \langle r_{i}, x \rangle \langle r_{j}, y \rangle \langle y, x \rangle + \langle r_{i}, x \rangle \langle r_{j}, x \rangle \langle y, y \rangle = \langle r_{i}, r_{j} \rangle + \langle r_{j}, y \rangle \langle r_{i}, x \rangle - \langle r_{j}, x \rangle \langle r_{i}, y \rangle + \langle r_{i}, y \rangle \langle x, r_{j} \rangle - \langle r_{i}, x \rangle \langle y, r_{j} \rangle = \langle r_{i}, r_{j} \rangle.$$
(5)

This implies that $AA^T = A(x,y)A(x,y)^T$. By Lemma 2.1 (i), C(A,x,y) is self-dual.

Remark 3.2. For self-dual codes over finite fields, the corresponding result can be found in [23].

Remark 3.3. We modify the above theorem as follows. Let C be a \mathbb{Z}_m -code of length n+n' having generator matrix $(I_n \ A)$, where A is an $n \times n'$ matrix. Then the code C(A, x, y) having generator matrix $(I_n \ A(x, y))$ defined in Theorem 3.1 satisfies $AA^T = A(x, y)A(x, y)^T$.

We consider a method for constructing Type II \mathbb{Z}_{2k} -codes.

Theorem 3.4. Suppose that C is a Type II \mathbb{Z}_{2k} -code of length 2n having generator matrix $(I_n \ A)$. Suppose that x and y are vectors of \mathbb{Z}_{2k}^n satisfying the condition $\operatorname{wt}_E(x) \equiv \operatorname{wt}_E(y) \equiv 0 \pmod{4k}$ and $\langle x, y \rangle = 0$. Then the code C(A, x, y) having generator matrix $(I_n \ A(x, y))$ defined in Theorem 3.1 is a Type II \mathbb{Z}_{2k} -code of length 2n.

Proof. Let ρ be the map given in (2). For a vector $v = (v_1, v_2, \ldots, v_n) \in \mathbb{Z}_{2k}$, define the positive integer $v * v = \sum_{i=1}^{2k-1} i^2 n_i(v)$, where $n_i(v) = |\{j \in \{1, 2, \ldots, n\} \mid \rho(v_j) \equiv i \pmod{2k}\}| \ (i = 0, 1, \ldots, 2k-1)$. By [1, Lemma 2.1], wt_E(v) $\equiv v * v \pmod{4k}$. This implies that $\langle x, x \rangle = \langle y, y \rangle = 0$. Thus, by Theorem 3.1, C(A, x, y) is self-dual. Let r_i and r'_i denote the *i*-th row of A and A(x, y), respectively. By Lemma 2.1 (ii), it is sufficient to show that wt_E(r'_i) $\equiv 4k-1 \pmod{4k}$ ($i=1,2,\ldots,n$). From (5), we have

$$\operatorname{wt}_{E}(r'_{i}) \equiv r'_{i} * r'_{i}$$

$$\equiv r_{i} * r_{i} + x * x + y * y$$

$$\equiv \operatorname{wt}_{E}(r_{i}) + \operatorname{wt}_{E}(x) + \operatorname{wt}_{E}(y)$$

$$\equiv \operatorname{wt}_{E}(r_{i})$$

$$\equiv 4k - 1 \pmod{4k}.$$

The result follows.

Remark 3.5. For the case k=1, the above theorem can be found in [23]. Theorem 5 in [14] can be obtained from the above theorem by putting k=2, x=t and $y=(2,2,\ldots,2)$.

4 Extremal Type II \mathbb{Z}_{2k} -codes of length 24

For k = 4, 5, ..., 20, in this section, we construct extremal Type II \mathbb{Z}_{2k} -codes of length 24, by four-negacirculant codes and by Theorem 3.4.

4.1 Motivation and results

For lengths 8 and 16, every Type II \mathbb{Z}_{2k} -code is extremal. In other words, the smallest possible length for which there is a nontrivial extremal Type II \mathbb{Z}_{2k} -code is 24. Meanwhile, an extremal Type II \mathbb{Z}_{2k} -codes of length 24 are related to the Leech lattice, which is arguably the most remarkable lattice. In particular, by considering the existence of 2k-frames in the Leech lattice,

it was shown that there is an extremal Type II \mathbb{Z}_{2k} -code of length 24 for every positive integer k with $k \geq 2$ [5] and [12].

There are nine inequivalent binary doubly even self-dual codes of length 24 [25]. The seven codes are indecomposable and these code are denoted by A24, B24, ..., G24 in [25, Table II]. Only the code G24 has minimum weight 8, that is, G24 is extremal, and it is well known as the extended Golay code. As usual we denote the two decomposable codes by e_8^3 and $d_{16}e_8$. For every binary doubly even self-dual code B of length 24, there is an extremal Type II \mathbb{Z}_{2k} -code C such that $C^{(2)} \cong B$ when k=2 [14, Proposition 9] (see [4, Postscript]) and k=3 [20, Theorem 3]. In this section, we establish the following theorem by constructing extremal Type II \mathbb{Z}_{2k} -codes of length 24 for $k=4,5,\ldots,20$ by using four-negacirculant codes and by Theorem 3.4 explicitly.

Theorem 4.1. Suppose that $k \in \{2, 3, ..., 20\}$. For every binary doubly even self-dual code B of length 24, there is an extremal Type II \mathbb{Z}_{2k} -code C such that $C^{(2)} \cong B$.

Although the following lemma is trivial, we give a proof for completeness.

Lemma 4.2. Suppose that $k \geq 3$ and $n \in \{24, 32, 40\}$. Let C be a Type II \mathbb{Z}_{2k} -code of length n. Let $A_{2k}(C)$ denote the even unimodular lattice given in (4). Then C is extremal if and only if $A_{2k}(C)$ has minimum norm 4.

Proof. Since $k \geq 3$, a vector of norm 2 of $A_{2k}(C)$ is written as

$$\frac{1}{\sqrt{2k}}(\rho(c_1),\rho(c_2),\ldots,\rho(c_n))$$

for some codeword $c=(c_1,c_2,\ldots,c_n)\in C$ such that $\operatorname{wt}_E(c)=4k$, where ρ is the map given in (2). Thus, C contains no codeword of Euclidean weight 4k if and only if $A_{2k}(C)$ contains no vector of norm 2. It is known that the minimum norm $\min(L)$ of an even unimodular lattice in dimension n is bounded by $\min(L) \leq 2\lfloor \frac{n}{24} \rfloor + 2$ (see [7, Chapter 7]). Thus, $\min(A_{2k}(C)) \leq 4$ for $n \in \{24, 32, 40\}$. Hence, C contains no codeword of Euclidean weight 4k if and only if $A_{2k}(C)$ has minimum norm 4. The result follows.

By the above lemma, we determined the extremality for all Type II \mathbb{Z}_{2k} codes found in this section and the next section.

4.2 Extremal Type II \mathbb{Z}_8 -codes of length 24

By considering four-negacirculant codes, our computer search found extremal Type II \mathbb{Z}_8 -codes $C_{8,24,1}$ and $C_{8,24,2}$ of length 24 (see the end of Section 2 for four-negacirculant codes). The codes $C_{8,24,1}$ and $C_{8,24,2}$ have generator matrices of form (3), where the first rows r_A and r_B of the negacirculant matrices A and B are as follows

$$(r_A, r_B) = ((5, 3, 3, 3, 3, 1), (5, 5, 5, 6, 7, 7))$$
 and $((0, 0, 0, 1, 1, 3), (0, 1, 0, 3, 3, 1)),$

respectively. By Theorem 3.4, our computer search found more extremal Type II \mathbb{Z}_8 -codes $C_{8,24,3}, C_{8,24,4}, \ldots, C_{8,24,8}$ and $C_{8,24,9}$ as C(A, x, y) from $M_{8,24,i}$ (i = 1, 2, 4), where $M_{8,24,i}$ denotes the right half of the generator matrix of $C_{8,24,i}$. In Table 1, we list the matrices A and the vectors x, y in Theorem 3.4.

Codes (0,0,0,1,1,2,4,6,0,7,2,1) $C_{8,24,3}$ (1,1,1,4,7,3,3,0,2,5,5,6) $M_{8,24,1}$ $C_{8,24,4}$ $M_{8,24,1}$ (5,4,1,4,1,1,3,1,3,0,7,4)(0, 1, 1, 1, 7, 4, 6, 6, 1, 3, 3, 7) $C_{8,24,5}$ $M_{8,24,1}$ (6,5,1,4,5,5,6,6,4,6,0,2)(0,0,0,5,2,2,2,5,6,2,3,1) $C_{8,24,6}$ $M_{8,24,2}$ (3,3,3,6,1,0,3,5,1,2,5,4)(0,0,0,3,1,6,4,5,2,1,4,2) $C_{8,24,7}$ $M_{8,24,2}$ (4,3,7,0,0,0,6,2,6,0,1,5)(0, 1, 6, 1, 2, 4, 5, 6, 0, 0, 0, 5) $C_{8,24,8}$ $M_{8,24,2}$ (4, 1, 1, 7, 7, 5, 2, 3, 0, 3, 5, 6)(0,0,2,3,0,2,2,0,3,3,0,3)(5, 2, 2, 6, 6, 2, 6, 1, 5, 2, 1, 0)(0,0,1,6,5,0,6,0,1,0,3,2) $M_{8,24,4}$ $C_{8,24,9}$

Table 1: Extremal Type II \mathbb{Z}_8 -codes of length 24

4.3 Extremal Type II \mathbb{Z}_{2k} -codes of length **24** for $k = 5, 6, \dots, 20$

For \mathbb{Z}_{2k} -codes $(k = 5, 6, \dots, 20)$, to save space, we list the results only. Note that the approach is similar to that in the previous subsection.

For k = 5, 6, ..., 20, by considering four-negacirculant codes, our computer search found extremal Type II \mathbb{Z}_{2k} -codes $C_{2k,24,i}$ of length 24 $(i = 1, 2, ..., i_k)$, where $i_k = 3$ if $k \in \{5, 6, ..., 10\}$ and $i_k = 4$ if $k \in \{11, 12, ..., 20\}$ The codes have generator matrices of form (3), where the first rows r_A and r_B of the negacirculant matrices A and B are listed in Table 7. By Theorem 3.4, our computer search found more extremal Type II \mathbb{Z}_{2k} -codes $C_{2k,24,j}$ $(j = i_k + 1, i_k + 2, ..., 9)$ as C(A, x, y) from $M_{2k,24,i}$, where $M_{2k,24,i}$ denotes

the right half of the generator matrix of $C_{2k,24,i}$ (i = 1, 2, ..., 9). In Tables 8, 9 and 10, we list the matrices A and the vectors x, y in Theorem 3.4.

4.4 Binary parts of $C_{2k,24,1}, C_{2k,24,2}, \ldots, C_{2k,24,9}$ for $k = 4, 5, \ldots, 20$

For a given binary doubly even self-dual code B of length 24 and a given k, we list in Table 2 the number i such that $C_{2k,24,i}^{(2)} \cong B$. From Table 2, we have Theorem 4.1 combining with known results in [14] and [20].

- Remark 4.3. (i) For $B = e_8^3$ and G24, an extremal Type II \mathbb{Z}_8 -code C of length 24 such that $C^{(2)} \cong B$ can be found in [11] and [13], respectively.
 - (ii) Extremal Type II \mathbb{Z}_{10} -codes C of length 24 such that $C^{(2)} \cong e_8^3$ can be found in [11].
- (iii) For B = E24 and e_8^3 , an extremal Type II \mathbb{Z}_{12} -code C of length 24 such that $C^{(2)} \cong B$ can be found in [11].
- (iv) An extremal Type II \mathbb{Z}_{14} -code C of length 24 such that $C^{(2)} \cong e_8^3$ can be found in [11].
- (v) An extremal Type II \mathbb{Z}_{16} -code C of length 24 such that $C^{(2)} \cong E24$ can be found in [11].
- (vi) An extremal Type II \mathbb{Z}_{22} -code C of length 24 such that $C^{(2)} \cong E24$ can be found in [12]. For B = E24 and G24, an extremal Type II \mathbb{Z}_{22} -code C of length 24 such that $C^{(2)} \cong B$ can be found in [11].

In addition, by Lemma 2.2, we have the following corollary.

Corollary 4.4. Suppose that $k \in \{2, 3, ..., 20\}$. Then there are at least nine inequivalent extremal Type II \mathbb{Z}_{2k} -codes of length 24.

We end this section with the following question.

Question 1. Suppose that $k \geq 21$. For every binary doubly even self-dual code B of length 24, is there an extremal Type II \mathbb{Z}_{2k} -code C such that $C^{(2)} \cong B$.

			J 1		- 2n,2	4,6	, - ,	,	- /
\overline{k}	A24	B24	C24	D24	E24	F24	G24	e_8^3	$d_{16}e_{8}$
4	3	8	5	6	1	2	7	4	9
5	3	6	5	4	2	9	7	1	8
6	3	5	6	4	2	7	9	1	8
7	3	4	7	6	2	8	9	1	5
8	4	7	6	5	2	3	8	1	9
9	3	5	4	8	2	7	9	1	6
10	3	5	7	4	2	8	9	1	6
11	3	5	8	7	2	4	9	1	6
12	2	8	6	7	3	4	9	1	5
13	4	5	8	6	3	2	9	1	7
14	2	7	9	5	4	3	8	1	6
15	2	5	7	6	4	3	9	1	8
16	2	5	7	8	4	3	9	1	6
17	2	7	5	8	3	4	9	1	6
18	2	5	8	6	3	4	9	1	7
19	2	6	5	8	4	3	9	1	7

Table 2: Binary parts of $C_{2k,24,i}$ (k = 4, 5, ..., 20)

5 Extremal Type II \mathbb{Z}_{2k} -codes of length 32

8

5

20

For k = 4, 5, ..., 10, in this section, we construct extremal Type II \mathbb{Z}_{2k} -codes of length 32, by four-negacirculant codes and by Theorem 3.4.

4

3

9

1

6

There are five inequivalent binary extremal doubly even self-dual codes of length 32 [6]. The five codes are denoted by C81, C82, ..., C85 in [6, Table A]. For every binary extremal doubly even self-dual code B of length 32, there is an extremal Type II \mathbb{Z}_{2k} -code C such that $C^{(2)} \cong B$ when k = 2 [10, Theorem 5] and k = 3 [15, Proposition 3].

In this section, by constructing extremal Type II \mathbb{Z}_{2k} -codes of length 32 (k = 4, 5, ..., 10) by using four-negacirculant codes and by Theorem 3.4, we explicitly establish the following theorem.

Theorem 5.1. Suppose that $k \in \{2, 3, ..., 10\}$. For every binary extremal doubly even self-dual code B of length 32, there is an extremal Type II \mathbb{Z}_{2k} -code C such that $C^{(2)} \cong B$.

The approach is similar to that in the previous section. To save space, we list the results only. By considering four-negacirculant codes, our computer

Table 3: Extremal Type II four-negacirculant \mathbb{Z}_{2k} -codes of length 32 (k = 4, 5, ..., 10)

Codes	r_A	r_B
$C_{8,32,1}$	(7,4,3,6,3,5,6,3)	(5,6,0,2,0,4,7,4)
$C_{8,32,2}$	(7,4,3,6,4,5,2,5)	(6,5,2,2,0,7,1,2)
$C_{10,32,1}$	(5,1,4,5,1,8,0,4)	(4,6,0,0,2,3,1,5)
$C_{10,32,2}$	(5, 1, 4, 5, 4, 0, 5, 0)	(6,6,5,2,8,4,9,7)
$C_{12,32,1}$	(2,5,11,3,1,7,6,10)	(5, 10, 5, 6, 8, 6, 0, 4)
$C_{12,32,2}$	(2, 5, 11, 3, 3, 4, 10, 2)	(0,6,10,1,2,0,5,1)
$C_{14,32,1}$	(3, 11, 6, 1, 3, 10, 13, 12)	(4, 10, 3, 0, 7, 12, 6, 8)
$C_{14,32,2}$	(3, 11, 6, 1, 2, 2, 11, 8)	(12, 0, 2, 13, 1, 10, 6, 9)
$C_{16,32,1}$	(2, 13, 4, 2, 11, 2, 0, 15)	(4, 15, 9, 0, 2, 0, 15, 11)
$C_{16,32,2}$	(2, 13, 4, 2, 14, 7, 7, 0)	(6, 12, 7, 7, 5, 1, 6, 10)
$C_{18,32,1}$	(14, 17, 15, 11, 1, 3, 14, 2)	(12, 10, 3, 4, 5, 6, 4, 4)
$C_{18,32,2}$	(14, 17, 15, 11, 5, 16, 16, 14)	(8, 16, 5, 6, 0, 3, 15, 14)
$C_{20,32,1}$	(2,0,8,9,15,5,11,7)	(10, 0, 15, 8, 5, 4, 8, 16)
$C_{20,32,2}$	(2,0,8,9,1,14,12,1)	(6, 18, 0, 1, 1, 19, 17, 16)

search found extremal Type II \mathbb{Z}_{2k} -codes $C_{2k,32,1}$ and $C_{2k,32,2}$ of length 32 $(k=4,5,\ldots,10)$. The codes have generator matrices of form (3), where the first rows r_A and r_B of the negacirculant matrices A and B are listed in Table 3. By Theorem 3.4, our computer search found more extremal Type II \mathbb{Z}_{2k} -codes $C_{2k,32,3}$, $C_{2k,32,4}$ and $C_{2k,32,5}$ as C(A,x,y) from $M_{2k,32,i}$, where $M_{2k,32,i}$ denotes the right half of the generator matrix of $C_{2k,32,i}$ $(i=1,2,k=4,5,\ldots,10)$. In Table 11, we list the matrices A and the vectors x,y in Theorem 3.4. For a given binary extremal doubly even self-dual code B of length 32 and a given k, we list in Table 4 the number i such that $C_{2k,32,i}^{(2)} \cong B$ $(k=4,5,\ldots,10)$. From Table 4, we have Theorem 5.1 combining with known results in [10] and [15]. In addition, by Lemma 2.2, we have the following corollary.

Corollary 5.2. Suppose that $k \in \{2, 3, ..., 10\}$. Then there are at least five inequivalent extremal Type II \mathbb{Z}_{2k} -codes of length 32.

We end this section with the following question.

Question 2. Suppose that $k \geq 11$. For every binary extremal doubly even self-dual code B of length 32, is there an extremal Type II \mathbb{Z}_{2k} -code C such that $C^{(2)} \cong B$.

Table 4: Binary parts of $C_{2k,32,i}$ (k = 4, 5, ..., 10)

\overline{k}	C81	C82	C83	C84	C85
4	2	4	1	3	5
5	2	4	1	5	3
6	2	5	1	4	3
7	2	4	1	3	5
8	2	4	1	5	3
9	2	4	1	5	3
10	2	5	1	3	4

6 Extremal Type II \mathbb{Z}_4 -codes of lengths 56 and 64

In this section, by four-negacirculant codes and by Theorem 3.4, we construct new extremal Type II \mathbb{Z}_4 -codes of lengths 56 and 64.

6.1 Extremality

We describe how to determine the extremality of a given Type II \mathbb{Z}_4 -code of lengths 48,56 and 64. Although the following lemma is trivial, we give a proof for completeness.

Lemma 6.1. Suppose that $n \in \{48, 56, 64\}$. Let C be a Type II \mathbb{Z}_4 -code of length n. Let $A_4(C)$ denote the even unimodular lattice given in (4). Then C is extremal if and only if $A_4(C)$ has minimum norm 4 and kissing number 2n.

Proof. It is trivial that C has a codeword of Euclidean weight 8 if and only if $A_4(C)$ has a vector of norm 2. If C has a codeword of Euclidean weight 16, then $A_4(C)$ has a vector of norm 4. In addition, $A_4(C)$ has 2n vectors of norm 4, which have the following form

$$(\pm 2, 0, 0, \dots, 0), (0, \pm 2, 0, \dots, 0), \dots, (0, 0, \dots, 0, \pm 2).$$

Hence, $A_4(C)$ has minimum norm 4 and kissing number 2n if and only if C has no codeword of Euclidean weight 16.

By the above lemma, we determined the extremality for all Type II \mathbb{Z}_4 -codes found in this section.

6.2 Extremal Type II \mathbb{Z}_4 -codes of length 64

The first example of an extremal Type II \mathbb{Z}_4 -code of length 64 was found in [16]. Note that the binary part of the code is not self-dual.

By considering four-negacirculant codes, our computer search found four extremal Type II \mathbb{Z}_4 -codes $C_{4,64,i}$ (i=1,2,3,4) of length 64. These codes have generator matrices of form (3), where the first rows r_A and r_B of the negacirculant matrices A and B are listed in Table 5.

Table 5: Extremal Type II four-negacirculant \mathbb{Z}_4 -codes of length 64

Codes	r_A	r_B
$C_{4,64,1}$	$ \begin{array}{c} (1,0,1,1,3,3,2,3,3,2,2,3,3,2,3,2) \\ (3,0,3,2,3,0,3,2,0,3,0,1,2,2,0,0) \\ (3,0,0,0,2,0,1,0,2,3,2,2,3,0,1,1) \end{array} $	(2,0,1,1,3,1,2,3,3,2,3,1,2,0,3,0)
$C_{4,64,2}$	(3,0,3,2,3,0,3,2,0,3,0,1,2,2,0,0)	(0, 2, 0, 0, 1, 2, 3, 2, 1, 2, 0, 0, 3, 0, 2, 1)
$C_{4,64,3}$	(3,0,0,0,2,0,1,0,2,3,2,2,3,0,1,1)	(1, 2, 1, 3, 1, 3, 2, 0, 0, 3, 3, 2, 1, 0, 1, 2)
$C_{4,64,4}$	(0,0,3,3,3,1,1,1,0,1,0,3,0,3,3,2)	(1,0,1,2,1,2,2,0,3,1,2,3,1,0,3,3)

In addition, by Theorem 3.4, our computer search found one more extremal Type II \mathbb{Z}_4 -code $C_{4,64,5}$ as $C(M_{4,64,2}, x, y)$, where $M_{4,64,2}$ denotes the right half of the generator matrix of $C_{4,64,2}$, and

$$x = (1, 1, 3, 0, 1, 3, 1, 0, 3, 2, 3, 0, 2, 2, 1, 1, 1, 2, 1, 3, 2, 2, 1, 0, 0, 0, 0, 0, 0, 0, 1, 1),$$

 $y = (1, 3, 2, 0, 0, 3, 2, 2, 0, 2, 0, 3, 1, 2, 1, 0, 2, 0, 2, 2, 0, 2, 2, 3, 1, 1, 1, 1, 2, 1, 2, 2).$

By Gleason's theorem (see [24]), the weight enumerator of a binary doubly even self-dual code of length 64 and minimum weight at least 8 is written as

$$W_{64}(a) = 1 + ay^8 + (2976 + 20a)y^{12} + (454956 + 2a)y^{16} + \cdots,$$

where a is the number of codewords of weight 8. Our computer search verified that the binary parts $C_{4,64,i}^{(2)}$ (i=1,2,3,4,5) have weight enumerators $W_{64}(16)$, $W_{64}(64)$, $W_{64}(0)$, $W_{64}(0)$, $W_{64}(19)$, respectively. In addition, our computer search verified that the binary parts $C_{4,64,3}^{(2)}$ and $C_{4,64,4}^{(2)}$ are inequivalent. By Lemma 2.2, we have the following proposition.

Proposition 6.2. There are at least six inequivalent extremal Type II \mathbb{Z}_4 -codes of length 64.

6.3 Extremal Type II \mathbb{Z}_4 -codes of length 56

The first example of an extremal Type II \mathbb{Z}_4 -code of length 56 was found in [16]. Note that the binary part of the code is not self-dual. Two more extremal Type II \mathbb{Z}_4 -codes of length 56 were constructed by considering double circulant codes in [17]. The two codes are denoted by $\mathcal{D}_{56,1}$ and \mathcal{C}_{56} in [17]. Note that the binary parts of these codes are binary extremal doubly even self-dual codes.

By considering four-negacirculant codes, our computer search found three extremal Type II \mathbb{Z}_4 -codes $C_{4,56,i}$ (i=1,2,3) of length 56. These codes have generator matrices of form (3), where the first rows r_A and r_B of the negacirculant matrices A and B are listed in Table 6.

By Gleason's theorem (see [24]), the weight enumerator of a binary doubly even self-dual code of length 56 and minimum weight at least 8 is written as

$$W_{56}(a) = 1 + ay^8 + (8190 + 6a)y^{12} + (622314 - 83a)y^{16} + \cdots$$

where a is the number of codewords of weight 8. Our computer search verified that the binary parts $C_{4,56,i}^{(2)}$ (i=1,2,3) have weight enumerators $W_{56}(35)$, $W_{56}(42)$, $W_{56}(0)$, respectively. In addition, our computer search verified that $\mathcal{D}_{56,1}^{(2)}$, $\mathcal{C}_{56}^{(2)}$ and $C_{4,56,3}^{(2)}$ are inequivalent. By Lemma 2.2, we have the following proposition.

Proposition 6.3. There are at least six inequivalent extremal Type II \mathbb{Z}_4 -codes of length 56.

Table 6: Extremal Type II four-negacirculant \mathbb{Z}_4 -codes of length 56

Codes	r_A	r_B
$C_{4,56,1}$	(2,2,0,1,2,0,2,0,2,2,1,1,2,1)	(0,0,3,3,2,0,3,1,3,1,1,0,0,2)
$C_{4,56,2}$	(2, 2, 2, 0, 1, 0, 1, 0, 1, 2, 3, 0, 3, 1)	(2,3,0,0,0,1,3,2,1,2,0,2,2,1)
$C_{4,56,3}$	(1,2,2,0,3,2,2,3,1,1,1,3,1,0)	(2,1,3,1,1,2,2,2,0,1,3,3,0,0)

By an approach is similar to that used in the previous subsection, we tried to construct a new extremal Type II \mathbb{Z}_4 -code of length 56 by Theorem 3.4. However, our extensive search failed to construct such a code.

Acknowledgments. This work was supported by JSPS KAKENHI Grant Number 19H01802. The author would like to thank the anonymous reviewers for useful comments.

References

- [1] E. Bannai, S.T. Dougherty, M. Harada and M. Oura, Type II codes, even unimodular lattices, and invariant rings, *IEEE Trans. Inform. Theory* **45** (1999), 1194–1205.
- [2] A. Bonnecaze, P. Solé, C. Bachoc and B. Mourrain, Type II codes over Z₄, IEEE Trans. Inform. Theory 43 (1997), 969–976.
- [3] W. Bosma, J. Cannon and C. Playoust, The Magma algebra system I: The user language, J. Symbolic Comput. 24 (1997), 235–265.
- [4] A.R. Calderbank and N.J.A. Sloane, Double circulant codes over \mathbb{Z}_4 and even unimodular lattices, *J. Algebraic Combin.* **6** (1997), 119–131.
- [5] R. Chapman, Double circulant constructions of the Leech lattice, J. Austral. Math. Soc. Ser. A 69 (2000), 287–297.
- [6] J.H. Conway, V. Pless and N.J.A. Sloane, The binary self-dual codes of length up to 32: a revised enumeration, J. Combin. Theory Ser. A 60 (1992), 183–195.
- [7] J.H. Conway and N.J.A. Sloane, Sphere Packing, Lattices and Groups (3rd ed.), Springer-Verlag, New York, 1999.
- [8] S.T. Dougherty, T.A. Gulliver and M. Harada, Type II self-dual codes over finite rings and even unimodular lattices, J. Algebraic Combin. 9 (1999), 233–250.
- [9] S.T. Dougherty, M. Harada and P. Solé, Self-dual codes over rings and the Chinese remainder theorem, *Hokkaido Math. J.* **28** (1999), 253–283.
- [10] P. Gaborit and M. Harada, Construction of extremal Type II codes over \mathbb{Z}_4 , Des. Codes Cryptogr. **16** (1999), 257–269.
- [11] S. Georgiou, M. Harada and C. Koukouvinos, Orthogonal designs and Type II codes over \mathbb{Z}_{2k} , Des. Codes Cryptogr. **25** (2002), 163–174.
- [12] T.A. Gulliver and M. Harada, Orthogonal frames in the Leech lattice and a Type II code over \mathbb{Z}_{22} , J. Combin. Theory Ser. A **95** (2001), 185–188.

- [13] T.A. Gulliver and M. Harada, Extremal self-dual codes over \mathbb{Z}_6 , \mathbb{Z}_8 and \mathbb{Z}_{10} , AKCE Int. J. Graphs Comb. 2 (2005), 11–24.
- [14] M. Harada, New extremal Type II codes over \mathbb{Z}_4 , Des. Codes Cryptogr. **13** (1998), 271–284.
- [15] M. Harada, On the existence of extremal Type II codes over \mathbb{Z}_6 , Discrete Math. **223** (2000), 373–378.
- [16] M. Harada, Extremal Type II \mathbb{Z}_4 -codes of lengths 56 and 64, J. Combin. Theory Ser. A 117 (2010), 1285–1288.
- [17] M. Harada, Note on the residue codes of self-dual \mathbb{Z}_4 -codes having large minimum Lee weights, $Adv.\ Math.\ Commun.\ 10\ (2016),\ 695-706.$
- [18] M. Harada, W. Holzmann, H. Kharaghani and M. Khorvash, Extremal ternary self-dual codes constructed from negacirculant matrices, *Graphs Combin.* **23** (2007), 401–417.
- [19] M. Harada and H. Kimura, New extremal doubly-even [64, 32, 12] codes, Des. Codes Cryptogr. 6 (1995), 91–96.
- [20] M. Harada and M. Kitazume, \mathbb{Z}_6 -code constructions of the Leech lattice and the Niemeier lattices, European J. Combin. 23 (2002), 573–581.
- [21] M. Harada and T. Miezaki, An upper bound on the minimum weight of Type II \mathbb{Z}_{2k} -codes, J. Combin. Theory Ser. A **118** (2011), 190–196.
- [22] M. Harada, P. Solé and P. Gaborit, Self-dual codes over \mathbb{Z}_4 and unimodular lattices: a survey, Algebras and combinatorics (Hong Kong, 1997), 255–275, Springer, Singapore, 1999.
- [23] K. Ishizuka and K. Saito, Construction for both self-dual codes and LCD codes, Adv. Math. Commun., (to appear), arXiv: 2108.12544.
- [24] C.L. Mallows and N.J.A. Sloane, An upper bound for self-dual codes, Inform. Control 22 (1973), 188–200.
- [25] V. Pless and N.J.A. Sloane, On the classification and enumeration of self-dual codes, *J. Combin. Theory Ser. A* **18** (1975), 313–335.

- [26] E. Rains and N.J.A. Sloane, "Self-dual codes," Handbook of Coding Theory, V.S. Pless and W.C. Huffman (Editors), Elsevier, Amsterdam 1998, pp. 177–294.
- [27] V.D. Tonchev, Self-orthogonal designs and extremal doubly even codes, J. Combin. Theory Ser. A 52 (1989), 197–205.

Table 7: Extremal Type II four-negacirculant \mathbb{Z}_{2k} -codes of length 24 $(k = 5, 12, \dots, 20)$

Codes	r_A	r_B	Codes	r_A	r_B
$C_{10,24,1}$	(9, 8, 2, 5, 4, 2)	(3, 8, 6, 0, 6, 0)	$C_{26,24,4}$	(14, 13, 16, 2, 8, 20)	(7,7,7,11,11,9)
$C_{10,24,2}$	(1,7,3,1,9,3)	(7, 5, 3, 1, 1, 8)	$C_{28,24,1}$	(15, 8, 10, 7, 6, 8)	(22, 16, 9, 16, 22, 14)
$C_{10,24,3}$	(5, 8, 1, 6, 5, 0)	(5,7,2,5,2,1)	$C_{28,24,2}$	(2, 23, 20, 27, 24, 1)	(22, 3, 0, 3, 21, 25)
$C_{12,24,1}$	(6,4,8,8,5,6)	(10, 9, 0, 2, 5, 10)	$C_{28,24,3}$	(13, 15, 1, 18, 18, 20)	(3, 7, 5, 22, 9, 22)
$C_{12,24,2}$	(3,7,9,7,11,11)	(9, 9, 2, 7, 11, 1)	$C_{28,24,4}$	(17, 5, 3, 13, 25, 14)	(21, 1, 5, 1, 27, 25)
$C_{12,24,3}$	(6, 9, 8, 7, 4, 3)	(1,0,1,6,9,3)	$C_{30,24,1}$	(1, 8, 28, 9, 12, 18)	(6, 8, 4, 2, 1, 20)
$C_{14,24,1}$	(2,6,11,4,0,5)	(6,4,6,0,8,11)	$C_{30,24,2}$	(29, 28, 23, 13, 21, 14)	(22, 11, 4, 3, 0, 27)
$C_{14,24,2}$	(11, 9, 13, 7, 9, 5)	(10, 5, 9, 7, 3, 7)	$C_{30,24,3}$	(29, 28, 23, 13, 21, 14)	(22, 11, 7, 21, 18, 0)
$C_{14,24,3}$	(7, 8, 11, 3, 11, 12)	(13, 10, 1, 6, 5, 0)	$C_{30,24,4}$	(18, 19, 29, 5, 23, 11)	(27, 1, 1, 3, 27, 23)
$C_{16,24,1}$	(13, 4, 2, 11, 4, 4)	(15, 8, 14, 0, 8, 14)	$C_{32,24,1}$	(2, 16, 16, 4, 29, 2)	(18, 6, 1, 2, 0, 7)
$C_{16,24,2}$	(1, 3, 13, 15, 5, 14)	(5, 9, 3, 7, 9, 5)	$C_{32,24,2}$	(16, 5, 16, 21, 22, 5)	(27, 27, 0, 29, 2, 7)
$C_{16,24,3}$	(8,9,8,3,15,1)	(6,0,2,3,15,9)	$C_{32,24,3}$	(14, 11, 19, 17, 30, 0)	(29, 13, 2, 27, 6, 1)
$C_{18,24,1}$	(3, 2, 8, 15, 8, 14)	(16, 5, 2, 14, 0, 0)	$C_{32,24,4}$	(15, 3, 31, 23, 29, 27)	(27, 10, 13, 3, 19, 3)
$C_{18,24,2}$	(15, 3, 8, 17, 13, 5)	(17, 3, 17, 1, 3, 5)	$C_{34,24,1}$	(16, 19, 30, 22, 25, 14)	(0, 3, 4, 32, 2, 0)
$C_{18,24,3}$	(5,9,13,17,5,7)	(4, 10, 2, 8, 2, 1)	$C_{34,24,2}$	(33, 13, 25, 6, 33, 8)	(11, 12, 1, 12, 33, 16)
$C_{20,24,1}$	(17, 10, 16, 5, 10, 4)	(12, 11, 12, 8, 14, 8)	$C_{34,24,3}$	(4, 25, 3, 27, 15, 13)	(3, 29, 9, 5, 31, 27)
$C_{20,24,2}$	(13, 3, 17, 17, 3, 17)	(5, 2, 15, 5, 15, 1)	$C_{34,24,4}$	(18, 7, 9, 17, 10, 18)	(21, 10, 3, 24, 23, 13)
$C_{20,24,3}$	(1, 9, 8, 17, 2, 13)	(5,6,1,4,13,8)	$C_{36,24,1}$	(20, 7, 24, 22, 11, 6)	(2, 8, 0, 7, 4, 24)
$C_{22,24,1}$	(11, 0, 10, 12, 10, 14)	(17, 2, 0, 19, 6, 10)	$C_{36,24,2}$	(2,11,34,35,6,1)	(11, 2, 1, 25, 23, 8)
$C_{22,24,2}$	(7, 21, 11, 5, 2, 3)	(21, 7, 1, 7, 15, 13)	$C_{36,24,3}$	(15, 15, 25, 7, 3, 10)	(35, 5, 1, 33, 13, 19)
$C_{22,24,3}$	(16, 9, 14, 11, 9, 21)	(16, 17, 4, 5, 18, 5)	$C_{36,24,4}$	(21, 34, 13, 17, 17, 25)	(31, 20, 2, 31, 22, 22)
$C_{22,24,4}$	(1, 10, 7, 19, 5, 19)	(12, 15, 10, 6, 11, 4)	$C_{38,24,1}$	(16, 36, 16, 32, 25, 26)	(16, 29, 0, 22, 25, 14)
$C_{24,24,1}$	(7, 4, 6, 17, 8, 18)	(9, 8, 4, 10, 4, 12)	$C_{38,24,2}$	(32, 33, 18, 1, 20, 1)	(18, 7, 20, 19, 7, 9)
$C_{24,24,2}$	(0, 3, 16, 13, 22, 13)	(15, 5, 1, 0, 17, 2)	$C_{38,24,3}$	(17, 17, 27, 17, 26, 9)	(2, 2, 3, 34, 22, 13)
$C_{24,24,3}$	(9, 11, 5, 23, 11, 19)	(5, 19, 2, 21, 3, 15)	$C_{38,24,4}$	(29, 3, 4, 15, 17, 31)	(11, 3, 3, 33, 13, 17)
$C_{24,24,4}$	(7, 3, 16, 12, 14, 3)	(2, 11, 0, 19, 23, 23)	$C_{40,24,1}$	(34, 26, 39, 2, 38, 6)	(26, 10, 1, 0, 4, 23)
$C_{26,24,1}$	(17, 2, 14, 8, 8, 10)	(10, 15, 0, 0, 1, 10)	$C_{40,24,2}$	(25, 38, 31, 19, 37, 8)	(7, 20, 1, 0, 9, 22)
$C_{26,24,2}$	(15, 20, 5, 15, 17, 8)	(14, 0, 21, 11, 1, 12)	$C_{40,24,3}$	(5, 1, 32, 15, 30, 35)	(24, 25, 3, 3, 6, 28)
$C_{26,24,3}$	(1, 7, 14, 23, 5, 21)	(3, 25, 1, 11, 11, 13)	$C_{40,24,4}$	(37, 13, 15, 19, 27, 15)	(3, 15, 5, 31, 31, 10)

Table 8: Extremal Type II \mathbb{Z}_{2k} -codes of length 24 $(k = 5, 6, \dots, 9)$

Codes	A	x	y
$C_{10,24,4}$	$M_{10,24,1}$	(7,1,6,2,4,1,0,3,5,1,7,7)	(0,0,0,0,1,5,9,2,4,0,3,2)
$C_{10,24,5}$	$M_{10,24,1}$	(6,5,6,0,3,6,3,5,4,0,8,2)	(0,0,0,4,0,4,4,6,9,5,7,1)
$C_{10,24,6}$	$M_{10,24,1}$	(1,7,1,1,0,6,0,4,5,7,1,9)	(0,0,0,0,3,0,8,1,5,8,4,9)
$C_{10,24,7}$	$M_{10,24,4}$	(1,9,5,9,8,5,5,3,5,2,2,4)	(0,0,0,0,8,2,3,1,2,1,6,1)
$C_{10,24,8}$	$M_{10,24,6}$	(9,6,0,6,0,4,7,6,7,8,2,7)	(0,0,0,5,4,6,7,2,1,3,4,8)
$C_{10,24,9}$	$M_{10,24,7}$	(5,4,3,8,8,6,7,0,0,4,0,9)	(0,0,0,0,0,0,1,6,5,5,8,7)
$C_{12,24,4}$	$M_{12,24,1}$	(5,7,9,5,6,7,7,1,0,8,1,10)	(0,0,0,0,1,5,1,8,10,2,6,9)
$C_{12,24,5}$	$M_{12,24,1}$	(11, 4, 10, 8, 7, 8, 2, 3, 0, 0, 1, 10)	(0,0,0,0,10,3,11,10,5,11,6,4)
$C_{12,24,6}$	$M_{12,24,2}$	(8,4,3,9,8,10,0,5,2,8,7,10)	(0,0,0,0,9,2,0,0,7,3,8,3)
$C_{12,24,7}$	$M_{12,24,4}$	(7, 2, 11, 8, 2, 0, 2, 10, 6, 1, 5, 0)	(0,0,0,1,3,10,2,4,6,11,3,4)
$C_{12,24,8}$	$M_{12,24,5}$	(3,4,5,1,7,8,4,7,3,4,7,9)	(0,0,0,3,6,7,7,6,1,8,10,8)
$C_{12,24,9}$	$M_{12,24,7}$	(5,6,6,10,1,2,5,9,1,5,5,11)	(0,0,1,0,6,7,8,7,0,11,6,10)
$C_{14,24,4}$	$M_{14,24,1}$	(7, 8, 6, 6, 11, 2, 6, 8, 2, 1, 8, 5)	(0,0,0,0,9,10,13,0,6,7,11,12)
$C_{14,24,5}$	$M_{14,24,1}$	(0,9,6,7,12,8,5,7,4,12,0,8)	(0,0,0,0,11,10,1,3,2,2,9,10)
$C_{14,24,6}$	$M_{14,24,3}$	(4,5,13,0,6,7,1,2,1,3,1,5)	(0,0,0,0,7,8,4,13,10,7,7,6)
$C_{14,24,7}$	$M_{14,24,3}$	(4,11,12,13,8,13,10,2,1,4,8,0)	(0,0,0,0,6,9,11,0,7,6,2,3)
$C_{14,24,8}$	$M_{14,24,6}$	(6, 13, 6, 11, 7, 0, 5, 6, 1, 1, 3, 7)	(0,0,0,0,12,5,4,11,11,4,13,2)
$C_{14,24,9}$	$M_{14,24,8}$	(6,6,10,1,2,6,8,3,4,5,13,6)	(0,0,0,1,3,3,2,7,7,11,11,1)
$C_{16,24,4}$	$M_{16,24,1}$	(1,0,7,7,5,9,7,9,2,13,10,8)	(0,0,0,0,7,12,6,3,15,10,2,13)
$C_{16,24,5}$	$M_{16,24,1}$	(11, 14, 13, 10, 5, 11, 12, 5, 1, 8, 9, 13)	(0,0,0,3,0,7,9,7,14,0,12,12)
$C_{16,24,6}$	$M_{16,24,1}$	(4,11,3,10,13,7,0,1,1,2,7,11)	(0,0,0,0,10,14,9,2,0,5,5,7)
$C_{16,24,7}$	$M_{16,24,1}$	(14, 14, 15, 10, 2, 7, 15, 11, 2, 0, 0, 8)	(0,0,0,0,1,9,1,6,15,6,10,8)
$C_{16,24,8}$	$M_{16,24,3}$	(2,6,2,14,13,10,7,10,3,7,4,10)	(0,0,0,0,15,5,14,11,8,1,6,2)
$C_{16,24,9}$	$M_{16,24,7}$	(10, 14, 9, 14, 3, 14, 7, 1, 5, 13, 9, 7)	(0,0,0,0,9,4,3,9,12,14,1,12)
$C_{18,24,4}$	$M_{18,24,2}$	(2, 14, 14, 12, 17, 5, 7, 17, 1, 3, 7, 3)	(0,0,0,0,6,0,3,8,13,3,1,0)
$C_{18,24,5}$	$M_{18,24,3}$	(8,4,13,3,13,4,2,10,0,5,2,0)	(0,0,0,0,13,13,7,7,11,17,3,3)
$C_{18,24,6}$	$M_{18,24,3}$	(3, 15, 11, 15, 7, 16, 1, 8, 1, 14, 8, 11)	(0,0,0,0,6,13,12,9,12,5,11,0)
$C_{18,24,7}$	$M_{18,24,5}$	(11, 17, 9, 14, 12, 3, 11, 1, 0, 17, 6, 3)	(0,0,0,0,8,11,0,13,12,1,10,11)
$C_{18,24,8}$	$M_{18,24,5}$	(9, 16, 0, 14, 10, 1, 16, 10, 0, 3, 3, 6)	(0,0,0,0,13,5,10,14,5,5,6,6)
$C_{18,24,9}$	$M_{18,24,7}$	(4,3,0,9,0,6,0,3,4,2,3,12)	(0,0,0,0,4,3,0,1,4,7,11,2)

Table 9: Extremal Type II \mathbb{Z}_{2k} -codes of length 24 $(k=10,12,\ldots,15)$

G 1	1 4		
Codes	A	x	y
$C_{20,24,4}$	$M_{20,24,1}$	(19,7,14,7,1,7,4,10,0,3,17,9)	(0,0,0,0,14,1,10,11,15,0,14,19)
$C_{20,24,5}$	$M_{20,24,1}$	(16, 10, 16, 5, 7, 10, 15, 2, 0, 0, 17, 16)	(0,0,0,0,10,1,1,9,14,18,19,4)
$C_{20,24,6}$	$M_{20,24,1}$	(2, 1, 15, 10, 17, 11, 0, 18, 4, 2, 10, 4)	(0,0,0,0,5,15,9,14,2,17,8,14)
$C_{20,24,7}$	$M_{20,24,2}$	(11, 0, 17, 19, 0, 1, 0, 4, 0, 4, 0, 14)	(0,0,0,0,14,10,19,0,9,6,5,9)
$C_{20,24,8}$	$M_{20,24,3}$	(8,0,12,0,1,17,1,18,2,1,14,4)	(0,0,0,0,0,6,10,5,5,10,5,7)
$C_{20,24,9}$	$M_{20,24,4}$	(13, 6, 19, 6, 7, 3, 6, 4, 0, 2, 0, 2)	(0,0,0,1,8,14,5,8,9,3,2,14)
$C_{22,24,5}$	$M_{22,24,1}$	(18, 20, 15, 19, 8, 18, 1, 6, 0, 2, 4, 7)	(8, 12, 15, 21, 3, 20, 9, 17, 1, 14, 9, 15)
$C_{22,24,6}$	$M_{22,24,1}$	(20, 10, 4, 2, 16, 3, 0, 19, 2, 12, 11, 9)	(18, 3, 2, 18, 1, 13, 12, 10, 14, 14, 19, 14)
$C_{22,24,7}$	$M_{22,24,3}$	(21, 12, 11, 6, 20, 12, 1, 0, 0, 1, 18, 4)	(13, 11, 12, 6, 0, 10, 7, 9, 6, 4, 8, 8)
$C_{22,24,8}$	$M_{22,24,3}$	(19, 14, 3, 4, 15, 15, 7, 6, 0, 9, 17, 19)	(5,7,17,12,1,0,3,3,4,15,8,15)
$C_{22,24,9}$	$M_{22,24,4}$	(7, 18, 20, 4, 17, 17, 8, 15, 0, 8, 20, 16)	(2, 6, 20, 0, 1, 18, 4, 11, 3, 1, 0, 10)
$C_{24,24,5}$	$M_{24,24,1}$	(18, 21, 21, 18, 23, 9, 0, 0, 0, 0, 10, 8)	(10, 5, 1, 8, 8, 20, 12, 23, 4, 6, 21, 10)
$C_{24,24,6}$	$M_{24,24,1}$	(9, 3, 23, 3, 23, 14, 0, 0, 1, 1, 9, 10)	(0, 18, 21, 4, 2, 20, 14, 4, 17, 7, 5, 16)
$C_{24,24,7}$	$M_{24,24,2}$	(13, 21, 17, 12, 7, 11, 0, 0, 0, 5, 9, 19)	(10, 5, 21, 0, 19, 12, 0, 16, 20, 6, 3, 10)
$C_{24,24,8}$	$M_{24,24,2}$	(21, 16, 20, 22, 9, 13, 0, 0, 0, 2, 2, 15)	(5, 13, 8, 15, 5, 14, 3, 8, 7, 1, 9, 10)
$C_{24,24,9}$	$M_{24,24,4}$	(4, 10, 6, 3, 22, 3, 0, 0, 0, 0, 21, 21)	(6, 16, 15, 6, 20, 7, 20, 8, 17, 0, 18, 9)
$C_{26,24,5}$	$M_{26,24,1}$	(2, 11, 8, 18, 7, 22, 0, 0, 0, 0, 7, 7)	(13, 24, 16, 18, 2, 17, 22, 12, 16, 6, 2, 5, 3)
$C_{26,24,6}$	$M_{26,24,2}$	(9, 18, 16, 17, 23, 6, 0, 0, 0, 0, 19, 16)	(16, 20, 12, 23, 21, 11, 3, 23, 1, 1, 20, 17)
$C_{26,24,7}$	$M_{26,24,3}$	(15, 9, 2, 14, 11, 2, 9, 18, 0, 16, 24, 24)	(5, 9, 9, 21, 23, 13, 5, 13, 9, 13, 7, 25)
$C_{26,24,8}$	$M_{26,24,5}$	(20, 18, 16, 17, 4, 3, 0, 0, 0, 5, 7, 20)	(21, 1, 12, 4, 25, 24, 2, 16, 22, 8, 14, 25)
$C_{26,24,9}$	$M_{26,24,6}$	(20, 21, 2, 15, 15, 21, 0, 0, 0, 24, 12, 14)	(8,7,17,12,13,4,15,14,1,7,15,9)
$C_{28,24,5}$	$M_{28,24,1}$	(21, 21, 4, 10, 2, 8, 0, 0, 0, 1, 21, 2)	(2, 27, 26, 23, 23, 12, 23, 27, 11, 25, 20, 5)
$C_{28,24,6}$	$M_{28,24,1}$	(20, 19, 24, 13, 25, 12, 20, 4, 0, 10, 8, 13)	(13, 17, 10, 15, 8, 22, 20, 18, 11, 18, 24, 2)
$C_{28,24,7}$	$M_{28,24,2}$	(16, 26, 10, 6, 20, 27, 0, 0, 0, 3, 23, 17)	(0, 13, 0, 21, 22, 27, 26, 3, 17, 27, 25, 7)
$C_{28,24,8}$	$M_{28,24,3}$	(23, 20, 12, 18, 14, 26, 0, 0, 0, 3, 3, 11)	(4, 2, 27, 18, 20, 22, 18, 25, 14, 3, 17, 4)
$C_{28,24,9}$	$M_{28,24,4}$	(3, 16, 16, 15, 22, 7, 0, 0, 0, 0, 8, 13)	(16, 16, 22, 22, 15, 14, 5, 14, 20, 1, 25, 18)
$C_{30,24,5}$	$M_{30,24,1}$	(29, 29, 27, 4, 2, 17, 8, 29, 0, 3, 11, 5)	(2, 12, 9, 13, 26, 21, 2, 7, 13, 29, 19, 11)
$C_{30,24,6}$	$M_{30,24,2}$	(16, 18, 23, 3, 2, 13, 0, 0, 0, 0, 5, 8)	(16, 4, 14, 26, 27, 24, 2, 1, 23, 28, 18, 17)
$C_{30,24,7}$	$M_{30,24,2}$	(19, 18, 21, 18, 7, 22, 0, 0, 0, 1, 26, 10)	(14, 15, 28, 28, 26, 6, 9, 16, 7, 4, 4, 29)
$C_{30,24,8}$	$M_{30,24,2}$	(29, 9, 9, 4, 27, 2, 3, 25, 0, 11, 22, 3)	(3, 10, 29, 22, 19, 19, 22, 13, 25, 1, 6, 3)
$C_{30,24,9}$	$M_{30,24,6}$	(3, 1, 16, 3, 29, 26, 0, 0, 0, 14, 6, 4)	(6, 15, 25, 9, 16, 24, 1, 23, 21, 17, 15, 26)

Table 10: Extremal Type II \mathbb{Z}_{2k} -codes of length 24 $(k=16,17,\ldots,20)$

Codes	A	x	y
$C_{32,24,5}$	$M_{32,24,1}$	(18, 10, 25, 10, 3, 22, 23, 6, 0, 0, 28, 23)	(15, 8, 10, 23, 30, 10, 24, 4, 27, 20, 29, 0)
$C_{32,24,6}$	$M_{32,24,1}$	(14, 4, 9, 10, 22, 3, 31, 18, 0, 7, 14, 12)	(30, 4, 29, 22, 3, 31, 16, 16, 31, 12, 8, 10)
$C_{32,24,7}$	$M_{32,24,2}$	(21, 23, 19, 23, 3, 1, 23, 15, 0, 8, 8, 0)	(9, 5, 19, 4, 23, 24, 29, 13, 24, 9, 29, 0)
$C_{32,24,8}$	$M_{32,24,2}$	(27, 27, 7, 25, 0, 25, 28, 31, 0, 0, 9, 5)	(2, 13, 27, 27, 14, 16, 30, 6, 18, 4, 12, 9)
$C_{32,24,9}$	$M_{32,24,3}$	(22, 24, 21, 19, 31, 31, 0, 4, 0, 8, 22, 18)	(8, 18, 29, 12, 3, 10, 20, 0, 14, 28, 21, 5)
$C_{34,24,5}$	$M_{34,24,1}$	(3, 26, 20, 2, 1, 9, 20, 32, 0, 2, 18, 33)	(14, 31, 26, 5, 0, 32, 21, 25, 24, 10, 28, 10)
$C_{34,24,6}$	$M_{34,24,1}$	(3, 2, 32, 30, 30, 12, 5, 22, 0, 11, 20, 1)	(10, 10, 15, 19, 2, 8, 8, 26, 32, 13, 20, 3)
$C_{34,24,7}$	$M_{34,24,2}$	(32, 25, 16, 3, 16, 22, 27, 29, 0, 0, 12, 18)	(31, 16, 17, 3, 25, 17, 29, 8, 30, 9, 32, 13)
$C_{34,24,8}$	$M_{34,24,4}$	(15, 20, 3, 23, 6, 23, 6, 6, 0, 12, 20, 28)	(6, 26, 25, 22, 4, 7, 28, 6, 15, 14, 30, 23)
$C_{34,24,9}$	$M_{34,24,4}$	(13, 10, 26, 7, 11, 19, 6, 24, 1, 1, 23, 13)	(13, 13, 7, 12, 31, 26, 7, 31, 7, 22, 13, 8)
$C_{36,24,5}$	$M_{36,24,1}$	(35, 16, 10, 11, 0, 33, 20, 23, 0, 2, 2, 4)	(35, 25, 34, 0, 22, 11, 35, 0, 35, 11, 3, 19)
$C_{36,24,6}$	$M_{36,24,1}$	(28, 15, 24, 19, 13, 16, 8, 21, 0, 0, 12, 2)	(3, 9, 5, 33, 7, 24, 23, 3, 2, 21, 22, 32)
$C_{36,24,7}$	$M_{36,24,1}$	(34, 25, 25, 32, 20, 4, 27, 20, 0, 0, 31, 20)	(8, 8, 0, 26, 31, 24, 33, 4, 34, 35, 11, 26)
$C_{36,24,8}$	$M_{36,24,3}$	(19, 29, 26, 9, 30, 6, 30, 18, 0, 1, 2, 22)	(3, 29, 31, 29, 20, 3, 13, 10, 28, 33, 10, 31)
$C_{36,24,9}$	$M_{36,24,4}$	(4, 28, 15, 7, 33, 26, 13, 19, 1, 1, 2, 21)	(33, 9, 20, 4, 6, 32, 17, 24, 15, 12, 22, 10)
$C_{38,24,5}$	$M_{38,24,1}$	(18, 5, 17, 23, 6, 5, 29, 1, 0, 0, 23, 21)	(3, 33, 36, 1, 15, 11, 8, 12, 19, 17, 10, 33)
$C_{38,24,6}$	$M_{38,24,1}$	(1, 22, 16, 12, 19, 14, 31, 19, 0, 0, 36, 24)	(31, 5, 17, 28, 10, 21, 18, 11, 35, 25, 5, 20)
$C_{38,24,7}$	$M_{38,24,1}$	(34, 16, 34, 7, 37, 2, 17, 19, 0, 0, 6, 32)	(21, 4, 28, 35, 26, 34, 9, 30, 36, 11, 0, 26)
$C_{38,24,8}$	$M_{38,24,3}$	(2, 22, 1, 1, 37, 5, 18, 9, 0, 1, 19, 35)	(30, 11, 12, 23, 11, 32, 24, 17, 29, 35, 19, 33)
$C_{38,24,9}$	$M_{38,24,3}$	(4, 28, 24, 35, 37, 13, 3, 34, 0, 0, 28, 12)	(30, 18, 10, 10, 13, 32, 15, 36, 7, 6, 23, 32)
$C_{40,24,5}$	$M_{40,24,1}$	(11, 26, 7, 16, 35, 33, 17, 9, 0, 1, 7, 2)	(24, 36, 27, 6, 1, 1, 34, 18, 10, 7, 16, 34)
$C_{40,24,6}$	$M_{40,24,1}$	(12, 2, 17, 30, 8, 28, 22, 39, 0, 0, 9, 17)	(32, 38, 35, 4, 28, 9, 30, 39, 19, 30, 12, 20)
$C_{40,24,7}$	$M_{40,24,2}$	(19, 20, 22, 15, 5, 23, 15, 9, 0, 1, 10, 33)	(3, 1, 29, 34, 19, 21, 1, 1, 28, 15, 28, 6)
$C_{40,24,8}$	$M_{40,24,3}$	(6, 39, 12, 39, 21, 27, 1, 32, 1, 1, 15, 34)	(35, 29, 21, 13, 5, 37, 28, 14, 23, 1, 28, 34)
$C_{40,24,9}$	$M_{40,24,3}$	(23, 16, 0, 28, 5, 9, 22, 22, 0, 2, 32, 27)	(7, 29, 11, 38, 30, 21, 8, 12, 26, 10, 22, 24)

Table 11: Extremal Type II \mathbb{Z}_{2k} -codes of length 32 $(k=4,5,\ldots,10)$

Codes	A	x	y
$C_{8,32,3}$	$M_{8,32,1}$	(0,0,7,6,4,1,1,3,5,0,3,0,0,1,3,2)	(0,0,0,0,3,7,2,2,7,1,0,4,2,4,6,2)
$C_{8,32,4}$	$M_{8,32,1}$	(5,5,0,5,0,1,6,3,1,3,3,0,1,1,1,1)	(0,0,6,3,4,7,1,6,4,3,4,5,5,7,1,4)
$C_{8,32,5}$	$M_{8,32,2}$	(6,0,2,4,4,1,3,6,4,5,1,0,0,0,0,4)	(0,0,0,0,0,1,3,7,5,1,7,4,0,1,4,5)
$C_{10,32,3}$	$M_{10,32,1}$	(4, 1, 3, 6, 5, 4, 2, 6, 6, 3, 0, 6, 1, 1, 3, 5)	(0,0,0,0,2,7,8,7,4,1,0,8,0,6,6,9)
$C_{10,32,4}$	$M_{10,32,1}$	(3,3,8,8,2,9,0,4,7,3,1,4,0,1,1,4)	(5, 2, 3, 6, 1, 1, 2, 9, 4, 1, 7, 5, 2, 2, 0, 0)
$C_{10,32,5}$	$M_{10,32,2}$	(3,9,9,8,2,6,4,1,5,8,6,3,0,0,3,5)	(0,0,0,1,1,1,8,8,2,5,4,1,3,3,9,2)
$C_{12,32,3}$	$M_{12,32,1}$	(6,7,1,0,8,7,4,3,2,8,0,11,0,1,3,3)	(0,0,0,0,8,4,1,8,2,8,4,7,1,2,10,5)
$C_{12,32,4}$	$M_{12,32,1}$	(10, 3, 9, 4, 3, 11, 6, 11, 11, 2, 0, 0, 0, 1, 1, 2)	(2, 10, 11, 10, 1, 0, 0, 7, 5, 7, 0, 1, 0, 7, 7, 10)
$C_{12,32,5}$	$M_{12,32,1}$	(1, 8, 4, 3, 7, 1, 2, 11, 6, 0, 1, 7, 0, 0, 3, 0)	(0, 11, 3, 5, 7, 9, 6, 9, 11, 3, 11, 5, 0, 1, 4, 5)
$C_{14,32,3}$	$M_{14,32,1}$	(0, 2, 13, 11, 1, 7, 4, 3, 7, 3, 12, 6, 0, 1, 0, 6)	(0, 11, 11, 11, 11, 8, 11, 13, 8, 11, 9, 7, 0, 11, 7, 7)
$C_{14,32,4}$	$M_{14,32,1}$	(11, 12, 1, 1, 3, 9, 11, 7, 13, 5, 12, 13, 0, 5, 8, 5)	(8,3,13,13,9,13,4,1,5,0,13,3,3,3,7,2)
$C_{14,32,5}$	$M_{14,32,2}$	(10, 6, 7, 5, 10, 2, 5, 0, 5, 4, 13, 3, 0, 0, 1, 1)	(1,4,7,7,10,11,5,11,4,0,11,12,0,1,2,10)
$C_{16,32,3}$	$M_{16,32,1}$	(15,0,3,3,12,5,11,0,15,5,14,6,0,0,0,3)	(1, 2, 9, 9, 13, 15, 5, 12, 11, 7, 3, 14, 3, 2, 11, 13)
$C_{16,32,4}$	$M_{16,32,1}$	(3, 11, 1, 0, 12, 3, 10, 10, 10, 11, 6, 12, 1, 1, 3, 0)	(15, 8, 14, 1, 1, 13, 2, 2, 2, 6, 5, 11, 0, 1, 3, 6)
$C_{16,32,5}$	$M_{16,32,2}$	(1, 11, 15, 3, 11, 15, 8, 5, 10, 8, 12, 0, 0, 0, 2, 7)	(2, 14, 3, 10, 2, 11, 11, 0, 11, 13, 1, 1, 0, 1, 0, 12)
$C_{18,32,3}$	$M_{18,32,1}$	(9,4,16,10,2,7,4,7,2,3,2,5,1,0,3,5)	(13, 0, 9, 4, 17, 16, 2, 0, 0, 12, 8, 2, 0, 14, 3, 10)
$C_{18,32,4}$	$M_{18,32,1}$	(6, 12, 5, 7, 5, 6, 6, 7, 16, 15, 5, 9, 1, 0, 2, 4)	(3,6,6,4,13,12,2,13,7,12,4,1,3,1,4,9)
$C_{18,32,5}$	$M_{18,32,2}$	(15, 17, 15, 12, 10, 2, 5, 7, 6, 6, 12, 7, 0, 1, 1, 2)	(12, 16, 1, 17, 13, 16, 10, 14, 7, 9, 4, 9, 1, 10, 0, 13)
$C_{20,32,3}$	$M_{20,32,1}$	(1, 5, 8, 10, 14, 18, 4, 18, 17, 11, 18, 16, 0, 0, 2, 6)	(6, 12, 15, 14, 6, 2, 5, 16, 6, 18, 6, 10, 3, 8, 5, 18)
$C_{20,32,4}$	$M_{20,32,1}$	(6,6,15,6,6,3,1,18,1,19,11,10,0,0,3,5)	(8, 5, 5, 9, 4, 18, 9, 4, 0, 5, 7, 15, 2, 15, 14, 18)
$C_{20,32,5}$	$M_{20,32,1}$	(5, 13, 11, 19, 5, 2, 18, 8, 6, 9, 19, 12, 0, 0, 6, 3)	(19, 14, 13, 18, 2, 14, 3, 2, 10, 12, 12, 8, 19, 16, 12, 2)