

Basic Bicategories

Tom Leinster

Department of Pure Mathematics, University of Cambridge

Email: leinster@dpmms.cam.ac.uk

Web: <http://www.dpmms.cam.ac.uk/~leinster>

1 May, 1998

Minor amendments September 1998

Abstract

A concise guide to very basic bicategory theory, from the definition of a bicategory to the coherence theorem.

Contents

0	Introduction	1
1	Definitions	1
1.0	Bicategories	1
1.1	Morphisms	3
1.2	Transformations	5
1.3	Modifications	6
1.4	Strength Terminology	7
2	Coherence	7
2.0	Functor Bicategories	7
2.1	The Yoneda Embedding	7
2.2	Biequivalence	8
2.3	The Coherence Theorem	8
2.4	Coherence and Commuting Diagrams	8
	References	10

0 Introduction

This is a minimalist account of the coherence theorem for bicategories. The definitions of a bicategory, of a morphism between them, and so on, are given first; from here, a straight-line path is taken to the coherence theorem. No motivation or context is given, and only such examples as are necessary to the development of the theory. More discursive literature on bicategories is available: for instance, the original paper of Bénabou, or Gray's book (see bibliography). In particular, section 9 of Street's paper [7] covers much the same material as this paper.

Nothing here is new (although I haven't seen 2.4 put this way before). The definitions are culled from Bénabou's paper and Gray's book, and the bare bones of the coherence theorem from the papers of Street [6] and of Gordon, Power and Street. Many points are also covered in Lack's thesis.

Acknowledgements

This work was supported by a PhD grant from EPSRC. It was prepared in \LaTeX using Paul Taylor's diagrams package. I am grateful to Jeff Egger for initiating the series of talks which induced me to learn about bicategories.

1 Definitions

1.0 Bicategories

A *bicategory* \mathcal{B} consists of the following data subject to the following axioms:

DATA

- Collection $\text{ob } \mathcal{B}$ (with elements *0-cells* A, B, \dots)
- Categories $\mathcal{B}(A, B)$ (with objects *1-cells* f, g, \dots and arrows *2-cells* α, β, \dots)
- Functors

$$\begin{array}{ccc} c_{ABC} : \mathcal{B}(B, C) \times \mathcal{B}(A, B) & \longrightarrow & \mathcal{B}(A, C) \\ (g, f) & \longmapsto & g \circ f = gf \\ (\beta, \alpha) & \longmapsto & \beta * \alpha \end{array}$$

and $I_A : \mathbf{1} \longrightarrow \mathcal{B}(A, A)$ (thus I_A is a 1-cell $A \longrightarrow A$).

- Natural isomorphisms

$$\begin{array}{ccc}
 \mathcal{B}(C, D) \times \mathcal{B}(B, C) \times \mathcal{B}(A, B) & \xrightarrow{1 \times c_{ABC}} & \mathcal{B}(C, D) \times \mathcal{B}(A, C) \\
 \downarrow c_{BCD} \times 1 & \nearrow a_{ABCD} & \downarrow c_{ACD} \\
 \mathcal{B}(B, D) \times \mathcal{B}(A, B) & \xrightarrow{c_{ABD}} & \mathcal{B}(A, D)
 \end{array}$$

$$\begin{array}{ccc}
 \mathcal{B}(A, B) \times \mathbf{1} & & \mathbf{1} \times \mathcal{B}(A, B) \\
 \downarrow 1 \times I_A & \searrow \sim & \downarrow I_B \times 1 \\
 \mathcal{B}(A, B) \times \mathcal{B}(A, A) & \xrightarrow{c_{AAB}} & \mathcal{B}(A, B) & \mathcal{B}(B, B) \times \mathcal{B}(A, B) & \xrightarrow{c_{ABB}} & \mathcal{B}(A, B) \\
 & \nearrow r_{AB} & & \nearrow l_{AB} & &
 \end{array}$$

thus 2-cells

$$\begin{array}{l}
 a_{hgf} : (hg)f \xrightarrow{\sim} h(gf) \\
 r_f : f \circ I_A \xrightarrow{\sim} f \\
 l_f : I_B \circ f \xrightarrow{\sim} f.
 \end{array}$$

AXIOMS

The following commute:

$$\begin{array}{ccc}
 ((kh)g)f & \xrightarrow{a * 1} & (k(hg))f \\
 \searrow a & & \searrow a \\
 (kh)(gf) & & k((hg)f) \\
 \searrow a & & \swarrow 1 * a \\
 & k(h(gf)) &
 \end{array}$$

$$\begin{array}{ccc}
 (gI)f & \xrightarrow{a} & g(I f) \\
 \searrow r * 1 & & \swarrow 1 * l \\
 & gf &
 \end{array}$$

Variants

If a , l and r are identities, so that $(hg)f = h(gf)$, $If = f = fI$, and similarly for composition of 2-cells, then \mathcal{B} is called a *2-category*. In this case the axioms hold automatically.

Example

There is a 2-category **Cat** whose 0-cells are small categories, whose 1-cells are functors, and whose 2-cells are natural transformations.


Internal Equivalence

As **Cat** is a bicategory, we may imitate certain definitions from category theory in an arbitrary bicategory \mathcal{B} . In particular, an (*internal*) *equivalence* in \mathcal{B} consists of a pair of 1-cells $A \begin{matrix} \xrightarrow{f} \\ \xleftarrow{g} \end{matrix} B$ together with an isomorphism $1 \longrightarrow g \circ f$ in the category $\mathcal{B}(A, A)$ and an isomorphism $f \circ g \longrightarrow 1$ in the category $\mathcal{B}(B, B)$. We also say that f is an equivalence and that A is equivalent to B (inside \mathcal{B}).

The Opposite Bicategory

Given a bicategory \mathcal{B} , we may form a dual bicategory \mathcal{B}^{op} by reversing the 1-

cells but not the 2-cells. Thus if \mathcal{B} has a 2-cell $A \begin{matrix} \xrightarrow{f} \\ \downarrow \alpha \\ \xrightarrow{g} \end{matrix} B$ then \mathcal{B}^{op} has a



1.1 Morphisms

A morphism F (or strictly speaking, (F, ϕ)) from \mathcal{B} to \mathcal{B}' consists of the following data subject to the following axioms:

DATA

- Function $F : \text{ob } \mathcal{B} \longrightarrow \text{ob } \mathcal{B}'$
- Functors $F_{AB} : \mathcal{B}(A, B) \longrightarrow \mathcal{B}'(FA, FB)$

- Natural transformations

$$\begin{array}{ccc}
\mathcal{B}(B, C) \times \mathcal{B}(A, B) & \xrightarrow{c} & \mathcal{B}(A, C) & \mathbf{1} & \xrightarrow{I_A} & \mathcal{B}(A, A) \\
F_{BC} \times F_{AB} \downarrow & & \nearrow \phi_{ABC} & \Downarrow & \nearrow \phi_A & \downarrow F_{AA} \\
\mathcal{B}'(FB, FC) \times \mathcal{B}'(FA, FB) & \xrightarrow{c'} & \mathcal{B}'(FA, FC) & \mathbf{1} & \xrightarrow{I'_{FA}} & \mathcal{B}'(FA, FA)
\end{array}$$

thus 2-cells $\phi_{gf} : Fg \circ Ff \longrightarrow F(g \circ f)$ and $\phi_A : I'_{FA} \longrightarrow FI_A$.

AXIOMS

The following commute:

$$\begin{array}{ccc}
(Fh \circ Fg) \circ Ff & \xrightarrow{\phi * 1} & F(h \circ g) \circ Ff & \xrightarrow{\phi} & F((h \circ g) \circ f) \\
a' \downarrow & & & & \downarrow Fa \\
Fh \circ (Fg \circ Ff) & \xrightarrow{1 * \phi} & Fh \circ F(g \circ f) & \xrightarrow{\phi} & F(h \circ (g \circ f))
\end{array}$$

$$\begin{array}{ccc}
Ff \circ I'_{FA} & \xrightarrow{1 * \phi} & Ff \circ FI_A & \xrightarrow{\phi} & F(f \circ I_A) & I'_{FB} \circ Ff & \xrightarrow{\phi * 1} & FI_B \circ Ff & \xrightarrow{\phi} & F(I_B \circ f) \\
r' \downarrow & & \downarrow Fr & & \downarrow Fl & l' \downarrow & & \downarrow Fl \\
Ff & \xlongequal{\quad\quad\quad} & Ff & & Ff & \xlongequal{\quad\quad\quad} & Ff
\end{array}$$

Variants

If ϕ_{ABC} and ϕ_A are all natural isomorphisms, so that $Fg \circ Ff \cong F(g \circ f)$ and $FI \cong I'$, then F is called a *homomorphism*. If ϕ_{ABC} and ϕ_A are all identities, so that $Fg \circ Ff = F(g \circ f)$ and $FI = I'$, then F is called a *strict homomorphism*.

Representables

If A is a 0-cell of a bicategory \mathcal{B} , there arises a homomorphism $\mathcal{B}(-, A) : \mathcal{B}^{\text{op}} \longrightarrow \mathbf{Cat}$. The 2-cells “ ϕ_{gf} ” and “ ϕ_B ” come from a and r , respectively.

Local Properties

Let P be a property of functors. We say a morphism F is *locally P* if each functor F_{AB} has the property P : thus *locally faithful*, *locally an equivalence*, ...

1.2 Transformations

A transformation $F \begin{matrix} \mathcal{B} \\ \xrightarrow{\sigma} \\ \mathcal{B}' \end{matrix} G$, where $F = (F, \phi)$ and $G = (G, \psi)$ are mor-

phisms, is defined by the following data and axioms. Below, we use the notation $h_* : \mathcal{B}(C, D) \longrightarrow \mathcal{B}(C, E)$ for the functor induced by a 1-cell $D \xrightarrow{h} E$ of a bicategory \mathcal{B} , and similarly $h^* : \mathcal{B}(E, C) \longrightarrow \mathcal{B}(D, C)$.

DATA

- 1-cells $FA \xrightarrow{\sigma_A} GA$
- Natural transformations

$$\begin{array}{ccc} \mathcal{B}(A, B) & \xrightarrow{F_{AB}} & \mathcal{B}'(FA, FB) \\ \downarrow G_{AB} & \nearrow \sigma_{AB} & \downarrow (\sigma_B)_* \\ \mathcal{B}'(GA, GB) & \xrightarrow{(\sigma_A)^*} & \mathcal{B}'(FA, GB) \end{array}$$

thus 2-cells $\sigma_f : Gf \circ \sigma_A \longrightarrow \sigma_B \circ Ff$.

AXIOMS

The following commute:

$$\begin{array}{ccc} (Gg \circ Gf) \circ \sigma_A \xrightarrow{a'} Gg \circ (Gf \circ \sigma_A) \xrightarrow{1 * \sigma_f} Gg \circ (\sigma_B \circ Ff) \xrightarrow{a'^{-1}} (Gg \circ \sigma_B) \circ Ff \xrightarrow{\sigma_g * 1} (\sigma_C \circ Fg) \circ Ff \xrightarrow{a'} \sigma_C \circ (Fg \circ Ff) \\ \downarrow \psi * 1 \qquad \qquad \qquad \downarrow 1 * \phi \\ G(g \circ f) \circ \sigma_A \xrightarrow{\sigma_{gf}} \sigma_C \circ F(g \circ f) \\ \downarrow \psi * 1 \qquad \qquad \qquad \downarrow 1 * \phi \\ I'_{GA} \circ \sigma_A \xrightarrow{l'} \sigma_A \xrightarrow{r'^{-1}} \sigma_A \circ I'_{FA} \\ \downarrow \psi * 1 \qquad \qquad \qquad \downarrow 1 * \phi \\ GI_A \circ \sigma_A \xrightarrow{\sigma_{I_A}} \sigma_A \circ FI_A \end{array}$$

Variants


If σ_{AB} are all natural isomorphisms then σ is called a *strong transformation*. If σ_{AB} are all identities then σ is called a *strict transformation*.

Representables

If $A \xrightarrow{f} B$ is a 1-cell in a bicategory, then there arises a strong transformation $f_* = \mathcal{B}(-, f) : \mathcal{B}(-, A) \longrightarrow \mathcal{B}(-, B)$. The 2-cells “ σ_g ” come from the associativity isomorphism a .


1.3 Modifications

A modification



consists of the following data subject to the following axioms:

DATA



AXIOMS

The following commute:


$$\begin{array}{ccc} Gf \circ \sigma_A & \xrightarrow{1 * \Gamma_A} & Gf \circ \tilde{\sigma}_A \\ \sigma_f \downarrow & & \downarrow \tilde{\sigma}_f \\ \sigma_B \circ Ff & \xrightarrow{\Gamma_B * 1} & \tilde{\sigma}_B \circ Ff \end{array}$$

Variants

None.

Representables

If $A \begin{array}{c} \xrightarrow{f} \\ \downarrow \alpha \\ \xrightarrow{g} \end{array} B$ is a 2-cell in a bicategory, then there arises a modification



1.4 Strength Terminology

The terminology used to describe whether something holds strictly, up to isomorphism, or just up to a connecting map, has evolved messily. Here is a summary of the definitions given above; note that for representables, everything is at the ‘iso’ level.

	bicategories	morphisms	transformations
map	—	(plain)	(plain)
iso	bicategory	homomorphism	strong transformation
equality	2-category	strict homomorphism	strict transformation

2 Coherence

2.0 Functor Bicategories

Given a pair of bicategories \mathcal{B} and \mathcal{B}' , one can define (canonically) a ‘functor bicategory’ $\mathbf{Lax}(\mathcal{B}, \mathcal{B}')$, whose 0-cells are morphisms $\mathcal{B} \rightarrow \mathcal{B}'$, whose 1-cells are transformations, and whose 2-cells are modifications. This is not in general a 2-category, but it is if \mathcal{B}' is. We will take a particular interest in the sub-bicategory $[\mathcal{B}, \mathcal{B}']$ of $\mathbf{Lax}(\mathcal{B}, \mathcal{B}')$, consisting of homomorphisms, strong transformations and modifications (i.e. everything at the ‘iso’ level).

2.1 The Yoneda Embedding

By the observations of the previous section, there is for each bicategory \mathcal{B} a 2-category $[\mathcal{B}^{\text{op}}, \mathbf{Cat}]$. Moreover, the representables constructed in Section 1 provide a ‘Yoneda’ homomorphism $Y : \mathcal{B} \rightarrow [\mathcal{B}^{\text{op}}, \mathbf{Cat}]$. It is straightforward to calculate that locally Y is full, faithful, and essentially surjective on objects— in other words, that Y is a local equivalence. (In order for Y to exist it is

necessary that \mathcal{B} should be locally small, but we do not emphasize this issue here.)

2.2 Biequivalence

Let \mathcal{B} and \mathcal{B}' be bicategories. A *biequivalence* from \mathcal{B} to \mathcal{B}' consists of a pair of homomorphisms $\mathcal{B} \begin{matrix} \xrightarrow{F} \\ \xleftarrow{G} \end{matrix} \mathcal{B}'$ together with an equivalence $1 \longrightarrow G \circ F$ inside the bicategory $[\mathcal{B}, \mathcal{B}]$ and an equivalence $F \circ G \longrightarrow 1$ inside $[\mathcal{B}', \mathcal{B}']$. We also say that F is a biequivalence and that \mathcal{B} is biequivalent to \mathcal{B}' . Now, just as for equivalence of plain categories, there is an alternative criterion for biequivalence: namely, that a homomorphism $F : \mathcal{B} \longrightarrow \mathcal{B}'$ is a biequivalence if and only if F is locally an equivalence and is surjective-up-to-equivalence on objects. The latter condition means that if B' is any 0-cell of \mathcal{B}' then there is some 0-cell B of \mathcal{B} such that FB is (internally) equivalent to B' .

2.3 The Coherence Theorem

Theorem *Every bicategory is biequivalent to a 2-category.*

Proof Let \mathcal{B} be a bicategory and $Y : \mathcal{B} \longrightarrow [\mathcal{B}^{\text{op}}, \mathbf{Cat}]$ the Yoneda map. Let \mathcal{B}' be the full image of Y : that is, the sub-2-category of $[\mathcal{B}^{\text{op}}, \mathbf{Cat}]$ whose 0-cells are those in the image of Y , and with all 1- and 2-cells of $[\mathcal{B}^{\text{op}}, \mathbf{Cat}]$ between them. Let $Y' : \mathcal{B} \longrightarrow \mathcal{B}'$ be the restriction of Y . Then:

- Y' is a homomorphism, since Y is
- Y' is surjective on 0-cells, by construction
- Y' is a local equivalence, since Y is (observed in 2.1).

Thus Y' is a biequivalence from \mathcal{B} to the 2-category \mathcal{B}' . □

2.4 Coherence and Commuting Diagrams

Coherence theorems sometimes have the form ‘all diagrams of a certain kind commute’; for instance, the coherence theorem for monoidal categories states that all diagrams built out of the associativity and identity isomorphisms commute. This result for monoidal categories also holds for bicategories. Without giving a precise statement or proof, we indicate by an example how it is a corollary of our coherence theorem, 2.3.

A typical instance of what we wish to prove is that the diagram

$$\begin{array}{ccc}
 & (h(Ig))f & \\
 a^{-1} * 1 \swarrow & & \searrow a \\
 ((hI)g)f & & h((Ig)f) \\
 (r * 1) * 1 \downarrow & & \\
 (hg)f & & \\
 a \swarrow & & \searrow 1 * (l * 1) \\
 & h(gf) &
 \end{array} \tag{1}$$

should commute, for any composable 1-cells f, g, h in any bicategory. Let us say that a bicategory \mathcal{B} has the *coherence property* if all diagrams ‘like this’ in \mathcal{B} commute; our goal is to show that every bicategory has the coherence property. To achieve this, first observe that every 2-category has the coherence property, since a, l and r are all 1. Then, for any \mathcal{B} we have an ‘embedding’ $Y : \mathcal{B} \longrightarrow [\mathcal{B}^{\text{op}}, \mathbf{Cat}]$ of \mathcal{B} into a bicategory with the coherence property, and this implies that \mathcal{B} too has the coherence property, as now explained.

Let $(F, \phi) : \mathcal{B} \longrightarrow \mathcal{B}'$ be a morphism of bicategories, and suppose that \mathcal{B}' has the coherence property. We want to deduce that, subject to certain conditions on (F, ϕ) , the bicategory \mathcal{B} also has the coherence property. Consider the diagram (1) in \mathcal{B} . Let α be the composite down the left-hand side, and β down the right; let α' be the composite

$$(Fh \circ (I' \circ Fg)) \circ Ff \xrightarrow{a'^{-1} * 1} ((Fh \circ I') \circ Fg) \circ Ff \xrightarrow{(r' * 1) * 1} (Fh \circ Fg) \circ Ff \xrightarrow{a'} Fh \circ (Fg \circ Ff),$$

and similarly β' . We know that $\alpha' = \beta'$ and want to conclude that $\alpha = \beta$. Consider, then, the diagram in Figure 2, where the 2-cells called ϕ are built up from ϕ_A 's and ϕ_{qp} 's. By definition of morphism, the diagram commutes—that is, $F\alpha \circ \phi = \phi \circ \alpha'$. Similarly, $F\beta \circ \phi = \phi \circ \beta'$, so $F\alpha \circ \phi = F\beta \circ \phi$. If F is a homomorphism then the 2-cells ϕ are isomorphisms, so $F\alpha = F\beta$; if also F is locally faithful then we may conclude that $\alpha = \beta$.

Our example therefore demonstrates: if $F : \mathcal{B} \longrightarrow \mathcal{B}'$ is a locally faithful homomorphism and \mathcal{B}' has the coherence property, then so does \mathcal{B} . Applying this to $Y : \mathcal{B} \longrightarrow [\mathcal{B}^{\text{op}}, \mathbf{Cat}]$, for any bicategory \mathcal{B} , shows that all bicategories have the coherence property.

$$\begin{array}{ccc}
(Fh \circ (I' \circ Fg)) \circ Ff & \xrightarrow{\phi} & F[(h(Ig))f] \\
\downarrow a'^{-1} * 1 & & \downarrow F[a^{-1} * 1] \\
((Fh \circ I') \circ Fg) \circ Ff & \xrightarrow{\phi} & F[((hI)g)f] \\
\downarrow (r' * 1) * 1 & & \downarrow F[(r * 1) * 1] \\
(Fh \circ Fg) \circ Ff & \xrightarrow{\phi} & F[(hg)f] \\
\downarrow a' & & \downarrow F[a] \\
Fh \circ (Fg \circ Ff) & \xrightarrow{\phi} & F[h(gf)]
\end{array}$$

Figure 2: diagram of 2-cells in \mathcal{B}'

References

- [1] Jean Bénabou, Introduction to bicategories (1967). In: *Reports of the Midwest Category Seminar*, ed. Bénabou et al, Springer LNM 47.
- [2] R. Gordon, A. J. Power, Ross Street, Coherence for tricategories (1995). *Memoirs of the AMS*, vol. 117, no. 558.
- [3] J. Gray, *Formal Category Theory: Adjointness for 2-Categories* (1974). Springer LNM 391.
- [4] G. M. Kelly, R. Street, Review of the elements of 2-categories (1974). In: *Category Seminar*, Springer LNM 420, pp 75–103.
- [5] Stephen Lack, *The Algebra of Distributive and Extensive Categories* (1995). PhD thesis, University of Cambridge. Available via <http://cat.maths.usyd.edu.au/~stevl>.
- [6] Ross Street, Fibrations in bicategories (1980). *Cahiers Top. Geom. Diff.*, vol. XXI, no. 2.
- [7] Ross Street, Categorical structures (1995). In: *Handbook of Algebra*, Vol. 1, ed. M. Hazewinkel, Elsevier North-Holland.