

Reference	<p>On, the patient underwent right frontal craniotomy for resection of tubercular sellar meningioma. There were no intraoperative complications. Postoperatively the patient was monitored in the Surgical Intensive Care Unit. She was awake, alert, and oriented times three. Pupils were 5 down to 3 mm bilaterally. The face was symmetric. She had no drift. Her strength was in all muscle groups. She continued to have a left superior field cut. Her dressing was clean, dry, and intact. She was transferred to the regular floor. Postoperatively she was very sleepy and difficult to arouse. It took several days for the patient's mental status to clear and she was finally awake, alert, and oriented times three and up out of bed on postoperative day #3 with full extraocular movements, no drifts, and face symmetric. She was seen by physical therapy and occupational therapy and found to be safe for discharge to home.</p>
Original	<p>Pt was admitted to the neurosurgery service and monitored closely in the ICU. She remained neurologically stable and was transferred to the floor on. She was seen by physical therapy and occupational therapy and found to be safe for discharge to home. She was discharged to home on in stable condition with follow up arranged with Dr. in 4 weeks with a noncontrast head CT. She will also follow up in the Brain Clinic in 2 weeks with an MRI MRA of the brain to assess the extent of the lesion and to determine if it is communicating with the vascular system. She is to call Dr. office for a follow up appointment.</p>
Filtered	<p>41 yo female with pmh of HTN, NIDDM, migraines, and anxiety disorder admitted with worsening HAs, blurred vision, L eye swelling. s p craniotomy for resection of parasellar meningioma on. Neuro: lethargic, yet easily with verbal stimuli. PERRLA, speech clear. Oriented x3. MAE, eqaul strength. R frontal incision with dsd, old serosang drainage noted, no new drainage. flat affect, pt appears depressed. Dr. aware, saw pt, assessed lethargy. CV: HR 50s sinus brady. SBP 120-135.</p>
Extractive Revision	<p>Transfer note 41 yo female with pmh of HTN, NIDDM, migraines, and anxiety disorder admitted with worsening HAs, blurred vision, L eye swelling. S p craniotomy for resection of parasellar meningioma on. condition update d: PT TRANSFERRED FROM PACU AT 1900- LETHARGIC BUT, IN PLACE, NO GTTS HEMODYNAMICS- HR 42-54 SB WITHOUT ECTOPY, SBP 110-134 NPO EXCEPT FOR MEDS, SM AMT BILIOUS EMESIS X1- MEDICATED WITH ZOFRAN WITH SOME RELIEF</p>
Abstractive Revision	<p>This is a 41 year old female with a history of HTN, NIDDM, migraines, and anxiety disorder who presented with worsening HAs, blurred vision, L eye swelling, and was found to have a suprasellar meningioma. She was admitted to the neurosurgery service and underwent a craniotomy for resection of the lesion on. She was transferred to the PACU in stable condition. She had no further episodes of bradycardia. Her neuro status was stable post-operatively. Her speech was clear and she was able to tolerate po intake. Her O2 sats were 95% on room air.</p>