

## Remembering Spillover Erosion of Grand Canyon

**Steven A. Austin**, Austin Research Consulting Inc., Sewickley, Pennsylvania, 15143

**Edmond W. Holroyd III**, Carmel Consulting, Arvada, Colorado, 80002

**David R. McQueen**, CSB Geology, Mangham, Louisiana, 71259


### Abstract

Grand Canyon of Arizona is the best-known example of hydraulic erosion on planet earth. No other canyon has been so carefully scrutinized by geologists. Therefore, Grand Canyon is the most important landscape on planet earth to be explained by competing creationist and evolutionist paradigms. Four hypotheses have been proposed for the erosion of Grand Canyon: (1) drainage spillover, (2) antecedent river, (3) stream piracy, and (4) flood drainage. The first geologist ever to explore Grand Canyon, John Newberry, recognized in 1858 that it was carved through a highland arch that separated topographic basins. Newberry proposed drainage spillover of the elevated terrain by what has later been called ancient "Hopi Lake" (aka "Lake Bidahochi"). We trace the history of "spillover" and "breached dam" hypotheses from Newberry's first discovery, but we focus on the last 50 years through both creationist and evolutionist thinking. Fifty years ago, when uniformitarian doctrine was stifling imaginative thinking about Grand Canyon erosion, Hopi Lake was "the lake that gets no respect," and spillover was "the forgotten transverse drainage hypothesis." Also, 50 years ago, three creationists (Henry M. Morris, Jr., Clifford Burdick, and Bernard Northrup) affirmed the creationist consensus that Grand Canyon was eroded by catastrophic spillover of a post-Flood lake. That recent erosion formed a young Colorado River. In 1988 Steve Austin and Ed Holroyd were working on the configuration of lakes east and north of Kaibab Upwarp. Bob Scarborough had surveyed lake sediments to understand better the western boundary of ancient Hopi Lake. Also, in 1988, Norm Meek had "rediscovered" Afton Canyon of the Mojave Desert in California that had been carved by spillover of ancient Lake Manix producing a one-tenth scale analog to Grand Canyon. By 1988, pieces of the larger spillover puzzle were being assembled. By the year 2000, Bob Scarborough, Norm Meek, Ron Dom, Jon Spencer, Philip Pearthree, John Douglass, Kyle House and Todd Dallegge were developing these ideas within the evolutionist community. Also, by 2000, Andrew Snelling and Tom Vail were communicating these ideas within the creationist community. By 2012, the spillover explanation was called "a favored concept for two decades." Responding to spillover's popularity, a self-appointed panel of experts challenged the notion. "Afton Canyon Controversy" focused on a different model for Lake Manix with headward erosion of Afton Canyon. Fact rechecking by U.S. Geological Survey silenced critics in the "Afton Canyon Controversy" by 2014. The "Crooked Ridge Miocene River" called for a much different basin configuration without Hopi Lake against the Kaibab Upwarp. Spillover critics that promoted the "Crooked Ridge Miocene River" experienced an almost simultaneous "spirit of repentance" killing quickly the "Crooked Ridge River" in 2016. Since then, Lake Manix and Hopi Lake have been restored, silently, as viable spillover candidates. Why no fanfare? These lakes continue to receive distressing abuse from "the establishment," verifying they are the lakes that get no respect. This spillover story shows how pervasive and deep-seated evolutionary assumptions are within the "Grand Debate." Today, more than 20 earth scientists (most with Ph.D. degrees) have constructively affirmed spillover theory, at the same time, maintaining an attitude of respect, not contempt, toward the ancient lake in northeastern Arizona. Geologists have been searching for the "Miocene River" for 150 years. If it is found, it would be the primary alternate to the spillover hypothesis. We should remember an important fact—creationist and evolutionist thinking about spillover continues to make a significant contribution to our understanding of erosion of Grand Canyon.

**Keywords:** Colorado River, Grand Canyon, erosion, Kaibab Upwarp, East Kaibab monocline, transverse drainage, spillover model, overtopping, ponding and overflow, breached dam hypothesis, Hopi Lake, Lake Bidahochi, Crooked Ridge, Bidahochi Formation, Mount St. Helens, tufa, Lake Manix, Afton Canyon, Pliocene lake sedimentation, Miocene river sediments, tuff ring, maar, scoria cone, paleontology, tectonics, isostasy, framework of Scripture, geologic time

The morning of Sunday, April 10, 1988 began like most early spring days at the iconic landscape in extreme eastern Grand Canyon National Park. The location is at the Canyon's rim in a place so obscure that National Park rangers refer to it simply as "No Name Point." Although one never finds No Name Point located on maps, it does have a guard rail at the Grand Canyon rim. As the sun

rose that morning and its rays grazed the extreme southeastern rim, the colors of the eastern Canyon wall began to gleam. The browns and grays of early morning soon became reddish, yellowish, and greenish hues as cliffs above the Colorado River reflected the increasing intensity of the sun. The normal chatter of ravens and squirrels began among the pine and juniper forest at the Canyon rim, but



**Fig. 1.** Grand Canyon is positioned across the elevated Kaibab Upwarp in northern Arizona. No Name Point is ideally located at 7,100ft elevation for discussion of erosion of Grand Canyon. Colorado River within the Canyon is at elevation of 2,545ft. Oblique aerial view is toward the northwest along the axis of the East Kaibab Monocline. Upper Colorado River enters from upper right and flows into Grand Canyon at Kaibab Upwarp (higher elevation forested area in center) and exits photo on the upper left. No Name Point is 13mi east of Grand Canyon Village. Width of this view is 100mi. International Space Station photo ISS039-E-5258 acquired on March 25, 2014.

this Sunday morning was going to be something new and unusual. No Name Point was going to inspire a memorable discussion on the erosion of Grand Canyon (fig. 1).

### Church on the Canyon's Rim?

Just after 9a.m. on Sunday, April 10, 1988, a few cars appeared at the obscure gravel and pavement turnout adjacent to the Canyon rim. People stepped out of cars and began walking by trail northward through the forest to a railing at the Canyon rim. No Name Point began to bustle with activity. Then, a most unusual event occurred. Two large buses and more cars dispatched passengers as a hundred people assembled at the rail fence. Then, a third bus arrived! It may have been the largest assembly of Christians yet on the Canyon's rim (fig. 2). People standing there began to marvel at the spectacle visible from that Canyon rim overlook at elevation of 7,100ft. As the crowd squeezed together behind the railing at the Canyon rim, some sat on folding chairs, but most sat on the flat limestone surface. Then, Tom Manning, the event host, with a loud voice announced the remote assembly was to open

in a prayer of thanksgiving. Almost spontaneously after the prayer, the crowd erupted in song with the words, "When through the woods and forest glades I wander...When I look down from lofty mountain grandeur..." These words are the second verse from



**Fig. 2.** People begin assembling at the lecture location called No Name Point on the rim of Grand Canyon. Here the third bus has just arrived. People interested in Grand Canyon erosion are getting seated. About 140 attended that lecture on Sunday morning April 10, 1988. Photo by John D. Morris.

the well-known Christian hymn titled “How Great Thou Art.” It was apparent to all that this Sunday morning at No Name Point was to be a worship and teaching event at the rim of Grand Canyon! The park ranger couldn’t remember any assembly like that at No Name Point.<sup>1</sup>


After some formalities at the Canyon rim, a Scripture from the Book of Job was read (“He cuts out rivers amongst the rocks...”) and a geologist, Dr. Steve Austin stood up and delivered a morning “message” titled “Erosion of Grand Canyon.” That teaching about Canyon erosion was unusual because it deviated from the expected explanation that the Park rangers give that the Colorado River eroded Grand Canyon very slowly over tens of millions of years. Instead, Dr. Austin reported evidence that Grand Canyon was eroded in just a few weeks just thousands of years ago by catastrophic drainage of lakes. The regional drainage of the entire Colorado

River is conveniently divided into upper and lower basins (fig. 3). Austin explained that a computer plot of elevation data shows that an enormous lake or series of lakes bigger than one of the Great Lakes could be contained within the present topographic region extending from the lower to the upper Colorado River basins. The series of lakes could form east and north of Grand Canyon if the eastern Grand Canyon was plugged by an enormous dam. The location of ancient lakes affected the entire Colorado River drainage basin (fig. 4). Austin pointed to lime sediment layers east of Grand Canyon visible at distance from No Name Point as evidence that at least one ancient lake once sat above 6,000ft elevation over Cape Solitude east of Grand Canyon (fig. 4). That lake was named Hopi Lake. Then, he described the topographic similarity of Grand Canyon to the spillway of a breached landscape at Mount St. Helens that was eroded in a single day on March 19, 1982.



**Fig. 3.** Colorado River drainage basin extends into parts of seven states and Mexico. The drainage basin is divided into “lower basin” and “upper basin,” with the boundary being drawn by the Bureau of Reclamation at Lees Ferry on the river northeast of Grand Canyon. Blue lines show major rivers, and black lines depict canals and pipes which the Bureau of Reclamation uses to transport water for nearby urban areas. Copyright International Mapping Associates, used by permission.

<sup>1</sup> National Park rangers tell us that Grand Canyon rim overlooks are for small (linear) assemblies of people against a railing. Contrast smaller rim assemblies against a railing with much larger assemblies around a location (e.g. Yellowstone’s Old Faithful Geyser).



**Fig. 4.** Eastern Grand Canyon location map shows the places and features used in discussions of the breached dam (spillover) hypothesis for erosion of Grand Canyon. Map includes Austin's and Holroyd's proposed locations of three ancient lakes and their seven spillover points. Shaded relief base map is from US Geological Survey 10m DEM processed using ESRI software.

That teaching about rapid erosion from spillover was well received by the crowd and a lively discussion about canyon erosion followed as the people dispersed that Sunday morning over 30 years ago. That teaching event on the rim of eastern Grand Canyon in April 1988 was an early statement of what has been called the breached dam hypothesis for the erosion of Grand Canyon.<sup>2</sup>

### Are Spilling Lakes All It Takes?

How geologists started thinking about erosion of Grand Canyon is also an interesting story worth remembering. John Newberry in April 1858 was the first geologist to explore, interpret and report upon erosion in Grand Canyon (Newberry 1861; Newberry 1862).<sup>3</sup> He recognized that strata were continuous through basins and arches, but strata continuity was broken by water erosion. But, what style of water erosion? Newberry wrote: "Doubtless in earlier times it [Colorado River]

filled these basins to the brim....its accumulated waters, pouring over the lowest points in the barriers which opposed their progress towards the sea, have cut them down from summit to base, forming that remarkable series of the deep and narrow canyons through which its turbid waters now flow...." (Newberry, 1861, 19, 20). Newberry observed lake clay deposits (now called Bidahochi Formation) along the Little Colorado River, and he proposed lake overflow of the topographic surface of Arizona was the cause of Grand Canyon's unique style of erosion. Notice that Newberry's explanation was that the geologic structure (strata of uplifted plateaus, faults and folds) formed *before* the Colorado River eroded the surface. The river was the latest addition to the landscape, therefore, a "young river." It was an extraordinary achievement. The first geologist ever to explore Grand Canyon, John Newberry, recognized in 1858 that it was carved by drainage spillover of what has later been called ancient "Hopi Lake."

<sup>2</sup> Austin delivered a similar breached-dam lecture at the South Rim on April 12, 1987. Participants at the April 10, 1988 lecture received study material (Austin, 1988). That 1988 Guidebook contains a five-page description of catastrophic drainage explanations for erosion of Grand Canyon and has attached an extensive annotated bibliography. It was a republication document written for field trip participants, and was not for sale or wider distribution.

<sup>3</sup> Newberry's investigation of western Grand Canyon at Diamond Creek caused him to appreciate the elevated land as a topographic barrier. Although he never visited Kaibab Plateau, he understood, from his field work in the Little Colorado River area, that it was a lower region where sediment showed natural ponding. Thus, Newberry proposed that the lake breached to erode Grand Canyon.

Observations in 1869 and 1871 on two river expeditions encouraged John Wesley Powell to think differently. Powell believed the Colorado River was older than those geologic uplifts and topographic barriers (Powell 1875). The issue was not what he could see, but what he could imagine. He supposed those uplifts were raised across the ancestral river's path; therefore, he imagined the "old river" was *antecedent* to the geologic structure. Two other geologists Charles Walcott (Walcott 1890) and William Morris Davis (Davis 1901) added to Powell by suggesting that the "old river" exhumed older strata that once overlay the rim of Grand Canyon. They suggested the "old river" was *superimposed* as it was let down through strata and structure that have been removed, thus, explaining why there is a lack of evidence for the hypothesis! Again, prominent in geologists' minds was what was *not* seen, and their presuppositional agenda as they interpret the river.

Seventy years after Newberry's spillover explanation, geologists in the early 1930s were carefully studying the lower Colorado River corridor for a site to build Hoover Dam and Lake Mead. One of these was Eliot Blackwelder, chairman of the Geology Department at Stanford University (Blackwelder 1934). Another was Chester Longwell, professor of geology at Yale University (Longwell 1928). Both Blackwelder and Longwell were proficient at distinguishing river gravels and river sand from lake-deposited clay and limestone. Focus of study was the area directly west of Grand Canyon, the region around today's Lake Mead (fig. 3). That is the area described by Newberry. Blackwelder and Longwell identified Pleistocene river sediment sitting directly on top of thick Pliocene green clay, limestone and gypsum salts (what was called Bouse Formation). Blackwelder recognized a straightforward explanation, and he knew that Newberry had already understood the issue *70 years* earlier.<sup>4</sup> Like Newberry, Blackwelder suggested the Southwest was once a series of closed drainage basins with big lakes.<sup>5</sup> He imagined, "a chain of lakes strung upon a river" (Blackwelder 1934, 562). He supposed highlands with basins between that filled forming lakes that eventually overtopped barriers spilling as rivers and eroding bedrock canyons into the adjacent basins. He visualized a "young river" where a lake overflowed Kaibab Upwarp.

One of Blackwelder's perceptive insights was that a spillover explanation ought to apply to

landscapes outside the Colorado River drainage. Elmer Ellsworth, Blackwelder's graduate student, described the noteworthy example in 1932 at Afton Canyon on the Mojave River in Southern California. Like Grand Canyon, Afton Canyon was understood to have been eroded across a mountain by catastrophic drainage of a big lake (Blackwelder and Ellsworth 1936). By 1936 the mudstone and sandstone strata of the Bidahochi Formation were formally recognized by geologist Howel Williams to be a deposit from an ancient lake in eastern Arizona. Williams (1936) proposed ancient "Hopi Lake" (aka "Lake Bidahochi"), a 12,000-square-mile-area lake that occupied the basin on the east side of the Kaibab and Coconino plateaus at elevation above 6,000ft. Scientists at that time began to use the technical term "transverse drainage" to describe rivers with canyons that cut across mountains. Things looked promising for the spillover explanation of Grand Canyon.

However, the second half of the twentieth century was the time that geologists embraced uniformitarian doctrine to such an extent that theories of landscape evolution were championed. That thinking expressed itself in the former century's "old river" Grand Canyon erosion narrative that still pervades our culture. According to the narrative, rivers evolve landscapes through tens of millions of years of erosion from lowlands to highlands. That process is called "headward erosion," or, more correctly "drainage-head erosion" (Hilgendorf et al. 2020). Canyons are part of landscapes that evolve slowly from the bottom up ("headward"). The process of "stream capture" is supposed to involve a precocious gully that eroded headward from the west to the east across Kaibab Upwarp diverting the upper drainage basin toward the Pacific Ocean. Those earliest explanations of Newberry, Blackwelder, and Williams understood erosion to occur over high-country barriers, then extending into lowlands. Spillover was erosion from the top down, *backwards* from the uniformitarian doctrine. As the last half of the twentieth century unfolded, spillover was lost from memory. According to Dr. Norman Meek the twentieth century was the time when "ponding and overflow became the forgotten transverse drainage hypothesis" (Meek 2002).

Fifty years ago, three creationists explored Grand Canyon erosion ideas, affirming what had become

<sup>4</sup> Blackwelder (1934) clearly cites and acknowledges Newberry's earlier contribution to spillover erosion of Grand Canyon. Blackwelder was first to propose that the entire Colorado River drainage basin was eroded by multiple spillovers.

<sup>5</sup> A misstatement of historical fact occurs in Oard 2016, 39: "Geologist Eliot Blackwelder was the first to propose that Grand Canyon was eroded by rushing water derived from the spillover of a lake that was ponded northeast of the Kaibab Plateau." Mike Oard appears to be oblivious to the earlier work of John Strong Newberry.

a long-established creationist consensus.<sup>6</sup> Dr. Clifford Burdick, geologist from Tucson, supposed late Flood uplift and ponding of water behind the Kaibab Upwarp with the unnamed lake on the east side breaching in the post-Flood period to erode the Canyon.<sup>7</sup> Dr. Henry M. Morris, Jr., who affirmed Byron Nelson's (1931) timing of post-Flood erosion, spoke favorably of Burdick's detailed explanation of rapid post-Flood drainage of a big lake (Whitcomb and Morris 1961).<sup>8</sup> Dr. Bernard Northrup, seminary professor in Old Testament from San Francisco, favored post-Flood elevation of the Kaibab Upwarp, followed by basin filling forming enormous "Lake Kaiparowitz" that was quickly breached forming Ice Age meltwater floods.<sup>9</sup> Dr. Austin recalls, "In 1968 I spoke privately about Grand Canyon with Henry Morris when he was speaking at University of Washington in Seattle. Also, in 1968, Ed Nafziger, a Seattle science teacher and veteran Grand Canyon hiker, introduced me to Cliff Burdick and Bernie Northrup. Burdick, Northrup and I did field work together in summer of 1968. These men not only transferred to me a passion for exploring Grand Canyon, but an understanding of spillover as a powerful erosive agent that could carve solid rock layers in the years after Noah's Flood." I am humbled to recall these discussions about Grand Canyon

erosion. I reflect on the possibility that, at that time, we may have been the only people on planet earth that were talking about spillover."

### A Memorable Lunch Discussion

Eighteen years after Austin's field work with Burdick and Northrup, three scientists remembered those erosion ideas. It was a lunch meeting in the cafeteria at Duquesne University in Pittsburgh in August 1986. The three scientists were Dr. Steve Austin, professor of geology at Institute for Creation Research Graduate School, Mr. David McQueen, also professor of geology at Institute for Creation Research, and Dr. Edmond Holroyd, physicist with the US Bureau of Reclamation in Denver. The Pittsburgh lunch discussion in 1986 among the three scientists concerned the erosion of the Colorado Plateau including Grand Canyon. Figs. 3 and 4 are maps showing the locations and features that were points for discussion.

McQueen recalls that Pittsburgh meeting, "Dr. Austin talked about his observations that the silt deposits near Hopi Buttes, Arizona indicate that a very large and high elevation ancient lake existed behind the Kaibab Upwarp in northeastern Arizona that others had called Hopi Lake. He suggested also that sediment deposits on the extreme eastern rim of Grand Canyon could be evidence of that big lake

<sup>6</sup> The 1968 creationist consensus concerning Grand Canyon erosion appears to go back to the 1930s to the disciples of George McCready Price. Disciples of Price sought to understand post-Flood glaciation and how it related to sculpting of the landscape, with special reference to western North America. That directed their thinking to Grand Canyon. Byron C. Nelson, a reader of Price, wrote of Grand Canyon "...dug in the first centuries following the Flood" (Nelson 1931, 64). Henry M. Morris, Jr. was very familiar with Nelson's book, having read it in 1943 (see "Foreword" written by Morris for the 1968 reprint of *Deluge Story in Stone* by publisher Bethany Fellowship). Also, Harold W. Clark, who succeeded Price as geology professor at Pacific Union College, wrote, "The Grand Canyon of the Colorado gives every evidence of having been formed by a gigantic crack in the earth's surface" (Clark 1946, 91). Clark, who taught geology for 37 years, applied post-Flood spillover thinking to the upper Colorado River drainage basin. He wrote: "The Green River Basin in southwest Wyoming filled with sediments until an opening appeared in the Uinta uplift to the south. Whether a natural channel through a low spot allowed water to drain from the basin, or whether an earthquake crack started the present gorge, we will never know. Anyway, the gorges of the Green River witness to tremendous erosion." (Clark 1968, 147). Also, Clark commented on the Arkansas River, "In the South Park, in Colorado, sediments accumulated until the overflow was let out through Royal Gorge, which eventually cut down to a depth of 1,200 feet." (Clark 1968, 150). Clifford Burdick later summarized that creationist consensus explanation of Grand Canyon as rapid, post-Flood drainage of ponded water bodies through the cracked dome of the Kaibab Anticline.

<sup>7</sup> Austin and Burdick had conversations on Grand Canyon erosion in the summer of 1968 while doing field work in Glacier National Park. Burdick later published on Grand Canyon erosion (Burdick 1974, 27): "...when this Kaibab Anticline rose, it may have dammed up the water toward Colorado, Utah, and Wyoming, to form a giant inland sea somewhat similar to Lake Bonneville in Utah." Burdick recognized that "Kaibab Anticline" was cracked rock when the lake was contained behind it, and that the lake breached catastrophically in the years after the Flood.

<sup>8</sup> Whitcomb and Morris (1961) in their book *The Genesis Flood*, say on page 153, in the caption of a Grand Canyon photo, "Following the Flood, ... the great canyons were rapidly scoured out as the waters rushed down from the newly-uplifted peneplains to the newly-enlarged ocean basins." That reminds us of Byron Nelson's 1931 statement about Grand Canyon "...dug in the first centuries following the Flood." Elsewhere, Morris had written, "...great dammed-up lake full of water from the Flood suddenly broke and a mighty hydraulic monster roared down toward the sea, digging deeply into the path it had chosen..." (Vail 2003). These statements by Morris about post-Flood erosion of Grand Canyon are denied by Michael J. Oard who claims concerning Grand Canyon erosion "...the Flood hypothesis is not new; it was offered by Whitcomb and Morris..." (Oard 2016, 154). Simple reference to Whitcomb and Morris (1961, 153) shows these authors favor rapid post-Flood erosion of Grand Canyon, not late Flood erosion of Grand Canyon, as stated by Oard. This is not a trivial misstatement by Oard because it is often repeated. John K. Reed writes in the "Foreword" of Oard's book, "This book resurrects the concepts of Whitcomb and Morris..." (Oard, 2016, xi). In the "Preface" Oard writes, "This book will flesh out a hypothesis which was suggested by Whitcomb and Morris... that the canyon itself was carved late in the Flood" (Oard 2016, xii). Several pages later Oard develops his thesis of the late-Flood channelized erosion hypothesis, "This idea was introduced by Whitcomb and Morris" (Oard 2016, 7).

<sup>9</sup> Austin attended Northrup's public lecture on the catastrophic-Flood breaching for erosion of Grand Canyon in the summer of 1968 at Lucerne, California. Austin and Northrup also discussed the breached-dam matter when doing geologic field work also in 1968 and 1969. See Northrup (2004).


extending westward from New Mexico through Hopi Buttes a distance of 200 miles to the Kaibab Upwarp.” At that meeting in Pittsburgh, Austin described a “campfire discussion” on closed-basin spillover in April 1985 with geology graduate students while camped at Horseshoe Mesa in eastern Grand Canyon. Two of those geology graduate students wrote term papers on Grand Canyon erosion. John Whitmore’s term paper affirmed evidence for Hopi Lake and the proposition that the lake spilled over the Kaibab Upwarp (Whitmore 1985). Austin had just given a paper at the Pittsburgh meeting on the similarities between Grand Canyon and the newest landscape at Mount St. Helens that was breached by an overtopping mudflow on March 19, 1982 (Austin 1986). At that same Pittsburgh lunch discussion in August 1986, Holroyd explained how, from his Colorado home in Montrose, he frequently conducted field study of the Black Canyon of the Gunnison River. That canyon, he noted, is very similar to Grand Canyon. Holroyd also related that through his government-office computer he had access to a US government digital elevation model (DEM) and that he could plot the lake shoreline that could be contained by the present topography if Grand Canyon was blocked by a gigantic dam. McQueen recalls, “Dr. Holroyd was most interested in how the series of lakes drained, and how drainage could shape landforms upstream of Grand Canyon, especially cliffs on the Colorado Plateau that seemed to lack significant sandstone blocks on the margin with the valley.” Together, these three scientists (Austin, McQueen, and Holroyd) agreed to study these landforms of the Southwest more carefully.

### Mapping a “Big Series of Lakes” on the Colorado Plateau

Late in 1986 Ed Holroyd conducted a survey of the digital elevation model (DEM) data available through the Bureau of Reclamation. These elevation data were originally derived from 1×2° topographic maps at scale one-inch equals six miles (scale 1:250,000) and typically 200ft contour interval. The horizontal geographic resolution (30 arc seconds) was adequate for the regional analysis that was to be performed, but

the elevation resolution ( $\pm 200$ ft from contour maps) was of marginal acceptance for the tasks the computer was being asked to perform. Compare that original data set available to Ed in 1986 to modern data sets. Today’s state-of-the-art elevation databases (modern DEM’s) typically contain laser-aircraft measurements (called LiDAR) of the earth’s solid surface elevation, even through dense forest cover, to horizontal and vertical accuracy of plus-or-minus 1 m (3ft).

In 1986 Ed Holroyd understood the Bureau of Reclamation elevation data to be satisfactory for the regional analysis he intended to perform. Next, he used a scientific programming language called FORTRAN to plot elevations as colors on a geographic grid.<sup>10</sup> He produced two plots, one a regional elevation chart “without lakes” (fig. 5, left), and the other a



**Fig. 5.** Edmond Holroyd’s copy slides of the two original paper elevation maps of the Colorado River drainage basin. Left shows the computer-generated image “without lakes,” and right shows the image “with lakes.” Below both digital images are the scanner copy of the cardboard Kodak mounts with enclosed processed Ektachrome film.

<sup>10</sup> The program title was MTNMAPC.FTN. (These lines are from a copy archived 16 March 1988 in preparation for a job transfer to a different working group in a different city.) Near the start of the program coding is:

```
WRITE(6,102)
102 FORMAT(' If a lake level is desired, type its elevation in meters'
+/ ' mmmm      0 if not wanted')
READ(5,13)LAKE
During production of the map the DEM elevations for each pixel were called MTN(J). Later:
C allow lake colors below nominated elevation
IF(MTN(J).LT.LAKE)THEN
M=21 ; light blue
IF(LAKE-MTN(J).GT.100)THEN
M=22 ; blue
IF(LAKE-MTN(J).GT.200)M=23 ; dark blue
ENDIF
ENDIF
```

The variable M was obviously a plotting code for particular colors, using a call to the plotting routine:

```
CALL SELPAT(KOLORS(M))
```

regional elevation chart “with lakes” (fig. 5, right). The “with lakes” version has a “nominated surface elevation” of 1,700m inserted into the computer code with lake depths shown in shades of blue. Holroyd could have inserted “nominated surface elevation” of 1,800m, but the lake would spill out of its basin. He could have chosen 1,600m, but the lake would only partly fill the basin. Holroyd was also acutely aware of the  $\pm 200$ ft topographic error in the data sets.

How fig. 5 was produced is an extraordinary achievement considering the computer technology available in 1986. Paper output from a color printer was limited to geographic extents of  $2 \times 2$  degrees of latitude/longitude because of the original data sets. So, the  $2 \times 2$  color prints were “tiles” used to assemble a regional mosaic. As one might expect using 1986 technology, “tiles” were pages from the color printer that were cut by hand and taped together into a regional mosaic. Holroyd says, “In 1986 this tile assembly process might be styled as an innovative solution. Today, as we recall it, we laugh and dismiss it as archaic.”

So, with limited technology, Holroyd constructed two charts. The two color-paper, hand-taped mosaics were formed, one chart “without lakes” and one “with lakes.” Original Ektachrome 35mm slide-film photographs of each mosaic (fig. 5, bottom) each display Kodak’s date stamp “JAN87D” embossed into the paper frame. The slide photographs also bear pencil notations “MT-49 and “MT-50,” the slide sequence numbers for Holroyd’s “Missing Talus” oral presentation at the 1990 International Conference on Creationism (Holroyd 1990b). These two Ektachrome slides (fig. 5, bottom) are the January 1987 originals from which are made the transformations by today’s digital-scanning technology (fig. 5, top). Notice that traces are visible of the  $2 \times 2$  degree tile grid from which they were prepared.

### Office Politics

How did Holroyd interface with the Bureau of Reclamation office about his work associated with the DEM? Holroyd says: “In early 1987 I asked a supervisor about the possibility of eventually publishing an article presenting the ancient lakes of the Colorado Plateau as glacial, like Ice Age Lake Bonneville in Utah. I showed the supervisor the color  $8 \times 10$ in prints of the big series of lakes. He responded by strongly threatening with the loss of my job because of unauthorized use of government computer and property. However, it was an extension of my regular DEM work and amounted to only a few extra lines of FORTRAN coding. In fear, I burned the original paper mosaics and the color prints. I only retained the original 35mm Ektachrome slides. These I eventually used in my two oral presentations

at the International Conference on Creationism in 1990. With regard to the DEM work, I had to go into hiding, and I needed to keep quiet about what I had found. I felt like I was carrying the Precious Child in flight to Egypt to avoid murderous King Herod! Yet, to have my findings distributed, I authorized Steve to make use of my lakes outline without mentioning my name as the source.” Austin says, “Both Dave and I recognized the serious problem that Ed was enduring. We honored him by avoiding his name in any description of the DEM work. A coping strategy I had learned, when one encounters a difficult problem, is to tell oneself that the problem could be worse. One day in early 1987, as Dave and I joked, we recalled Ed’s problem. One of us imagined what *could* happen if Sierra Club got wind of what Ed was doing. We jested about a newspaper headline: Bureau of Reclamation Scientist Designs 3,000-Foot-High Dam for New Lake in Grand Canyon.” Yes, that would be a much bigger problem!

In his note in the June 1987 issue of *Creation Research Society Quarterly*, Holroyd wrote, “One could also imagine a series of lakes if the Colorado River was plugged by high ground between the Kaibab and Coconino Plateaus at about the Grand Canyon Visitor Center. A lake surface at about the 1,700m (5,600ft) level could be supported by the present regional topography without the water spilling out over another divide to the north. The resulting series of lakes along the Colorado, Little Colorado, Green, and San Juan Rivers would resemble several of the Great Lakes in size. (Some believe that the sudden release of such a great quantity of water through a fault-generated crack between the north and south rims of the Grand Canyon near the Visitor Center is responsible for the bulk of the carving of the Grand Canyon.)” (Holroyd 1987). Notice that in this June 1987 publication Holroyd was speaking with round numbers and trying to encourage other workers to investigate lakes of the Colorado River drainage basin. Holroyd said nothing about having already *plotted* the lakes on his computer.

### How Does One Describe Lake Elevation?

The U.S. government DEM (digital elevation model) used by Holroyd in 1986 has elevations in meters derived from topographic maps with typically 200ft contour interval at 1:250,000 scale (1in equals about 6mi). The analysis of Holroyd with “nominated surface elevation” of the lake shore plotted at elevation 1,700m (5,577ft). But how much higher could the lake fill before overtopping the modern terrain? That question was directing Ed, Steve, and Dave to imagine the lake’s elevation relative to another hypothetical level called a “pour point.” The DEM, because of the  $\pm 200$ ft errors,



could not supply the terrain pour point answer. We needed much better data than the 1986 DEM. Analysis of the detailed 1:24,000 scale topographic maps of Telegraph Flat area east of Kanab, Utah showed that the water could rise today to about 5,616ft (about 1,712m) before it would spill to the southwest through the modern drainage gap in Moenkopi Formation mudstone that is 20mi east of Kanab, Utah (location in fig. 4).<sup>11</sup> The elevation of the Telegraph Flat pour point is not known exactly. Recent ASTER satellite elevation data indicate the 1:24,000 scale USGS contour map is poorly drawn near the potential “pour point” and contains error of 10ft or more. The recent ASTER data indicate the pour point is actually closer to 5,626ft.

Another, very intuitive way to refer to the lake’s elevation would be simply by the height of the feasible dam consistent with the overall modern topography. The height of the feasible dam should be just above the terrain pour point. Teachers know that talking about terrain this way connects with students.

So, there are three ways to express the lake’s elevation: (1) nominated elevation, (2) pour point elevation, and (3) height of the feasible dam. Austin, Holroyd, and McQueen believe the best way to communicate this elevation is by speaking about the height of the dam. That elevation should be expressed as a round number. Therefore, the elevation of the dam should be rounded to the next 100ft above 5,616ft (the underestimate of the pour point elevation). The question can be stated, “What is the elevation rounded to the nearest 100ft of the dam that could contain the maximum amount of water above present Grand Canyon?” Austin and Holroyd calculated the answer to that question in March 1987 in order to write the caption to a bulletin-board map. The answer is 5,700ft (the rounded elevation of the dam that would form the largest hypothetical lake). Notice, there is only one right answer to the question as it was stated. That one correct answer is 5,700ft. Rounding upward is good engineering practice because the crest of the dam needs to be higher than the lake it is designed to contain. Also, rounding upward was beneficial for office politics reasons because it further distanced the map from Holroyd’s metric computer plot.

### The Bulletin-Board Map

Ed Holroyd recalls how the map was transferred: “I sent a non-colored version of the extents of the lakes to Steve Austin in San Diego in January 1987. Back then we did not have screen-sharing or screen-printing technology, so it was impossible to share directly my computer screen in Colorado with scientists in San Diego. So, I did it the old-fashioned way by exchanging photography through the US Postal Service.”

Austin was interested in Holroyd’s progress in terrain analysis. Austin remembers, “After Holroyd had completed the DEM analysis, he loaned me a black-and-white photographic print of the series of hypothetical lakes on the Colorado Plateau. The print was about 8in wide with washed out details from Ed’s much larger, color rendition. In January 1987 I traced the print photo onto clear acetate and added several familiar geographic features and the state boundaries. I had trouble tracing through washed out detail. Over the tracing I wrote ‘Lake Kaibab’ intending it to depict a series of lakes east and north of the Kaibab Plateau. I promptly returned the print photo to Holroyd by the Postal Service requesting the higher resolution color version. I retained the acetate tracing of Holroyd’s print photo in my archive.” Holroyd provided the 35mm color film through the Postal Service in February 1987.<sup>12</sup> In March 1987, Austin more accurately replotted the shoreline film onto the distribution of familiar topographic features producing for the first time, on a sheet of paper in prepublication form,<sup>13</sup> a user-friendly and thought-provoking lake map (fig. 6). Of course, such a sketch map requires a caption with words describing what the map is attempting to depict. Both Austin and Holroyd composed the caption during a phone conversation in March 1987. Dr. Holroyd remembers he and Austin agreed that the map’s caption would read, “A computer was asked to draw the shoreline of the lake which would form behind the Kaibab Upwarp if the Grand Canyon were blocked at the 5,700ft elevation.”<sup>14</sup> Together in early 1987, Holroyd and Austin composed those words including the elevation 5,700ft.


Because of his association with the government database, and because of the office problem with the boss at Bureau of Reclamation in Denver,

<sup>11</sup> Spillover to southwest at Telegraph Flat would occur at north latitude 37.0659 and west longitude 112.1563 at elevation of 5,616feet (interpolation between 20ft contour intervals on the Petrified Hollow, Utah 1:24,000 scale USGS topographic map). That elevation and position are likely incorrect.

<sup>12</sup> Processing date embossed on the margin of the cardboard slide frame reads JAN87D showing that the Ektachrome film was processed in January 1987 at Denver. Therefore, Holroyd’s DEM study was completed by the end of 1986. Austin received a black-and-white photographic print of Holroyd’s taped color paper mosaic in January 1987. Holroyd’s transmittal letter to Austin/Vardiman with Ektachrome slide film is dated February 26, 1987. Film was returned promptly by postal service to Holroyd after Austin made the tracing. A photocopy of the tracing is in Austin’s archives.

<sup>13</sup> Austin drew the boundaries by hand and Marvin Ross (graphic artist) did the fill in.

<sup>14</sup> When dictating a sentence over the phone, always double check that you are using correct English. The map’s caption has a singular subject (“Grand Canyon”) and a plural verb (“were”). The posted map contained the poor English composition for over two years.



**Fig. 6.** The bulletin-board map that was posted in 1987 for graduate students and faculty. The caption attached to that map read, “A computer was asked to draw the shoreline of the lake which would form behind the Kaibab Upward if the Grand Canyon were blocked at the 5,700-foot elevation.” The data was rendered on Ed Holroyd’s computer (fig. 5, right) and the map was drafted by Steve Austin from the Ektachrome transparency (fig. 5, lower right). The map with caption was first published in March 1989 by Austin, and two different renderings were published by Holroyd in 1990 and 1994.

Holroyd asked that the map (fig. 6) not be published immediately. However, he agreed that Austin and McQueen could talk about the work generally (without mentioning Holroyd as the data source) and post Austin’s sketch map with caption in a non-public venue. Early in 1987 the map with caption was posted on the bulletin board of the Institute for Creation Research Graduate School, in Santee, California.<sup>15</sup> Dave McQueen recalls, “I’ve never seen a bulletin board that compares. It was not attached to the wall but was freestanding in the front of the Geology classroom. As you walked up to it, you recognized it to be a hinged, flip display with several 30×40 in boards forming a ‘book’ of geologic maps. The huge vertical format caught everyone’s attention. It was the 1987 analog of today’s social media; whatever was posted became a topic of conversation. On page 3 of the book of maps was the computer-fit series of lakes on the Colorado Plateau.” That bulletin-board posting focused attention on the question, “How were the ancient lakes configured?” The bulletin-board map suggested the series of lakes could extend into four states (Arizona, Utah, New Mexico, and Colorado) being as big or bigger than Lake Superior!

In June of 1987 Holroyd published his short note on “Missing Talus” to stimulate new research

(Holroyd 1987, 15, 16). Gravity causes accumulation of boulders at the bases of slopes below cliffs that form steeply inclined rock piles called talus. If slopes backwear slowly during millions of years, the base of associated slopes should have substantial talus. Holroyd says: “I pointed to shale slopes with sandstone capping strata at Mesa Verde, Grand Junction, Book Cliffs and Monument Valley. Sandstone blocks should litter the shale valley floors. Where did those blocks go? One possibility is that a lake shore was up against the slope with wave action abrading the boulders. Another possibility is that catastrophic drainage of lakes could have swept sandstone blocks from those valleys. In my 1987 paper I stated in words that one could imagine “a big series of lakes” on the Colorado Plateau. What I didn’t say was that I had already plotted the possible lake shoreline on my computer! By publishing this mid-1987 note in *Creation Research Society Quarterly*, I notified the creation community of what I discovered in late 1986 by this Colorado Plateau terrain analysis. Thereafter, anyone could reproduce the lake simulations.”

Early in 1988, Ed Holroyd was transferred to a different working group within Bureau of Reclamation. Specified in Ed’s new job description was doing remote sensing and mapping research.

<sup>15</sup> In 1987 and 1988 the yet-to-be-published “blocked at 5,700-foot elevation” map was simply referred to in graduate school conversations as “the Colorado Plateau lake map,” or occasionally as “Austin’s map,” but never as “Holroyd’s map.” McQueen in his archive has a 1987 classroom photo that shows a partial view of that bulletin board.

Also, in early 1988, Ed transferred the DEM computer code to his home computer taking the work away from government computers. Talk about the “big series of lakes” and “blocked at the 5,700ft elevation” changed in the spring of 1988. Holroyd had made his home-office computer render DEM and satellite imaging software, now making new home-office plots clearly in the public domain. So, Austin requested to publish the “blocked at 5,700ft elevation” map (letter to Holroyd, January 23, 1989) and Holroyd (letter to Austin, February 2, 1989) gave permission. Finally, after being posted on the bulletin board for two years, the “blocked at 5,700ft elevation” map was distributed by publication in March 1989.

### Tectonic Tilting of Kaibab Upwarp

Also, in the summer of 1987, graduate school classes continued in San Diego with faculty and students expressing their opinions on the bulletin board map. Austin suggested that the actual ancient configuration of post-Flood lakes was somewhat different than suggested by the map. According to Austin, important topographic change occurred by tectonics and isostasy after drainage of the lakes. Understanding the tectonic change since the lakes existed, allows us to visualize more accurately the original lake basins and their bedrock dams.

As one stands on the South Rim at Grand Canyon Village and looks north and northwest, the strata of the Kaibab and Kanab plateaus on the far side of the Colorado River appear to be perfectly flat and level with the horizon. That is the iconic image that we all have of Grand Canyon. However, that impression of strata flatness is an illusion created by our viewing angle. If one pays attention to the Kaibab Formation at the Canyon rim and follows it northward through the Kanab and Kaibab plateaus, that formation decreases over a thousand feet of elevation before reaching the Utah border! Therefore, the Kaibab Formation is really inclined, dipping northward at over 50ft per mile (over 10m per km) through a distance of 40mi. All of the plateaus in Arizona north of the Colorado River have substantial northward inclination (up on the south sides, down on the north sides). Austin considers Kaibab Upwarp and most of the monocline structure of Kaibab Plateau formed *before* Canyon erosion. That early formed structure confined Hopi Lake topographically. However, Austin suggested that a major part of the *tilting* of the Kaibab Plateau occurred *after* spillover erosion of Grand Canyon. Therefore, Austin suggested that more than 1,000ft of present elevation needs to be *removed* from the south end of Kaibab Plateau to approximate the configuration of the East Kaibab Spillover when the structure was breached. Also, Austin proposed that more than 1,000ft of elevation

needs to be added to the north end of Kaibab Plateau near Telegraph Flat at the time of spillover. As early as 1987 Austin postulated rotational plateau tilting (up on the south, down on the north) was likely associated with the oblique strike-slip shear that continued after monocline flexure. That was Austin’s “tectonic tilting hypothesis” for Kaibab Upwarp formulated in late 1987. Geologists, lately, are still discussing the tectonic process, especially the “Pliocene and Pleistocene uplift” of the southern margin of the Colorado Plateau.

In addition to *tectonic uplift* of the south end of the Kaibab Plateau, one needs to imagine likely hundreds of feet of *isostatic uplift* because of the removed weight of sediment within the eroded canyon and plateau. Isostatic uplift also likely occurred by the removal of the weight of upstream lakes. Austin comments: “A noteworthy cluster of recent earthquakes occurs under the Kaibab Upwarp. Continued faulting appears to have uplifted the southern Kaibab Plateau many hundreds of feet after the Kaibab dam was breached. Some elegant structural geologic models have been developed to explain the recent northward tilting of the Kaibab Plateau that allows us to visualize the original configuration of the Kaibab dam.”


### There’s Power in the Flood!

Among the participants in the April 10, 1988 rim lecture at Grand Canyon was Paul MacKinney. Paul was a concrete engineer from Illinois with a special interest in Grand Canyon erosion. As early as 1985 Paul was calling attention to spillway erosion, especially the process of cavitation, whereby fluid vacuum bubbles form in high-velocity floods. Paul pointed out how shallow, high-velocity flow creates very low-pressure fluid around channel obstructions, creating vapor bubbles within the fluid. Those bubbles are vitally important because they *implode* inflicting explosive forces on rock spillways. The process of cavitation can produce greatly accelerating bedrock erosion on a colossal scale. As a real-world example of cavitation erosion, Paul pointed out the catastrophic failure of the concrete-and-steel-reinforced left spillway tunnel at Glen Canyon Dam (location in fig. 3). In June 1983, during an emergency water release, discharge of 93,000 cubic feet per second was sustained through the left tunnel. During that emergency release, earthquakes were felt within the dam, and the “rooster tail sweep” exiting the tunnel turned distinctly red! After shut-down of the spillway tunnel, engineers found at an elbow within the tunnel a new 63,000-cubic-foot-volume hole penetrating concrete, steel and bedrock. Paul believed the enormous erosion in the left spillway tunnel could have happened because of cavitation within seconds.

Paul wanted to stimulate technical study of cavitation so he encouraged his creationist friends Cliff Paiva and Ed Holroyd. Cliff Paiva completed his master's thesis on cavitation in 1988 (Pavia 1988). As a physicist with Bureau of Reclamation in Denver, Holroyd closely followed the government's study of cavitation after the 1983 Glenn Canyon Dam cavitation event (Falvey 1990). Holroyd wrote computer programs to simulate cavitation erosion on scales never observed by humans (Holroyd 1990a). Just below No Name Point in Grand Canyon is Papago Creek. Holroyd wrote computer simulations of spillover erosion down Papago Creek through a channel 1,000ft wide with initial flow speed of 10m per second (Holroyd 1990b). Catastrophic bedrock erosion occurred in the simulation. After 1985, popular breached-dam lectures on Grand Canyon erosion often described how cavitation can erode spillways catastrophically. After one lecture on the Canyon Rim, an attendee was humming a familiar Christian hymn, then he sang the concluding words: "There's Power, Power, wonder-working Power, in the overflowing flood of the dam!"

### Holroyd's Backyard

Dr. Ed Holroyd lived in Montrose, Colorado, from 1983 to 1988, adjacent to what is now the Black Canyon of the Gunnison National Park. That region, which displays the amazing course of the Gunnison River in southwestern Colorado, is displayed in fig. 7. The Gunnison River drainage begins in the topographic saddle in the upper right corner of fig. 7 between the West Elk Mountains (upper left of figure) and the San Juan Mountains (off the upper right corner of fig. 7). Then, between the Red Rocks and Cimarron faults, the Gunnison drainage turns abruptly northwestward (right side fig. 7) and enters the extraordinary gorge of Black Canyon. Then, after departing Black Canyon, the Gunnison drainage turns northward (foreground of fig. 7). Finally, the Gunnison River turns westward again, exiting in the front left corner of fig. 7. Holroyd says: "I sought to understand how Gunnison River was established across this fascinating landscape. I enrolled in the college course *Geology of Southwestern Colorado* offered by what was then Western State College, Gunnison. From that class and working with hand analyses of geological and topographic maps of



**Fig. 7.** The amazing course of the Gunnison River in southwestern Colorado. Foreground width is 18mi. View is here toward the southeast. Ed Holroyd noted that Black Canyon of the Gunnison (right-middle of this view) is positioned through elevated terrain that is aligned parallel to the axis of arched strata between two faults. Ed also noted a topographic low at "Cimarron Spillover" that appears to support his "crack-and-capture hypothesis." Ed Holroyd's home in Montrose, Colorado is just off the lower right corner of this view. (Landsat 8 photo acquired September 24, 2013.)

the Black Canyon region, I sought to understand the geologic history of the canyon and its region. I read carefully the explanation published by USGS geologist Wallace R. Hansen (Hansen 1965). I even discussed the landscape with Wally Hansen in 1986. Both the college course and the USGS geologist promoted the headward erosion and antecedent “old river” hypothesis to explain how the Gunnison River eroded upstream along the same course from the ancestral Gunnison River. I recognized that explanation is the same as the classic twentieth century story for Grand Canyon!” So, fig. 7 shows the magnitude of this river course location problem.


One detail of the Gunnison landscape fascinated Holroyd. That is the turning point, where between the Red Rocks and Cimarron faults, the Gunnison River enters Black Canyon. There appears to be a low point in the terrain as shown in fig. 8. Holroyd comments on fig. 8: “I was impressed by oblique views in overflight. Gunnison River should have maintained its westerly course out of the high country into the incredibly soft Mancos Shale at the village of Cimarron. That should have occurred if the classic ancestral river story is correct. I noticed in overflight that the strata at the entry of Black Canyon form a broad arch structure with the river and canyon parallel at the top of the arch between the two faults. What I saw convinced me that overtopping occurred at what we can call Cimarron Spillover. These observations led me to propose cracks in the bedrock. An arch structure is where one would expect tensional cracks that open upward, and, because of overtopping, could direct erosion into high country.” That was the genesis of what Holroyd calls his “crack-and-capture hypothesis.”

### Lees Ferry Spillover


During July 1987, Holroyd drove from his home

next to the Cimarron Spillover in southwestern Colorado to Marble Canyon, Arizona, just northeast of Grand Canyon. His paper “Missing Talus” had just been published and observations of Gunnison landscapes were on his mind. Holroyd was impressed with the central location of Lees Ferry between the upper and lower Colorado River basins (location in fig. 3). At Lees Ferry, Arizona in July 1987 he inspected Moenkopi Shale slopes below classic Shinarump Conglomerate cliffs. In fall 1987 Holroyd drew two sketches from slides of cliffs at Lees Ferry without significant talus (fig. 9). Here is how Ed explained that cliff in a phone conversation with Steve Austin: “The phone conversation was in December 1987. I explained to Steve my computer DEM plot of Prospect Lake that had formed within Grand Canyon by the lava dam at Vulcans Throne. Prospect Lake as I plotted it on present topography reached to Lees Ferry. My original thought before my July 1987 field trip was that the lake’s shoreline stood at the base of the cliff above Lees Ferry. Then, I saw the missing talus confirmation in July 1987 during my field work. Steve suggested on the phone a bigger lake upstream that spilled over and swept talus away. So, I was ready to describe the spillover point at Lees Ferry in a research proposal that I wrote in January 1988.”

Fig. 10 shows the spillway at Marble Canyon and Holroyd’s new understanding of its configuration. Austin describes how Holroyd explained it: “In a December 1987 phone conversation, Ed was excited about his ‘crack-and-capture hypothesis.’ The Colorado River channel at Marble Canyon that Ed observed in July 1987 was oriented down the axis of an uplifted arch structure just like the channel at the upstream end of Black Canyon of the Gunnison River that he observed in 1985 and 1986. I told Ed that I liked the way he was thinking about spillover



**Fig. 8.** Oblique aerial photos above “Cimarron Spillover” on the Gunnison River. Left image is looking northwest above the spillover where Gunnison River enters from the bottom right, turns abruptly to the northwest and enters upper Black Canyon. Right image is looking southwest above the village of Cimarron and the spillover where the Gunnison River should have continued westward into the soft shale in the lowland (upper right), but instead turns into upper Black Canyon (lower right). Ektachrome slide photos by Edmond Holroyd. Left image is from overflight on 29 March 1986, and right image is overflight on 23 June 1987.



**Fig. 9.** Ed Holroyd sketched the talus slope at the base of the cliff at the pullover parking area between Marble Canyon and Lees Ferry. Hard and grey Shinarump talus boulders are a remarkably thin deposit lying on the soft and red Moenkopi Shale. Holroyd recognized in July 1987 that paucity of talus at Lees Ferry argues this cliff did not backwear through millions of years, but that the slope was recently swept by a catastrophic spillway flood. The two sketches were drawn from Ektachrome film by Holroyd in fall of 1987.

at Lees Ferry into Marble Canyon and mentioned that a geology graduate student might want to do a research thesis on such a topic. I encouraged Ed to submit a written proposal so that a student can easily explore the idea. I requested that the proposal contain a clear and positive hypothesis that a student might test.”


The three-page proposal was titled, “Some Research Opportunities at Marble Canyon for Creationists (written January 1988 from observations in July 1987)” which Austin received with Holroyd’s cover letter dated January 24, 1988.<sup>16</sup> Holroyd’s thesis proposal includes, “...the existence of Lake Bidahochi means that there were similar ‘Great Lakes’ throughout the entire Four Corners region and into Wyoming.... When one stands on the hillside north of the Marble Canyon Lodge and looks southwest over the valley containing the Colorado River, the doming of the strata is obvious. The Colorado River has chosen a bed in the crest of an anticline. Rivers naturally chose lower rather than higher ground for their beds. This means that the top

of the anticline developed a crack which captured the Colorado River....the channel was dug by the process of cavitation resulting from the catastrophic release of water from a large lake upstream through that crack....Perhaps the catastrophic release of three Great Lakes-worth of water carved the Grand Canyon in a few weeks by means of cavitation process rather than in millions of years.” Holroyd’s perspective of Marble Canyon drainage spillway after the December 1987 phone conversation might be best portrayed by an oblique view above Lees Ferry Spillover (fig. 11).

### Where Are the Lake Deposits?

If an enormous lake breached a natural topographic dam through the east side of the Kaibab Upwarp, one would expect to find evidence in sediment diagnostic of such a lake. Quiet lake water traps wind-blown silt and clay, and the dissolved minerals in the water could precipitate calcium carbonate particles to form a type of limestone called tufa. Austin recalls, “The quiet water of a lake could allow accumulation of clay

<sup>16</sup> Austin considered Holroyd’s January 1988 proposal to be confidential, only to be communicated to graduate students who expressed special interest. Holroyd’s original proposal “Some Research Opportunities at Marble Canyon...” and Holroyd’s original signed cover letter are retained in Austin’s archive.



**Fig. 10.** Observations of Marble Canyon spillway allowed Ed Holroyd to understand a “crack-and-capture hypothesis” for catastrophic drainage of a big Utah lake. Note that the axis of the channel marks the crest of the arch structure. Inset Ektachrome photo was acquired by Holroyd on the talus-strewn slope above Marble Canyon village during field work on July 14, 1987.

and calcium carbonate particles just east of Grand Canyon. If such sediment layers exist, it would be like court-room reports of a smoking gun at the scene of a crime.” Austin further describes his thinking, “In 1987 we were suggesting that late-Flood and post-Flood uplift of the Colorado Plateau trapped water in the saucer-shaped depression on top of the Colorado Plateau. It would be a lake formed after the Global Flood of Noah’s day but just as the Ice Age began. Kaibab Upwarp is the key topographic barrier to retain a lake in northeastern Arizona as the lake map attempts to depict. If the early post-Flood Hopi Lake existed just above 6,000ft elevation in Arizona, then its presence would likely obligate a huge lake or lakes in Utah and Colorado.”


For several years Austin had been searching for silt and limestone deposits that might be critical evidence that Hopi Lake stood just above 6,000ft elevation on the east side of Kaibab Upwarp. Austin studied calcium carbonate deposits in the Cape Solitude area on Navajo lands just east of Grand Canyon and other deposits adjacent to Blue Moon Bench (see fig. 4). These deposits resemble the shoreline lake limestone

called tufa. Robert Scarborough, a geologist who conducted graduate research on the Hopi Buttes silt, agreed with Austin. Scarborough had also been looking for ancient lake sediment east of Grand Canyon. Together, in 1988, they affirmed privately that ancient Hopi Lake was impounded just east of the Kaibab Upwarp, and that spillover of that lake likely eroded Grand Canyon.<sup>17</sup>

#### Spillway at Mount St. Helens

What would the spillway look like if lakes catastrophically drained by spillover through the highlands of northern Arizona? That was a vital question that geologists were asking 30 years ago. Austin recalls, “The largest landslide deposit in human history has a volume of two-thirds cubic mile and occupies an area of 23 mi<sup>2</sup> on the north side of Mount St. Helens. That historic-record debris deposit formed as the volcano exploded on the morning of May 18, 1980. Since then, Mount St. Helens volcano has provided a laboratory for the study of catastrophic erosion. I started doing field work on spillover erosion at Mount St. Helens in 1983. I described in public

<sup>17</sup> Austin describes a phone conversation in 1988: “I was talking with Bob Scarborough. I asked him if he knew of other professional geologists who believed in overflow origin of Grand Canyon. There was an awkward moment of silence. Were there any others? I knew of Cliff Burdick. Neither of us knew Norman Meek at UCLA, but he had just rediscovered Afton Canyon and likely was thinking about similarity to Grand Canyon.” Obviously, 1988 was a *very small* beginning for spillover thinking.



**Fig. 11.** Oblique aerial view of Marble Canyon Spillway from just above Lees Ferry Spillover. The view is toward the southwest. Field work in July 1987 allowed Ed Holroyd to propose that Lees Ferry Spillover was the overflow location of a dam of the “big Utah lake.” Later, Brown (1989) also favored Lees Ferry being a drainage spillway. Image rendered by Ed Holroyd using Google Earth software.

lectures and in publications how the mudflow on March 19, 1982 breached the landslide debris north of the volcano, especially an elevated landslide debris dam. The breach was not straight through the debris dam but has a curving-to-the-right path. In significant ways the spillway of the breached dam at the volcano resembles the spillway below No Name Point in eastern Grand Canyon.” Austin pointed out in publications similarities as early as 1984<sup>18</sup> and later published in peer-reviewed geology publication a description of the “Little Grand Canyon” in association with the breached landslide debris dam (Austin 2009). When observed just downstream of their points of spillover (see fig. 12), both landscapes are characterized by U-shaped canyon cross-sections, modern meandering channels, amphitheater-headed alcoves, steep cliffs and elevated upland flats. The unbranched mud spillway just downstream of the breached explosion pit at Mount St. Helens is very similar to the unbranched eastern Grand Canyon through the Kaibab Upwarp.


As early as 1984 Austin was pointing out that there is much more to the correspondence between Mount St. Helens and Grand Canyon than just the unbranched spillways of these breached dams. As early as 1984,

Austin called attention to three erosional areas at Mount St. Helens: (1) the breached dam and its prominent unbranched spillway, (2) the downcut landscape above the breached dam with its distinctive rill-and-gully topography, and (3) the drainage-dissected landscape with prominent tributary canyons downstream of the spillway. Upstream of the breached topography at Mount St. Helens is the large basin with 700m long steam-explosion pit where mud pooled temporarily behind the barrier. After breaching, that upper basin has been downcut a depth of 100ft (30m) displaying new channels with fluting and rill-and-gully erosion. Those oversteepened channels and heightened topography in the upper basin at Mount St. Helens resemble some of the “canyonland topography” in Marble Canyon and Glen Canyon areas on the upper Colorado River. Also, downstream of the unbranched spillway at Mount St. Helens is the region where multiple side canyons enter the North Fork of the new upper Toutle River (Austin 2009). These downstream tributary drainages also appear to be related to the main mudflow breaching event in March 1982. In the upper left corner of fig. 12 (left), a large side canyon joins the channel just below the spillway (Scheele 2010).<sup>19</sup> These big side canyons at Mount St. Helens are similar to Kanab Creek

<sup>18</sup> Soon after the 1980 eruptions of Mount St. Helens, Austin was describing spillover erosion (Austin 1984).

<sup>19</sup> The caption of Scheele’s Figure 3 says: “The edges of the ‘Little Grand Canyon’ at Mount St. Helens are relatively straight and do not exhibit the branching structure of the Grand Canyon.” The figure at its center shows what Austin (2009) calls “the old channel of the main drainage” where it joins “the new channel of the main drainage” downstream of the unbranched spillway segment (left side of Scheele’s Figure 3). Thus, Scheele’s Figure 3 shows branching canyon structure downstream of the unbranched spillway and makes it an analog to side canyons like Kanab Creek and Havasu Creek in central Grand Canyon! Obviously, Scheele was unaware of Austin’s 2009 publication. Journal editor Tas Walker should get better peer-review to correct during review the unsupported statement “... the canyons of Mount St Helens (Figure 3) do not show branching structure exhibited by the GC.”





**Fig. 12.** Spillways at Mount St. Helens (left) and Grand Canyon (right). Both images are views from above their spillover points looking downstream. Mount St. Helens spillway (left) is on the North Fork of the Toutle River and is eroded by spillover 180 ft deep through unstratified May 18, 1980 debris avalanche deposit by mudflow of March 19, 1982. Grand Canyon spillway (right) is just downstream of the breach in the margin of Hopi Lake and is 4,500 ft deep eroded by water through distinctly stratified sandstone, shale and limestone to form the present channel of the Colorado River. These landscapes, just downstream of their points of spillover, are characterized by U-shaped canyon cross-sections, modern meandering channels, amphitheater-headed alcoves, very steep cliffs, and elevated upland flats. Grand Canyon spillway (right) has No Name Point in upper center. Grand Canyon spillway is 25 times deeper than the Mount St. Helens spillway. Photos copyright Steven A. Austin 2019.

and Havasu Creek (see fig. 4) in the central Grand Canyon.<sup>20</sup> Thus, some unanticipated similarities relate historic spillover erosion at Mount St. Helens to Grand Canyon. That would be expected if spillover is a good working hypothesis.

### The Forgotten “Grand Canyon” of the Mojave Desert


An awkward moment occurred in the summer of 1930 at Stanford University in California. Dr. Eliot Blackwelder, chairman of the Geology Department, was opening graduate school, but where was that new geology student named Elmer Ellsworth? When 23-years-old Ellsworth showed up late to school, he appeared to have an adequate geologic excuse. As he crossed the country in his Model A Ford from Wisconsin, he detoured to Grand Canyon. His mule train caused him to be delayed *within Grand Canyon!* (Ellsworth 1999).<sup>21</sup> Ellsworth’s uneasy start at Stanford was soon forgotten when he explained to his new professor how his graduate work at University of Wisconsin

had revealed sedimentary evidence of an Ice Age lake in central Wisconsin. That’s when Eliot Blackwelder recognized he had, contrary to first impression, a promising student that could do his Ph.D. dissertation study on an ancient Ice Age lake in Mojave Desert of Southern California. As Blackwelder described his spillover hypothesis for the entire Colorado River drainage basin, his new student recognized that the hypothesis could be tested at ancient Lake Manix and Afton Canyon on the Mojave River (Afton Canyon located in fig. 3, with Lake Manix reconstructed in fig. 13). Lake Manix had three prominent basins: (1) Coyote Basin on the northwest, (2) Troy Basin on the south, and (3) Afton Basin on the northeast.<sup>22</sup> Over the next two years Ellsworth carefully mapped the Ice Age lake’s sediments, interpreted the lake’s history, and turned in his Ph.D. dissertation, on time (Ellsworth 1932). Elmer Ellsworth made his professor proud! In 1936 the student and his professor coauthored a short description of their discovery in a peer-reviewed geology journal (Blackwelder and Ellsworth 1936).

<sup>20</sup> Does the spillover hypothesis hold water? Oard (2016) says “No”. A “fatal problem” with the spillover hypothesis, according to Oard, is “long tributary side canyons cannot be explained by a dam breach” (Michael J. Oard, recent online document <http://michael.oards.net/pdf/PostFloodBoundary/Chp38version3.pdf>). Yet, at Mount St. Helens, during the 1982 mud breaching event “long tributary side canyons” were produced downstream of the unbranched spillway. An extraordinary example of a “long tributary side canyon” is shown in Austin (2009, 350 and 351). Peter Scheele and Mike Oard need to become fully aware of the landscape features associated with the breached dam at Mount St. Helens. Both authors could benefit from peer review of their writings about Mount St. Helens landscape.

<sup>21</sup> During 25 years of work for American Association of Petroleum Geologists in Tulsa, Oklahoma, Elmer Ellsworth and his wife Helen were members of First Presbyterian Church (Reformed Presbyterian affiliation). He became ruling elder. When he retired in San Francisco Bay Area, he created a jail ministry program.

<sup>22</sup> Some geologists note that Lake Manix in map view resembled a “three-leafed clover.” Ed Holroyd, who drew fig. 13, thinks his map of Lake Manix looked like a “fire-breathing pterosaur.” As the pterosaur swooped eastward, it breathed fire on Afton Canyon!



**Fig. 13.** Mojave River drainage east of the city of Barstow, in San Bernardino County, California. Ice Age Lake Manix (elevation 543 m) breached its dam at Afton Canyon, eroded its spillway, and deposited outwash eastward (downstream) in Soda Lake basin. Shades of blue depict ancient Lake Manix. Deepest eastern arm of Lake Manix is Afton Basin where deepest erosion is evident. Red depicts Mojave River bedrock canyons, with short red segment directly east of deep Afton Basin being the famous Afton Canyon spillway. Semi-transparent pink shades depict braided delta deposits. (Map drawn by Ed Holroyd from DEM onto satellite image base with the specified elevation and spillover of Lake Manix.)

In the 1930s elevation control and topographic contour were much less accurate than today's data sets. So, in retelling the Lake Manix story, we use here newer and improved topographic control. Ellsworth believed that the lake shore was at elevation 1,781 ft (543 m). The surface area of Ice Age Lake Manix was almost 90 square miles, and the lake volume was about three-quarters cubic mile. Ellsworth used shoreline and lake-bed evidence to understand that the lake drained catastrophically by spillover. Greatly enlarged by flood erosion, Afton Canyon is now 440 ft deeper providing a new channel for Mojave River eastward through a mountain into Soda Lake basin. The uppermost one mile of the four-mile-long spillway (fig. 14) is eroded about 440 ft deep through bedrock within Afton Canyon. A major part of that deepening, according to Ellsworth, occurred because of spillover. The lower half of Afton Canyon (fig. 15) when viewed toward the east is less imposing. The arm of Lake Manix immediately adjacent to the breached dam is called Afton Basin, and it contains alluvial deposits,

especially conglomerates, now deeply channeled (figs. 16 and 17). Erosional landforms within Afton Basin have been referred to as “canyonland topography” (Meek 2019)<sup>23</sup> that resembles, both in appearance and variety, landforms upstream of Grand Canyon on the Colorado River. Afton Basin, remarkably, lacks fine-grained lake deposits.<sup>24</sup> Those Afton Basin erosional landforms resemble those in the upper Colorado River drainage basin, but are roughly at one-tenth scale. By 1936 it seemed like spillover was a good working hypothesis that could stand on its own!

The 1930s were not the best time to publish the lake overflow explanation for erosion of Afton Canyon. Ellsworth could not be employed by academic institutions because of the Great Depression, and, so, he went to work for the petroleum industry. Then, World War II came and Ellsworth joined the Air Force. Blackwelder had other projects with students demanding attention at Stanford. He personally was dealing with health issues, and other geologists seemed preoccupied in telling elaborate stories about

<sup>23</sup> On page 176, Meek notes, “Canyonland topography, albeit in relatively soft sediments, has formed >60 km upstream of Afton Canyon in <13,700 yr.” Meek understands “...canyonlands that result after a barrier is breached.”

<sup>24</sup> Meek has also noted general erosion of the lake's bed and missing fine-grained lake sediments generally in Afton Basin of the eastern arm of Lake Manix. To find good fine-grained lake sediments from ancient Lake Manix, according to Meek, one needs to go to the western region of the ancient lake. A similar observation of sediment distribution is made in Arizona at the Bidahochi Formation of ancient Hopi Lake. Fine-grained lake sediments are noticeably missing on the dissected “canyonland” bed of the ancient Hopi Lake just east of the Kaibab Upwarp.



**Fig. 14.** Afton Canyon looking westward into the topographic basin of ancient Lake Manix. Eliot Blackwelder and Elmer Ellsworth in 1936 understood Afton Canyon to be the spillway from catastrophic drainage of Lake Manix. Here the spillway was eroded through bedrock to a final depth of 440 ft. That makes Afton Canyon a one-tenth scale example relevant to the Grand Canyon debate. Union Pacific Railroad tracks are on the north side of the canyon. Photo by QT Luong, terragalleria.com, copyright 2019. Used by permission.



**Fig. 16.** View of Afton Basin from above the spillover point of Afton Canyon looking upstream (west) into the dissected bed of ancient Lake Manix. Several geologists have used the term “canyonland topography” to describe the unique assortment of flood erosion landforms in the old bed of Lake Manix. Drainage of Lake Manix occurred through the spillway in foreground. Photo by QT Luong, terragalleria.com, copyright 2018. Used by permission.

headward erosion of Grand Canyon. For the next 50 years little was said about Mojave River spillover. It was virtually ignored. Then, in the late 1980s a geology graduate student named Norman Meek at University of California Los Angeles revisited the evidence for Lake Manix. Meek said he was “shocked” to discover that spillover had been so well stated by Blackwelder and Ellsworth 50 years before, and he said he was greatly concerned as to why it had been virtually “forgotten.” Meek started publishing on Lake Manix and Afton Canyon (Meek 1989). He became a crusader for spillover. Three lines of evidence from Lake Manix



**Fig. 15.** Downstream end of Afton Canyon looking east. Catastrophic drainage of Lake Manix downcut the extreme east end of Afton Canyon, with the canyon dying out abruptly into the next downstream basin. In the canyon’s place downstream is an enormous outwash plain of flood debris entering Soda Lake basin near Baker, California. Photo by QT Luong, terragalleria.com, copyright 2019. Used by permission.



**Fig. 17.** Erosion in Afton Basin, just upstream of the breached dam at Afton Canyon. This “canyonland topography” was eroded in conglomerate of the bed of the Lake Manix during and after the flood’s drainage. Stream channel in front left is about 8 ft wide. Photo by QT Luong, terragalleria.com, copyright 2019. Used by permission.

promoted spillover: (1) the abrupt disappearance of the lake from its highstand elevation, (2) the Afton Canyon spillway and its upstream dissected “canyonland,” and (3) the thick and coarse buried flood sediment layer encountered in water wells downstream of Afton Canyon. In 1990 Meek completed his doctoral thesis on Lake Manix (Meek 1990).<sup>25</sup>

### The Afton Canyon Controversy

One might suppose that the rediscovery of the spillover hypothesis for Afton Canyon would now be received eagerly by the geology establishment. After all, the establishment should be seeking to assemble pieces of a great puzzle to solve important

<sup>25</sup> Meek’s dissertation is dedicated to Elmer W. Ellsworth.

problems like erosion of Grand Canyon. Add to that, *two* doctoral dissertations supported the hypothesis. Instead, rebuttal papers were written to dispute spillover in the Mojave River drainage basin (Enzel, Wells and Lancaster 2003; Wells and Enzel 1994; Wells et al. 2003). Three authors argued that the bedrock obstruction at Afton Canyon was eroded much more slowly from the east by an upstream migrating gully, not very much by an overtopping lake from the west (Enzel, Wells, and Lancaster 2003). The lake, according to spillover critics dried up slowly, leaving behind a progressive series of lower shorelines as Lake Manix finally disappeared. However, critics did not give specific locations of those recessional shorelines. Somebody asked the question: could they be arguing just from elevation measurement errors?

As the twenty-first century began, Lake Manix's history became embroiled in heated discussions which could be called the "Afton Canyon Controversy." An initial report by U.S. Geological Survey attempted to avoid the strong disagreements (Reheis and Redwine 2008). U.S. Geological Survey became available to conduct extremely detailed remapping in Afton Basin, the eastern bed of the big lake (Reheis et al. 2014). USGS used differential corrected GPS and LiDAR to measure shoreline elevations with errors less than one meter. After remapping by USGS, Lake Manix's history was "...punctuated by tectonic movements and a catastrophic flood that reconfigured the lake basin" (Reheis et al. 2014, 1). USGS had independently confirmed the Blackwelder, Ellsworth and Meek version of spillover! Critics appeared to be rebuffed. Then, the critics received criticism. Afton Canyon spillover critics are described as having a flawed method of educating geologists. This rebuttal unveils "a pedagogically engrained bias," and reveals, "...the continued omission of spillover as a possible transverse drainage hypothesis hints at a larger problem related to selective textbook content and a constrained paradigm that does not inform adequately about fundamental river development mechanisms" (Hilgendorf et al. 2020). That's fancy academic language describing *educational bias*. Is "Afton Canyon Controversy" an episode to inform us about bias one encounters in the much larger "Grand Debate" concerning erosion of Grand Canyon?

### Take a Look at My Backyard!

Four geologists open their 2009 geomorphology

research paper with the thought-provoking sentence, "The study of how rivers cross obstructing mountains, once popular in the early twentieth century, has seen a dramatic resurgence in the last decade" (Douglass et al. 2009). Researchers had worked out a technical "checklist" or "logic tree" for understanding rivers that cross mountains (Larson et al. 2017).<sup>26</sup> Once the options and methodology were stated technically, many of these scientists deliberately attempted to generate their own applications of "spillover" to landscapes where they were living. An earth scientist speaks about his or her "backyard" with enthusiasm and passion, often speaking with authority, even with a hint of pride. This emphasis on the geology of where the earth scientist lives has been called the "backyard effect."

It is easy to see the "backyard effect" in our story of Grand Canyon erosion. Remember that Ed Holroyd was living in Montrose, Colorado when he became fascinated by the Black Canyon of the Gunnison River. Then, he applied his thinking from the Gunnison River to the Grand Canyon spillover erosion problem. Another noteworthy "backyard" example is Dr. Norman Meek. He rediscovered catastrophic drainage of ancient Lake Manix explains erosion of Afton Canyon running through mountains along Mojave River (just northeast of his home in San Bernardino, California) (Meek 1989). Similarly, cooperation among eight researchers (Larson et al. 2014)<sup>27</sup> promoted understanding that lake overflow of Pemberton Basin established the modern course of the Salt and Verde rivers (north and east of their homes in Phoenix, Arizona). Of those eight Arizona researchers, only Phillip Larson was *not* living in Arizona at the time of publication. Larson had moved from the desert landscape of Phoenix back to his original home in the glacial landscape of Minnesota, where he found long-appreciated evidence of "spillover in glacial/proglacial environments" (Hilgendorf et al. 2020, 9–12). Many other earth scientist examples of the "backyard effect" could be cited.

### Reviving the "Grand Debate"

The forgotten transverse drainage hypothesis was proving itself, especially by the backyard effect, in technical thought and literature to have explanatory power! Then, an interesting chain of events happened. Lake spillover thinking transferred (would "overflowed" be a better word?) from technical science journals to Internet news releases, and finally, to television documentaries. A 2012 Internet

<sup>26</sup> Grand Canyon is featured in this paper.

<sup>27</sup> Larson, the senior author of this paper, is the only author on this paper who was not resident of the Phoenix area at time of publication.

<sup>28</sup> View the National Geographic documentary here: <http://www.youtube.com/watch?v=SeBPKE5eDU0>. The model of Dr. Douglass is 1:60,000 scale with ten times vertical exaggeration. Notice as the big lake drains in the model, it becomes two lakes with an elevated bench in between. Then, as Grand Canyon erodes downstream, the Little Colorado River and Marble Canyon are both eroding upstream. Note that topography at the first spillover causes the eastern Grand Canyon to form the broad sweeping, racetrack turn from south to west.

news release described Grand Canyon erosion with the lake-carved-the-Canyon theory as, “A favored concept for two decades” (Oskin 2012). The 2008 National Geographic made-for-TV documentary “Grand Canyon Spill-Over Theory” features Dr. John Douglass and his stream table experiment at scale 1:60,000 that modeled Grand Canyon’s overspilling lake.<sup>28</sup> A second stream table experiment of lake spillover by Douglass appears in the History Channel made-for-TV documentary “How the Earth Was Made—Grand Canyon” (2009, season 2, episode 1).<sup>29</sup> Detailed parameters of the spillover of Hopi Lake are proposed in two papers published by Douglass (Douglass 2011; Meek and Douglass 2001). These discussions of spillover hypothesis feature Hopi Lake (aka “Lake Bidahochi”) as the primary cause of the erosion that made Grand Canyon. As noted by geologists Jon Spencer and Philip Pearthree, spillover discussions come naturally because geologic features in the Grand Canyon region seem compatible, whereas the alternate hypothesis of headward erosion and stream capture remain difficult to visualize (Spencer and Pearthree 2001).

Dr. Jon Spencer and his coworkers feature catastrophic lake spillover prominently in explanation of the Lower Colorado River (Spencer et al. 2013). They propose that downstream of Grand Canyon was an almost continuous chain of five or six basins with five lakes—Lake Hualapai, Lake Las Vegas, Lake Mohave, Lake Havasu, and Lake Blythe (fig. 18). These Lower Colorado basins with lakes were supposed to have filled behind and spilled over natural bedrock dams. Freshwater lake evidence is noted from tufa layers in the Pliocene Bouse Formation along the descending stairway of basins and lakes that later formed the path for the Lower Colorado River (Spencer et al. 2013).<sup>30</sup> Lake sediment and fossils in the Blythe Basin just north of Yuma, Arizona indicates the abrupt Pliocene entry of the Colorado River (Bright et al. 2016). The entry of the Colorado River near the town of Blythe is marked by the appearance of green claystone. Discovery of multiple spillover events on the Lower Colorado River lead, by association, to consideration of the same process upstream of Grand Canyon (e.g., Bidahochi Basin) (House 2008, House, Pearthree and Perkins 2008).<sup>31</sup>

A final puzzle piece that remains to be integrated within the Grand Canyon spillover story is the ancient marine delta of the Colorado River. Dr. Rebecca Dorsey, geologist at University of Oregon, and her coworkers favor catastrophic spillover, pointing out that the Colorado River sediment appears very abruptly and recently within the marine mudstone of the Imperial Formation (Pliocene) in Southern California (Dorsey et al 2018). Here green claystone also marks the rapid influx of river water.

### A Very Polite Description of the Last Thirty Years

Let’s go back to the year 1988 and remember three important events that prepared the way for spillover discussions during the last 30 years. First, creationists in 1988 were exploring configuration of ancient Colorado Plateau lakes and their possible points of spillover. Austin was working on the “tectonic tilting hypothesis” for the configuration of Kaibab Upwarp at the time of overflow of Hopi Lake. Holroyd surveyed the upper Colorado River drainage basin and located “Cimarron Spillover” and “Lees Ferry Spillover.” Second, the mud deposits of the Bidahochi Formation were being appreciated with the overall basin geometry of the Colorado Plateau as evidence of a very big lake east of Grand Canyon. Bob Scarborough in 1988 had composed his manuscript “Cenozoic Erosion and Sedimentation in Arizona” that was ready for timely publication in 1989 (Scarborough 1989). Third, also in 1988, Afton Canyon on the Mojave River in Southern California was “rediscovered” as a landscape model for spillover. Meek was ready to publish his 1989 paper detailing the breaching process and the implication that it has wide application to landscapes.<sup>32</sup> In 1988 the spillover hypothesis was beginning as a small trickle of water but was soon to become a torrent!


How has thinking about erosion of Grand Canyon progressed during the last 30 years, since that lecture on the Canyon rim in 1988, since better appreciation of Arizona’s lake mud deposits in 1988, and since the rediscovery of Afton Canyon spillway in 1988? Grand Canyon ranger and geologist Wayne Ranney writes, “So the twentieth century closed without a widely accepted theory on Grand Canyon’s origin” (Ranney 2012, 97). That’s Ranney’s *very polite* way

<sup>28</sup> History Channel documentary “How the Earth Was Made—Grand Canyon” (2009, season 2, episode 1) was posted on YouTube for four years. A+E Networks asked that the clip be removed from YouTube on copyright grounds. Dr. John Douglass and spillover theory appears between time codes 21:00 and 26:22. This is a second stream table experiment. Lake Bidahochi (aka Hopi Lake) is reconstructed at ~7,000 ft elevation, has area of 20,000 square miles, and volume of 3,000 cubic miles making it bigger than Lake Michigan. Green clays of the upper Bidahochi Formation are understood to be evidence of low oxidation in this deep lake. For technical details of the stream table experiment see Douglass et al. (2020).

<sup>30</sup> The term “tufa” rather than “travertine” is used to describe Bouse Formation lake deposits. We agree with these authors on the definition of tufa and its distinction from travertine.

<sup>31</sup> Slide 57 of 66 in House (2008) explains breaching of the Kaibab Plateau.

<sup>32</sup> Meek (1989) is the publication where Afton Canyon is “rediscovered.” Meek (2019) was very well developed by 1992 (first manuscript submission date) of Meek (2019). Why did Meek’s manuscript remain unpublished for 27 years? Could it be the implications for Grand Canyon that others found unsettling?



**Fig. 18.** Lake system downstream of Grand Canyon indicated by tufa deposits of the Bouse Formation according to the fill-and-spill hypothesis of Jon Spencer and his coworkers. Also shown are the marine delta deposits of the Imperial Formation associated with abrupt appearance of the Lower Colorado River according to Rebecca Dorsey and her coworkers. Basins with paleodivides were likely breached from north to south by progressive filling and spill of lakes that established the channel of the Lower Colorado River. Therefore, the lakes did not all appear at the same time. Map is a modified version published by Jon Spencer (*Arizona Geology Magazine*, December 2011).

of saying the ancestral Colorado River did not carve Grand Canyon over tens of millions of years (the former century's National Park consensus ranger explanation). Ranney is also affirming, very politely, that a replacement theory among geologists has not been generally accepted. In less complementary words, one could say that chaos prevails in geologists' thinking about erosion of Grand Canyon! As the

twenty-first century began, Ranney adds, "... spillover theory was nudged to the forefront of ideas regarding the origin of the Colorado River" (Ranney 2012, 103).

### Progress in Creationist Modeling

So, these "fill and spill" ideas have proven beneficial and timely for geology. During the last thirty years creationists have continued to develop the idea of a

breached dam at the eastern Grand Canyon. Austin published in 1994 a map of lakes and description of the Bidahochi Formation with a defense of the breached dam hypothesis for erosion of Grand Canyon (Austin 1994). The global flood made strata of Grand Canyon, the retreat of flood waters beveled the plateau surface, and the structural dam breached in the post-Flood period. Austin favored the initial breach of Kaibab Upwarp at the northwestern margin of Hopi Lake. Also, Austin (2009) published a description in conventional geology literature of the catastrophically eroded landscape on the north slope of Mount St. Helens with a geologic map showing the breached dam and “Little Grand Canyon.” That’s where he described major side canyons associated with the main breach. The comparison of Mount St. Helens to Grand Canyon continues to be significant (see fig. 12).<sup>33</sup> Austin carried on study of breached dams, especially the Santa Cruz River breached dam and giant drained lake in southern Argentina (Austin and Strelin 2011).<sup>34</sup> That Argentina breached dam is just upstream of “Camp Darwin” where young Charles Darwin wrote in April 1834 his journal explaining why he was adopting Charles Lyell’s ideas of slow river canyon erosion.

Dr. Holroyd after 1988 went on to publish his thinking about the ancient big lakes on the Colorado Plateau and how they affected landforms. Many cliffs on the Colorado Plateau studied by Holroyd failed to confirm uniformitarian ideas about boulder aging leading to the term “missing talus” (Holroyd 1987; 1990b). Holroyd states: “Many of us have indelible

images in our minds of the red mudstone Moenkopi Formation slopes at Lees Ferry with missing talus. Many of us believe Lees Ferry is an ancient spillway for catastrophic drainage of a big lake.” Slope analysis of the Colorado Plateau performed by Holroyd (1994) seems to locate possible shorelines that could have been steepened by wave erosion or spring sapping at the edge of lakes. Although computer technology has much improved in the last 30 years, Holroyd’s work shows very early sophistication.

In 1989 Dr. Walter Brown further developed the breached dam hypothesis proposing that a big lake in Utah (he called “Grand Lake”) was the essential trigger agent in eroding Grand Canyon (Brown 2008).<sup>35</sup> Brown’s view is that Grand Lake first overtopped its southern barrier (Lees Ferry at Marble Canyon) at elevation of 5,700 ft. Brown supposed that the big Utah lake drained southward into Hopi Lake causing uplift on the Kaibab Upwarp and breaching of Hopi Lake through southern Kaibab Upwarp. In following years creationists reviewed various breach dam proposals and their evidences (Oard 1993; Williams, Meyers and Wolfrom 1992).

After two years of discussion, Austin’s “bulletin-board map” was published in 1989 (Austin 1989).<sup>36</sup> Also published later in 1989 was Walter Brown’s first Colorado Plateau lake map (Brown 1989).<sup>37</sup> These maps have initiated much discussion on the configuration of lakes and how the breaching of the topographic barriers occurred on Colorado Plateau. When telling a complex story about scientific discovery, one hopes it could be told without a priority dispute.<sup>38</sup> Sadly, “Grand Lake priority

<sup>33</sup> The spillway comparison in fig. 12 draws attention to the very odd comment in a recent publication, “The numerous, U-shaped, small canyons and gullies cutting into the loose material below Mount St. Helens do not look anything like the single, massive, V-shaped Grand Canyon” (Helble and Hill 2016, 170).

<sup>34</sup> A related video posting by Austin is called “Where Darwin Went Wrong” at: <https://www.youtube.com/watch?v=3darzVqzV2o>.

<sup>35</sup> Walter Brown’s breached dam hypothesis appeared as public oral presentation on KTIS Radio, Minneapolis, September 16, 1988. Brown’s breached dam hypothesis first appeared in print in July 1989 in the fifth edition of *In the Beginning: Compelling Evidence for Creation and the Flood*.

<sup>36</sup> Since March 1987, Austin was saying, even in public, “a computer was used to plot the big lake or series of lakes that could form on the Colorado Plateau.” Holroyd was less specific about the computer saying “one could imagine a series of lakes” (Holroyd 1987, 16) even though his Bureau of Reclamation computer had already plotted it, and even though his publication contained some details on the configuration of the “series of lakes.” It was obvious that Holroyd’s “one can imagine” statement was very well informed. Austin mentioned the computer again on the Canyon rim on April 10, 1988. He also described how Hopi Lake could form at 6,000 ft elevation on the east side of the Kaibab Upwarp. Now, talk about the “big series of lakes” and “blocked at the 5,700 ft-elevation” changed in the spring of 1988. Holroyd had made his home-office computer render DEM and imaging software, now making new home-office plots clearly in the public domain. So, Austin requested to publish the “blocked at 5,700 ft-elevation” map (letter to Holroyd, January 23, 1989) and Holroyd (letter to Austin, February 2, 1989) gave permission. Finally, after being posted on the bulletin board for two years, the “blocked at 5,700 ft-elevation” map was distributed by publication in March 1989.

<sup>37</sup> Brown (1989, 75) introduces “. . . what we call Grand Lake. . . . It stood at elevation of 5,700 feet above present sea level.” Eight pages later on page 83 is Brown’s map with no mention there of 5,700 ft elevation. Details of the Grand Lake breach process are offered on page 83: “The catastrophic dumping of Grand Lake took place through what is now the gap between Echo Cliffs and Vermillion Cliffs.”

<sup>38</sup> Scientific priority is recognition given to the individual or group of scientists that first make a discovery or propose a theory. Personal rewards like fame, prizes, or authority can accrue to the scientist who is first to publish a new finding even if several other researchers came to the same conclusion independently and at the same time. The priority game makes science publication a “winner take all” event. There are no second-place prizes, even if the new discovery is drawn from shoddy data. Harvard evolutionist Stephen Jay Gould commented that “debates about priority of ideas are usually among the most misdirected in the history of science” (Gould 1985, 35). Christians ought to exemplify the life of Christ and eschew prideful, self-centered, and narcissistic conduct seeking recognition and authority. That’s the Pharisee’s behavior that Christ denounced. If you must research the destructive effects of priority disputes upon science, you might begin by researching why Albert Einstein won the Nobel Prize in 1921 for “photoelectric effect” rather than for “general relativity.” Obviously, general relativity is Einstein’s chief contribution to science! Who remembers his photoelectric publications? Sadly, during Einstein’s career, relativity was embroiled in a priority dispute. Wikipedia Foundation had to deal with the ugliness of the Einstein priority debacle when the Einstein biography was written. The Wikipedia biography of Einstein ignores the priority dispute completely. One can find a hint of a problem at the “See Also” section at the end of the biography. Wikipedia gives a link to “relativity priority dispute.”

dispute<sup>39</sup> focused on the big lake in Utah, specifically its elevation<sup>40</sup> and lake position, spillover point and spillway configuration.<sup>41</sup> Austin calls it Canyonlands Lake (fig. 4) and maintains the dispute misdirects discussions of Grand Canyon erosion, because Hopi Lake (aka Lake Bidahochi) is most likely the primary cause of breaching Kaibab Upwarp.

Principal spillover advocates (Scarborough, Austin, Meek, Douglass, Spencer, and House) feature Hopi Lake as the primary agent for initial Grand Canyon erosion. Austin, like the other spillover advocates, does not invoke a separate lake in Utah as being a contributing cause to the initial breaching of Kaibab Upwarp.<sup>42</sup>

<sup>39</sup> Priority disputes are generally regarded as worthless and insipid by scientists. What if someone meets with you and the conversation turns toward the ancient “big Utah lake priority dispute?” It would be rude to walk away. You could respond with a caution about the overall misdirected and noxious history of priority disputes in science. Their track record is abysmal! You could mention Apostle Paul’s injunction, “Don’t have anything to do with foolish and stupid arguments, because you know they produce quarrels” (2 Timothy 2:23, NIV). Make sure you ask them why the priority of the ancient big Utah Lake, is worth discussing. Let’s say they entice you with pleasant and reasonable speech. If they convince you that the priority dispute is more than a banal enterprise, you must insist that the discussion be narrowly focused to the two 1989 maps and the authorship events they imply happened before the end of 1989. Further focus discussion by naming it “1989 big Utah lake map priority dispute.” You could ask four “when” questions about the “1989 big Utah lake map priority dispute.” The four “when” questions need an answer with specific date of 1989 or earlier because of the evidence in the two maps. (1) When was the elevation 5,700ft first used in reference to an ancient lake in Utah? (2) When was the location of the southern boundary of the Utah lake first described? (3) When was the spillover point at Lees Ferry first described? (4) When was a catastrophic understanding of the Marble Canyon flood spillway first proposed? Avoid discussions of hearsay, copyright infringement, plagiarism, fraud, and anything like tabloid journalism. Don’t be enticed! Set severe ground rules. Remember Apostle Paul’s concluding warning to Timothy, “Turn away from godless chatter and the opposing ideas of what is falsely called knowledge” (1 Timothy 6:20 NIV). If the guidelines you require fail to maintain their focus or are violated, just *walk out*.

<sup>40</sup> Austin’s “blocked at 5,700ft elevation” map (fig. 6) was published in February/March 1989. Four months later in July 1989, Walter Brown (1989, 75 and 83) published a different Colorado Plateau lake map with reference to Grand Lake “which stood at an elevation of 5,700 feet.” It would appear by date of publication that Austin has priority on the lake at 5,700ft elevation. However, since July 1989, people have asked how the two maps were composed, both maps with reference to elevation of lakes at 5,700ft. Both Austin and Holroyd believe they together composed the words “blocked at the 5,700foot elevation” in a phone conversation in March 1987. Austin believes he asked David McQueen to review his map with caption “blocked at the 5,700foot elevation” in March 1987. Austin, Holroyd, and McQueen believe that the two maps were composed independently with the 5,700foot elevation just being coincidental. Holroyd showed how to draw the big lake in his 1987 paper. Anyone could have drawn it by Holroyd’s method using topographic maps and a pencil. A different explanation of the map similarity is offered by Brown (2019, posted on Internet at [www.creationscience.com](http://www.creationscience.com), updated January 9, 2019, with more documents offered through snail mail on CD-ROM). Brown supposed Austin heard Brown’s public radio broadcast on KTIS Radio, September 16, 1988 where Brown first discussed in public a Colorado Plateau lake in Utah at 5,700ft elevation. Brown thinks the elevation 5,700ft is his “trademark” showing his pedigree on that lake elevation. Brown claims Austin repositioned pages deceptively in his 1989 *Guidebook* to make it appear he had priority. Thus, Brown is adding “fraud” to Austin’s account as Brown asserts his priority claim. However, Austin’s bulletin-board map, posted in March 1987, showed the hypothetical series of lakes upstream of Grand Canyon with the words “blocked at the 5,700ft elevation” 18 months before Brown’s radio broadcast. Austin’s map was drafted in March 1987 with Holroyd’s assistance and approval, and with McQueen’s review. That map remained unpublished on the bulletin board for two years until after Holroyd’s workplace computer problem at Bureau of Reclamation was resolved in spring 1988. Austin claims he knew nothing at that time about Brown’s radio broadcast. Austin, Holroyd, and McQueen believe the map elevation 5,700ft is just a coincidence. Another possible explanation is that a frequent visitor to ICR Graduate School, somebody like Paul MacKinney, could have informed Dr. Brown of details of the posted bulletin-board map.

<sup>41</sup> The priority dispute was not just the big Utah lake’s elevation. Walt Brown also maintains he has priority on the Utah lake’s location, breach point and characteristic spillway erosion. This priority issue appeared as a statement with low visibility in an endnote within chapter text in Brown (2008, 189 with endnote on pages 213–215) but was promoted to chapter text as a noticeable sidebar called “A Controversy” in the ninth edition of his book (Brown 2019, online ninth edition available at [www.creationscience.com](http://www.creationscience.com) updated 1/9/2019 with documents offered on CD-ROM). First, Brown read Austin’s 1988 *Guidebook* in April 1988 (letter from Brown, July 6, 1993). Second, he conducted field research near Marble Canyon in summer 1988. Finally, he began speaking publicly. Brown (2019) says in the sidebar section of ninth edition, “Beginning, in my fall 1988 seminars and on a 16 September 1988 radio broadcast over 200 stations, I described Grand Lake’s key features, location and how its breaching formed the Grand Canyon. This explanation was first published in July 1989” (quoting from sidebar in Brown 2019, posted on Internet, updated 1/9/2019). Then, Brown focuses the sidebar on Brown’s priority stating, “Austin probably knew the serious problems (mentioned above) that faced any proposal that the Grand Canyon was carved by the breaching of Hopi Lake. What he did not realize, as his writing exposes, was that a much larger and separate post-flood lake, Grand Lake, was north of Hopi Lake. Austin was unable to produce any spoken or written record showing that he knew, before 1989, anything about Grand Lake” (quoting from sidebar in Brown 2019, posted on Internet, updated 1/9/2019).

Let’s take a closer look at Brown’s priority argument for the big Utah lake. The argument is rather awkward because Brown is establishing what Austin did not think and what Austin did not do in 1988. Brown says he knows Austin’s thinking and actions in 1988 well enough to establish what did not happen! Certainly, according to Brown, Austin never thought in 1988 about a big lake in southern Utah breaching at Lees Ferry. Also, most certainly, according to Brown, Austin never communicated in 1988 with another person about failure of a big lake and dam in Utah. In other words, Brown had universal surveillance of Austin’s thoughts and actions in 1988. Brown is establishing his priority by a “God argument.”

Remember that Ed Holroyd conducted field work in Marble Canyon and at Lees Ferry in July 1987. He reported that work to Steve Austin in a phone call in December 1987. That phone call was of so much interest to Austin that he requested Holroyd to write it down as a continuing research proposal that a geology graduate student could use. Holroyd’s three-page research proposal was titled, “Some Research Opportunities at Marble Canyon for Creationists (written January 1988 from observations in July 1987)” with cover letter to Austin and Holroyd’s signature dated January 24, 1988. Holroyd described a big Utah lake north of Lees Ferry, a breach point at Lees Ferry, and a flood spillway within Marble Canyon. Walt Brown’s “God argument” about what Austin “did not realize” is incorrect. Austin was well informed in 1987 about the “big Utah lake,” its breach point at Lees Ferry, and its spillway at Marble Canyon. He learned about the “crack-and-capture hypothesis” directly from its author Ed Holroyd. Thus, Ed Holroyd appears to have priority over Walt Brown in describing the lake breach point at Lees Ferry, the lake location with southern boundary at Echo Cliffs Monocline, and spillway configuration at Marble Canyon. Both Austin and Holroyd have priority in March 1987 on the lake elevation at 5,700ft. Holroyd had a workable version of the “big Utah lake” (his version of Brown’s “Grand Lake Explanation”) that was well formulated in writing by January 1988.

<sup>42</sup> General spillover theory usually supposes a breach from a higher elevation basin into an adjacent lower elevation basin. Walter Brown’s specific “Grand Lake Explanation” is opposite of classic spillover theory. Brown abandons conventional spillover by having Grand Lake breaching from 5,700ft elevation into Hopi Lake at over 5,950ft elevation.



The early statement of spillover hypothesis by Austin (1994) and Holroyd (1994) has been improved through the years. Austin and Holroyd currently favor the popular “top-down spillover” scenario (outlined by Douglass et al. 2020; Hilgendorf et al. 2020; Meek 2019). That scenario postulates that highest elevation basins breach progressively into lower elevation basins leaving upstream a river within its canyon. Austin and Holroyd continue to favor Austin’s 1987 “plateau tilting model” for Grand Canyon plateaus, and the notion has been borne out by recent tectonic study (Crow et al. 2014). According to Austin, those plateaus on the north side of the Colorado River have tilted down on the north side and up on the south side since drainage of the lakes. We can suppose more than 1,000 ft (300 m) elevation change on Kaibab Plateau, Kanab Plateau, and Marble Platform (with its included Paria Plateau subunit) since drainage of the lakes. Understanding of the tilted plateaus allows the configuration of those lakes to be approximated as shown in fig 4.

Austin and Holroyd believe that Canyonlands Lake failed first. Fig 4 suggests its elevation was 6,400ft contained behind the formerly higher Echo Cliffs Monocline. We believe it spilled at Lees Ferry Spillover #1 (location in fig 4) into Hopi Lake basin that was largely vacant of a lake. The wide portion of Marble Canyon Spillway was excavated by flood flow from Canyonlands Lake. The addition of over 300ft of new water to Hopi basin deposited the upper member of the Bidahochi Formation, likely disturbed magma beneath the lake’s floor, and created instability of the dam on the west side of Hopi basin. Kaibab Upwarp was the western barrier of Hopi Lake that failed at East Kaibab Spillover #2. Austin believes the westward breach and sweeping turn-to-the-right of the Colorado River first occurred in the Moenkopi Formation on the north side of the turning erosional scarp of the very durable Shinarump Conglomerate as it originally covered the southern slope of the Kaibab Upwarp.<sup>43</sup> As Hopi Lake began to drain westward, eastern Grand Canyon within the Kaibab Upwarp first started to be carved. Soon, as partial drainage of Hopi Lake occurred, Blue Moon Bench (fig 4) was exposed as two lakes appeared—a smaller temporary lake on the northwest and a much bigger temporary lake on the southeast. The smaller temporary lake northwest of Blue Moon Bench drained southward from Nankoweap Spillover #3 into the enlarging eastern canyon. Spillover #3 formed the knickpoint that retreated northward incising the narrower inner channel of Marble Canyon. The larger temporary

lake on the south side of Blue Moon Bench drained westward at Little Colorado Spillover #4 with its knickpoint retreating eastward incising the inner channel of the Little Colorado River. The configuration of these spillovers with knickpoint retreats was successfully modeled in the remarkable stream table experiment described by Douglass et al. 2020.

Westward drainage of Hopi Lake over Kaibab Plateau allowed two topographic depressions downstream to fill completely merging into a single, larger Lake Toroweap (fig 4). Ponding on the western side of Coconino Plateau was separated by an arch from ponding on Kanab Plateau. To the west was Uinkaret Plateau with Toroweap Fault forming the topographic dam on Lake Toroweap’s west side. As water rose to elevation above 5,800ft, the single larger lake overtopped Uinkaret Plateau at the western side of Lake Toroweap. Overtopping occurred at Toroweap Spillover #5 (fig 4) forming the narrowest spillway constriction within Grand Canyon, just 4 miles wide at the Toroweap narrows. Toroweap dam and its new spillway were incised to the 4,000-foot elevation exposing the Esplanade Sandstone widely over the lake’s floor. The knickpoint within the spillway eroded east-northeastward along the broad structural arch between the two basins. Two structural synclines controlled the location of two other spillover points as the lake drained. As the level of the water lowered, two temporary lakes formed north and south of the new spillway. On the southside of the spillway Cataract Syncline (Billingsley 2000) formed a low area where Havasu Spillover #6 (fig 4) began incising Havasu Creek’s canyon along the axis of the syncline. The separate temporary lake on the north side had a low point along the Kanab Syncline (Billingsley 2000) forming Kanab Spillover #7 (fig 4) that incised Kanab Creek canyon. The two knickpoints eroded down to the Esplanade level and retreated upstream from “spillover points” through synclines to form very wide side canyons. Later, channelized flow narrowly focused erosion as canyons deepened. “Wine-glass canyons” (very wide above Esplanade and very narrow below Esplanade) are the most distinctive landform within the floor of former Lake Toroweap.

By 2003 the breached dam hypothesis had become the most popular creationist explanation for the erosion of Grand Canyon. Tom Vail’s book *Grand Canyon: A Different View* was for sale at Grand Canyon National Park in the rim bookstores. The introduction in Vail’s book contains the words of the father of the modern creationist movement Dr. Henry M. Morris, Jr., “...a

<sup>43</sup> This explanation of erosional scarp breaching of Kaibab Plateau follows Babenroth and Strahler (1945) and Douglass et al. (2020). The plateau tilting hypothesis of Austin suggests Hopi Lake overflowed at 6,100ft elevation (1860m) as proposed by Scarborough (2001). Douglass et al. (2020) do not invoke plateau tilting and suppose Lake Bidahochi breached Kaibab Plateau at about 7,400ft elevation (~2,260m). Obviously, these ideas should direct further future research.

great dammed-up lake full of water from the Flood suddenly broke and a mighty hydraulic monster roared down toward the sea, digging deeply into the path it had chosen...” (Vail 2003, 4). Tom Vail and Dr. Andrew Snelling, two veterans of Grand Canyon geology, featured the breached dam hypothesis (Snelling and Vail 2009). In 2007 Austin’s version of the breached dam hypothesis was displayed prominently in the Answers in Genesis Creation Museum in Petersburg, Kentucky (fig. 19). One recent review article summarizes creationist thinking about Grand Canyon erosion, “During the post-Flood years, water levels eventually grew and breached the natural dams to carve the canyon, a monument to catastrophe” (Mitchell 2013).<sup>44</sup>

Some creationists, however, do not believe that the post-Flood lake existed in northeastern Arizona. They suppose that the Bidahochi Formation is somehow a Flood deposit, so they dismiss spillover. These creationists believe that Grand Canyon was eroded by the retreating Flood. Peter Scheele understands that a wide channel (“river”) of retreating Flood water drained westward through northern Arizona eroding Grand Canyon from west to east through soft Flood

sediment (Scheele 2010). Meteorologist Mike Oard sees the retreat of the Flood differently. According to Oard (2016), the Grand Canyon strata were lithified late in the Flood while the Flood water still sat over northern Arizona. Then, Oard proposes, high velocity currents merged together flowing westward in deep ocean water late in the Flood. As ocean currents assembled, they eroded a submarine canyon system from east to west through the solid rock of the submerged Colorado Plateau. After the high velocity currents eroded the submarine canyon system, the retreat of the Flood exposed the carved rock surface of the Plateau which was ready made, sculpted with intricate detail through the Kaibab Upwarp for the modern Colorado River.<sup>45</sup> In Mike Oard’s view, Grand Canyon is a superposed submarine canyon! Oard’s hypothesis is completely new to geologists, never before proposed in 150 years of Grand Canyon discussion.

So, the initial focus of the Grand Canyon erosion discussion among creationists needs to be the Hopi Lake interpretation and better understanding of the sedimentology and paleontology of Bidahochi Formation. Is that



**Fig. 19.** The breached dam hypothesis became the most popular creationist explanation for the erosion of Grand Canyon as seen in this 2007 mural in the Answers in Genesis Creation Museum in Petersburg, Kentucky.

<sup>44</sup> This Answers in Genesis news release is a response to Oskin 2012 (Yahoo News) review of the technical paper by Dickinson 2013.

<sup>45</sup> Oard’s explanation, especially late-Flood submarine lithification with late-Flood submarine canyon erosion, prompted geologist William Hoesch to communicate to Austin and ask, “Is there anything that Noah’s Flood *cannot* do?”

evidence of the post-Flood lake? Or, is Bidahochi Formation deposited by the ocean flood upstream as the submarine canyon is being sculpted by deep marine currents toward the west?


### Does Hopi Lake Hold Water?

The most extensive part of the Bidahochi Formation is what has been called informally the “upper member” (see fig. 20). The prominent outcrop begins over 100 mi southeast of Grand Canyon. That prominent outcrop is 125 mi long on the east side of the Little Colorado River north of Holbrook, Arizona and extends eastward to just south of Gallup, New Mexico. It is often identified as “Pliocene” in geologists’ nomenclature, and usually assumed, somewhat tentatively, to be “about six million years old.” The uppermost Bidahochi averages 200 ft thick and is composed of light green and brown claystone, volcanic ash beds, and clayey sandstone, light brown cross-bedded sandstone and gray limestone. Sandstone strata of the “upper member” contain abundant calcite cement, as is common in lake sand deposits.

Tufa, a type of limestone, is the most diagnostic lacustrine sediment of the Bidahochi Formation. Tufa is the porous and laminated calcium carbonate lake deposit and is especially characteristic within eastern areas of uppermost Bidahochi outcrop. There is clear evidence that this tufa is not pedogenic (not a soil-alteration calcrete). Near the town of St Johns, tufa caps many terraces, mesas, and buttes at elevations of 6,000 to 6,700 ft near the Little Colorado River (Harris et al. 1998). Bidahochi limestone is also often seen around volcanic structures called maars in the Hopi Buttes area. Among the most famous freshwater

fish fossil localities is Coliseum Maar (fig. 21) where limestone intertongues with volcanic rocks (Uyeno and Miller 1965).<sup>46</sup> The strontium isotope ratio of upper Bidahochi limestone is not seawater’s ratio, but the ratio of nearby surface streams, again suggesting lake deposition (Douglass et al. 2020; Gross et al. 2001). The high strontium ratio of Bouse Formation limestone (generally acknowledged to be ponded lake tufa) resembles the high ratio of the upper Bidahochi limestone (Douglass et al. 2020). Outside of the area of primary Bidahochi outcrops, limestone resembling the main formation is known. Most interesting are thin outcrops of gray limestone around Preston Mesa at 6,000 ft elevation just north of Tuba City.

Volcanic structures are unexpectedly abundant within Bidahochi lake deposits, especially at Hopi Buttes volcanic field. There 300 *maar craters* have been identified (fig. 21). Magma near the bed of the lake exploded, then the crater in the lakebed collapsed creating steeply inclined strata within the crater. Also, the magma explosion at the lake floor in Coliseum Maar created a volcanic surge that buried freshwater fish in a 3 ft thick mass-kill limestone layer. The maar crater rim has a surrounding “ejecta ring” typically composed of volcanic ash of limited extent. Another, less abundant lake-floor volcanic structure is the *scoria cone* where dark colored basaltic cinders accumulated around a central neck or plug (fig. 21, central right). Sedimentary strata (especially limestone and tufa) with volcanic structures (maar craters and scoria cones) strongly support deposition of Bidahochi Formation in standing water (i.e., lake), not flowing water (i.e., Flood) (White 1990; Zelawski 2011).<sup>47</sup>



**Fig. 20.** Bidahochi Formation lake beds at Roberts Mesa about 15 mi north of Indian Wells, Arizona. Upper Bidahochi Formation is over 200 ft thick composed of green, tan, and reddish-brown claystone with tan siltstone and very fine sandstone forming the calcite-cemented cliff. Lower Bidahochi member is red mudstone on the left marking the base of the slope at 6,200 ft elevation. Photo by Brian Gootee, Arizona Geological Survey, March 2018.

<sup>46</sup> Four species of cyprinid fish (minnows) are described, including fish from the 3 ft thick, mass-kill limestone bed at Coliseum Maar.

<sup>47</sup> White (1990) says: “The temporal and geometric relationships among lithofacies of the Bidahochi Formation illustrate the dynamic interplay between volcanism and sedimentation in an essentially ‘closed’ sedimentary system.” Maar craters with surrounding ejecta rings, as well as scoria cones at central plugs, argue for stationary and closed lake depositional environment. Zelawski (2011) gives an extremely detailed facies analysis of marginal volcanic deposits.



**Fig. 21.** Oblique aerial image displays two typical lake-floor volcanic structures of the Bidahochi Formation in the Hopi Buttes area, west of Indian Wells, Arizona. The two lake-floor structures are the maar crater with ejecta ring and the scoria cone around a central vent. Front center is “Coliseum Maar,” a one-half-mile diameter lake-floor volcanic explosion crater. Magma near the bed of the lake exploded, then the crater in the lakebed collapsed, creating a volcanic surge that buried freshwater fish in a 3 ft thick limestone mass-kill layer. Latest lake-floor strata within the crater are inclined with inclination often more than 30° toward the center of the crater (making the interior of the crater appear like the Roman Coliseum). Around the crater rim is the flat-lying ejecta ring deposit of tuff. Just in back of the maar rim, slightly to the right side, is the scoria cone, a dark-colored hill of unsorted basaltic scoria and palagonitic tuff surrounding a central neck or plug of intrusive igneous rock. Ejecta rings surrounding maars are still associated with their eruption crater rims and mass-kill fish layers at present elevation of 6,000 ft above sea level. Scoria cones, although deeply weathered and eroded, are still adjacent to their eruptive centers (the intrusive plug or point of eruption of lava flows). Sedimentary strata with contained volcanic structures strongly support deposition of upper Bidahochi Formation in standing water (i.e., lake), not flowing water (i.e., Flood). Three hundred maar structures with their associated ejecta rings within lake beds are distributed through Hopi Buttes volcanic field in Arizona. Photo copyright by John Wark, used by permission.


If upper Bidahochi Formation was deposited at the end of the biblical Flood while a westward surging ocean current excavated Grand Canyon, ejecta rings would be eliminated from Bidahochi maar rims. Unidirectional flood flow must modify the crater rim and the sediment within the crater. Also, Flood current would redistribute scoria away from the eruptive pipe or plug destroying a scoria cone completely. Therefore, the “upper member” of Bidahochi Formation is best understood to represent the last deposits within the ancient, post-Flood, freshwater lake.

Freshwater fossils are abundant in the upper Bidahochi Formation, an observation which also supports the lake sedimentation view. Lake and

pond-dwelling snail fossils occur without land-dwelling snails (Taylor 1957).<sup>48</sup> The genus *Lymnaea*, the prominent larger pond snail in most fossil collections from upper Bidahochi, is today a versatile *freshwater* snail marketed to hobbyists for their home aquariums. Prominent fish fossils (Uyeno and Miller 1965) are minnows and chub strikingly similar to the same fast-swimming fish in modern freshwater lakes and rivers of the Colorado Plateau (fig. 22). The adult Colorado River Pikeminnow can attain huge size (up to 6ft) (Smith, Reynolds, and Stewart 2013)<sup>49</sup> and was a staple in the diet of Native Americans and early Utah settlers, but Pikeminnow size and abundance has diminished in recent times. Upper Bidahochi amphibian fossils (various frogs and toads)

<sup>48</sup> “The absence of land snails is somewhat puzzling. Possibly they were missed in collecting, but this seems unlikely” (Taylor 1957, 656).

<sup>49</sup> Colorado River Pikeminnow (*Ptychocheilus lucius*) attains a length of 1.5 m and indicates proximity to large pools and a big, swift river. The authors say that the Miocene fishes of the Lower Colorado River had been isolated from the upper Colorado basin until the Pliocene Colorado River spillover through Grand Canyon.



**Fig. 22.** Fossil fish from the upper Bidahochi Formation. These are juvenile Colorado River Pikeminnow that indicate a freshwater lake associated with swift river in northeastern Arizona. Photos from Robert L. Sutton of U.S. Geological Survey, reproduced at same scale.

(Parmley and Peck 2002)<sup>50</sup> resemble those today in ponds along the modern Colorado River in Grand Canyon. Fossil footprint trackways (Breed 1973) in upper Bidahochi include a wading bird (resembling heron) and a large web-footed bird (resembling goose), birds which today feed on the same kinds of lake snails, frogs and minnows in Grand Canyon. Beaver fossils (Stirton 1936) have even been reported from the upper Bidahochi Formation. Beavers live today within Grand Canyon. Dave McQueen comments, “The assemblage of fossils from the upper member Bidahochi appears to be the remains of the last complete lake ecosystem that is now gone. That’s like the post-Flood, freshwater bayou ecosystem near my home in Louisiana.”

Further support for the idea that Hopi Lake was a big lake came from an unlikely source. It was the fish fossils. Jon Spencer and his coauthors (Spencer, Pearthree, and House 2008) described the fossil Pliocene fishes of the upper member of the Bidahochi Formation and noted their dissimilarity to fossil fishes downstream in Los Angeles Basin in

Southern California. Instead, these authors noted that Bidahochi fishes possess striking similarity to fossil fishes in the Pliocene of the Snake River in Idaho. Spencer and his coauthors argued that the two Pliocene populations of fishes must be connected genetically when the upper Snake River was tributary to an enormous “closed drainage basin” on the Colorado Plateau that included a swiftly flowing river connected to Hopi Lake. Again, paleontology supports the lake. That explanation was reconfirmed and further developed by Douglass et al. (2020).

### Shoreline Deposits Next to Spillover Point

The upper member of Bidahochi Formation occurs prominently on the east side of the topographic basin north and east of Holbrook, Arizona, along the Little Colorado River. There a type of limestone called tufa is abundant. The uppermost Bidahochi contains noteworthy fossils representing a post-Flood freshwater ecosystem with associated green claystone layers indicating a large, and likely deep, lake. Could these lake deposits also occur on the *west side* of the topographic basin adjacent to the Kaibab Upwarp? Studies of the geometry of Bidahochi Basin reveal no structural closure on its west side as it extends westward across the Little Colorado River to Kaibab Upwarp and Mogollon Rim (Dallegge, Ort, and McIntosh 2003; Holm 2001). These discoveries seem to allow Hopi Lake to exist at high elevation west of the Little Colorado River in Pliocene time.

Most importantly, we have the observation of what appears to be lake deposits of the west side. Bob Scarborough and Richard Hereford studied a five-foot-thick freshwater limestone near Cape Solitude (location in fig. 4) just on the eastern side of Grand Canyon.<sup>51</sup> According to Scarborough (Scarborough 2001), the limestone contains small white domal algal structures that he calls “tufa,” associating it with the western shoreline of Hopi Lake. It seems unlikely that the Cape Solitude tufa is a carbonate-cemented soil (caliche or calcrete), and unlikely to be a local spring or cave deposit (travertine). Cape Solitude tufa is porous, algal laminated and forms a surface-encrusting deposit. It closely resembles in structure and composition Lake Bonneville “capping tufa” which is of undoubted lake origin (Felton et al. 2006).<sup>52</sup> Scarborough and Austin also discovered freshwater limestone and claystone on the north side

<sup>50</sup> “All of the taxa identified to living species occur in or near the White Cone area today” (Parmley and Peck 2002, 177). Four genera of frogs and toads (*Rana*, *Hyla*, *Bufo*, and *Scaphiopus*) have been identified in Upper Bidahochi Formation at White Cone. Species of each of these four genera have been identified living in Grand Canyon (northern leopard frog, canyon tree frog, red spotted toad, and spadefooted toad). Hikers are familiar with the distinctive croaking of *Bufo* (red spotted toad) next to Grand Canyon side streams.

<sup>51</sup> Cape Solitude tufa studied by Scarborough is located at latitude 36.169N, and longitude 111.798W. Elevation is 6,130 ft.

<sup>52</sup> The authors describe “Capping tufa” from Lake Bonneville shorelines as a level-surface deposit typically covering “beachrock.” Beachrock is a clast-supported calcium carbonate deposit. Cape Solitude tufa has composition, structure and position characteristics closely matching Lake Bonneville capping tufa.

of the Little Colorado River north of Cape Solitude. For good reason Scarborough supposes the western shore of the very large Hopi Lake was up against the Kaibab Upwarp at elevation above 6,000ft. Visible at distance from No Name Point, that outcrop of Cape Solitude tufa (straightforwardly understood to be freshwater limestone) is at elevation 6,130ft. Eastern deposits of upper Bidahochi Formation have an average elevation of 6,300ft (Chen and Maloof 2017).<sup>53</sup> The two college geology professors, Norman Meek and John Douglass add to Scarborough's work by affirming how the incision of the Grand Canyon was initiated by the spill from Hopi Lake across an erosional scarp formed by the very resistant dipping Shinarump Conglomerate (Douglass 2011; Meek and Douglass 2001).<sup>54</sup> That is what was being simulated in the stream table experiments by Douglass shown on made-for-television documentaries.

### The Lake That Gets No Respect

An American standup comedian named Rodney Dangerfield became famous for the line, "I don't get no respect!" In a very humorous way, he called attention to people's dismissive attitude when encountering something or some person they perceive to be odd, unexpected or quirky. We have a tendency to marginalize things that don't meet our expectations. Hopi Lake had 300 maar volcanoes that exploded within its basin spreading ash. Also, several hundred scoria cone volcanoes erupted less explosively (Zelawski 2011). Subaqueous mass flows buried freshwater fish abruptly and testify to very rapid deposition in the lake. Nothing about Hopi Lake commends it to our minds as a "normal" lake. Is it the lake that we must marginalize? Just because we marginalize it, that does not mean it was not a lake. Should familiarity breed contempt? Austin says, "I remember a geology society meeting in the 1980s when Dr. Gene Shoemaker spoke disapprovingly about Hopi Lake. He said that potassium-argon dating of Bidahochi volcanic rocks proved the geological oddity we call Hopi Lake was eight million years old, making it much older than

the erosion of Grand Canyon. Shoemaker said that we should dismiss Hopi Lake overflowing millions of years later to erode Grand Canyon because the lake basin by then was certainly filled with sediment and dried up before the Canyon appeared." Since then, many geologists have persisted in associating Hopi Lake with distain, imagining it to be peculiar or bizarre, even though its existence is appreciated from straightforward geologic evidence. Many have offered a litany of reasons why Hopi Lake should not exist, or why it should exist for only a very short time. It sits on the top of the landscape, and it is impossible to forget. We can often detect disrespect for Hopi Lake by listening to how geologists speak about it.

As discussed previously, post-Flood lakes and spillover erosion have been a favored interpretation among creationists. Leonard Brand and Arthur Chadwick wrote of "...compelling evidence for many large basin-filling lakes in the time following the flood..." (Brand and Chadwick 2016, 385). However, the most severe criticisms of Hopi Lake, sadly, come from creationists. Meteorologist Mike Oard writes of "no evidence for the lakes," "no lake-bottom sediments," and "no shorelines" (Oard 2010; 2016).<sup>55</sup> When Oard writes universal negatives about "no evidence," he seems to be expressing Godlike omniscience. Scientists know it takes just one affirmative example to disprove a universal negative. Austin responds to Oard saying: "There is good evidence for the post-Flood lake sediments in Bidahochi Formation. Tufa deposits are common on terraces, buttes and mesas overlooking the Little Colorado River near St. Johns, Arizona. Lake tufa deposits at Cape Solitude resemble "capping tufa" of undoubted lake origin on flatter surfaces at Lake Bonneville. Cape Solitude tufa is especially relevant for discussion because it is next to the potential spillover point at Kaibab Upwarp. Scientists are supposed to talk about what they see, not about what they don't see. I'm pointing to what I see. Oard is telling us what he hasn't seen, or, even worse, what he refuses to see. Let's do good science, and let's encourage those who are doing it!" Sadly, creation

<sup>53</sup> Ice Age Lake Bonneville shows why we want to be very cautious in using modern elevations uncritically to infer ancient lake shorelines. The weight of ancient glacial Lake Bonneville in Utah depressed the center of the region up to 76m (250ft) according to Chen and Maloof (2017). Now that Lake Bonneville is almost empty, the terrain has risen isostatically to different heights, higher rise in the center of the lake basin. Using DEM for shoreline positions of ancient lakes must be inaccurate. Upper Bidahochi member strata average over 6,300ft elevation today, are near the basin center, and could have risen 300ft or more. Cape Solitude likely rose 150ft. The Kaibab Upwarp on its south side, unloaded by erosion of more than 4,000ft of Grand Canyon strata, likely rose isostatically many hundreds of feet since the dam was breached. How much tectonic rise occurred due to further flexing of the East Kaibab monocline after the breach event? Austin believes that there are also hundreds of feet of vertical uplift on the south side of the East Kaibab monocline. He likes to think that the Kaibab Plateau has been tilted by rotation since the breach (up on the south side and down on the north side) likely caused by regional NNE-directed right-slip on basement faults. There is good evidence of isostatic and tectonic tilting of the Kanab Plateau and the Marble Platform (up on the south and down on the north).

<sup>54</sup> These authors propose Lake Bidahochi (aka Hopi Lake) was at high elevation of ~7,000ft next to Kaibab Upwarp. They do not describe recent isostatic or tectonic change in the spillway configuration.

<sup>55</sup> The editor of the *Creation Research Society Quarterly* should get better peer review, not letting "no evidence" statements concerning Bidahochi Formation to pass without receiving some qualification.

geologist Dr. Tim Clarey appears to agree with Oard (Clarey 2020).<sup>56</sup> Clarey dismisses Hopi Lake which he regards as one of those “Fictional lakes that some creation geologists propose emptied in a catastrophic manner to carve Grand Canyon but are based on little if any geological evidence” (Clarey 2018). McQueen responds to Oard and Clarey: “Hopi Lake should get better respect from these creationists. There is good evidence for the ancient post-Flood lake.”

### The Empire Strikes Back!

U.S. Geological Survey geologist Ivo Lucchitta closely studied what he supposed to be an abandoned ancient river channel gravel deposit just 30 mi east of the Kaibab Upwarp at Crooked Ridge (located in fig. 4). The gravel occurs at The Gap (location in fig. 4) at elevation of 5,600ft within what Lucchitta describes as a channel up to 6 mi wide that rises in elevation northeastward to White Mesa (location in fig. 4) at elevation of 6,700ft and beyond, perhaps a total distance of 55 mi. Lucchitta argued in 2011 and 2013 papers that the deep channel gravel is Miocene making it older than the Bidahochi Formation lake deposits (Lucchitta 2013; Lucchitta, Holm, and Lucchitta 2011). Furthermore, volcanic pebbles indicate the gravel source was the San Juan Mountains in southern Colorado. Putting “Crooked Ridge River” so near to Grand Canyon seemed to argue that the Kaibab Upwarp was already breached in Miocene time by a great river, so the supposed Pliocene Hopi Lake could not later have been confined topographically at high elevation by the Kaibab Upwarp. Lucchitta imagined the big Miocene “Crooked Ridge River Valley” extending westward from The Gap into the Kaibab Upwarp. That evidence for a lower elevation Miocene river channel, if interpreted correctly, would be a compelling argument against Pliocene Hopi Lake at high elevation and at later time extending westward to the Kaibab Upwarp.

William Dickinson, professor of geology at University of Arizona, was critical of Hopi Lake supposing (1) a very small lake of Miocene age on just

the eastern side of the basin, and (2) upper member Bidahochi Formation being river deposited without reference to a lake (Dickinson 2013). Like Lucchitta, Dickinson featured the Crooked Ridge gravel deposit as a deep, lower elevation, Miocene river channel located on the west side of the basin. Dickinson’s 2013 paper was titled “Rejection of the Spillover Model for Initial Incision of the Grand Canyon.” The Yahoo News Internet release (Oskin 2012) for Dickinson’s work is titled, “Grand Canyon Carved by Flood? Geologist Says No.” The news release says, “Tracing the history of the Grand Canyon is controversial... Dickinson hopes at least to lay to rest one hypothesis: That an ancient lake carved the canyon through a cascading series of waterfalls. A favored concept for two decades...” Then, University of New Mexico geologist Karl Karlstrom *with ten coauthors* published in the prestigious journal *Nature Geoscience* that Crooked Ridge paleoriver flowed westward from San Juan Mountains of Colorado into Grand Canyon through the Miocene East Kaibab paleocanyon (Karlstrom et al. 2014). Published in January 2014, the topographic analysis of Karlstrom and his ten coauthors prohibited Hopi Lake from extending westward at 6,000ft elevation to Kaibab Upwarp.

First Ivo Lucchitta of US Geological Survey, second William Dickinson of University of Arizona, and finally the eleven renowned *Nature Geoscience* authors asserted a Miocene age for the big river gravel deposit at Crooked Ridge just 30 mi east of the Kaibab Upwarp. These influential specialists bring up important questions. Is this the death knell for spillover theories? Should breached dam hypothesis for Grand Canyon now to be rejected? No, not yet!

### Hopi Lake Holds Water!

The popularity of this interpretation of the Miocene river was very short lived. Almost immediately, USGS geologist Richard Hereford assembled a diverse team of nine government and university geologists to reevaluate the Crooked Ridge gravel.

---

<sup>56</sup> “The biggest problem with the breached-dam hypothesis is the lack of evidence for these lakes.... Admittedly, the presumed Hopi Lake does contain a sedimentary unit known as the Bidahochi Formation, claimed by secular geologists to represent a lake environment. However,...” (Clarey 2020, 344, 345). The words “lack of evidence” and “However” kill the respectful and constructive attitude Clarey could display for Bidahochi Formation geologic evidence, and Clarey’s Bidahochi snub suspends all possible appreciation of a post-Flood Hopi Lake. Clarey is bringing up a very important big issue—the erosion of Grand Canyon. One might expect Clarey to engage in constructive discussion of Bidahochi Formation and possible lake features. That would be good scholarship. Clarey could mention tufa deposits in Bidahochi Formation resembling Lake Bonneville tufa. Oddly, Mike Oard denies the existence of Bidahochi tufa, but he accepts Lake Bonneville tufa. Clarey could mention the widespread “upper beds” with green clay and snail fossils, amphibian fossils and reptile fossils almost identical to modern Colorado Plateau freshwater species. Creationists need to hear about these. Clarey could mention hundreds of volcanic maar structures with surrounding tuff rings and scoria cones around central volcanic necks suggesting sedimentation in standing water (lake conditions), not by sedimentation in moving water (Flood conditions). Creationists need to hear about these if they are to get some informed perspective on Hopi Lake. Instead, Clarey reminds us of “lack of evidence” and “fictional lakes.” Clarey engages in the speech pattern that Rodney Dangerfield decried. Clarey illustrates geologists’ intolerance of quirkiness and their wholesale marginalizing of oddity, producing a cultural icon of “the lake that gets no respect.” Why should we expect Hopi Lake to be a normal lake? Give it respect. It’s *not* normal!

Those nine geologists submitted their response to a less-prestigious journal just seven months after *Nature Geoscience's* inditement of the western margin of Hopi Lake. Following an unprecedented 18 months in critical peer review, the conclusion of Hereford's group was finally published in April 2016 (Hereford et al. 2016). The Crooked Ridge gravel is "Early Pleistocene," not "Miocene." Also, these geologists assert the gravel on Crooked Ridge and on White Mesa (locations in fig. 4) was deposited by local streams, *not* a single regional river. That makes the river gravel *younger* than the Bidahochi Formation lake beds! Most interesting, *even remarkable*, is that the reevaluation was quickly accepted. Almost all published advocates of "Crooked Ridge Miocene River" repented by 2016.

Ironically, one of the nine geologists disputing the Miocene age assigned to the Crooked Ridge river gravel in the 2016 Hereford publication is William Dickinson, himself, a coauthor. Obviously, Dickinson repented about that supposedly Miocene River channel. Also, six of the *Nature Geoscience* authors (including Karlstrom, the paper's senior author) responded immediately to repent and affirm in print the Early Pleistocene assignment of Crooked Ridge and White Mesa in their next publication (Karlstrom et al. 2017)!

### After the Battle, the Smoke Is Clearing

We have seen a significant battle concerning the existence and identity of Hopi Lake. What can be said about this extraordinary episode? It appears we have witnessed an "Afton Canyon Controversy" repeat! It seems the breached dam hypothesis for Grand Canyon has withstood another critical test. A feature, which could have falsified specifics of the breached dam hypothesis, has, surprisingly, incorporated that evidence to make a better story. We can now imagine how small streams occupied Bidahochi Basin after Hopi Lake drained. It is now most likely that the local Ice Age stream channels at Crooked Ridge were eroded into the bed of the lake after the lake drained. The Crooked Ridge river gravel that, at first, appeared to be an insurmountable problem, has now contributed nicely to telling the story of the big lake and its drainage! Should we expect to see an Internet news release on that? Don't expect a news release because these matters concern "the lake that gets no respect." What we can expect is dismissive responses and critical speaking about spillover with little in the way of constructive proposals.

Is there any real collateral damage to Hopi Lake? Recall that William Dickinson scorned the

paleogeography of Hopi Lake by placing in its basin an imaginary Miocene Crooked River (Dickinson 2013). He used that river to reject the spillover hypothesis. Dickinson repented in print of that Miocene river. Now that the Miocene Crooked Ridge River is acknowledged to be imaginary, should Dickinson's other Hopi Lake pronouncements be considered valid?<sup>57</sup> Also, Karl Karlstrom no longer endorses a Miocene Crooked Ridge River, having repented in print. However, he and his coauthors persist in thinking that the Miocene Little Colorado River entered a small Miocene canyon through the eastern Kaibab Upwarp (Karlstrom et al. 2017). Sadly, for the *science* of geology, Hopi Lake is left out of the paleogeographic picture. Where is the geologic evidence of the Miocene Little Colorado River? Karlstrom maintains that *geologic theory* implies that river must have existed.

### Time Is Not a Magic Wand

To understand geologists' continuing fascination with the Miocene river, we must reflect on conventional geologists' notions of geologic time. Geologists assure us that the majority of the buckling of the Kaibab Upwarp and rise of the Colorado Plateau is "Laramide," as old or older than Eocene (Davis and Bump 2009), supposedly older than 40 million years. That's when the ocean left the Colorado Plateau for the last time. The uppermost member of the Bidahochi Formation, however, is Pliocene, conventionally "about 6 million years old." So, that creates an unimaginable condition: an uplifted plateau with an interior drainage basin, without a major river canyon system, after 35 million years! Now we understand why geologic time inspires geologists to dismiss Hopi Lake with disdain and to search for the Miocene river. According to theory, there must have been a Miocene river on the quarter-million-square-mile Colorado Plateau. Instead, with overtopping, geologists are acting like there is not. Therefore, spillover hypotheses are an affront to 35 million years of geologic time!

Normally, we are told that deep time (e.g., tens of millions of years) makes almost impossible things happen (e.g., natural selection causes slow organic evolution). We were also informed by that dominant twentieth century cultural myth that Grand Canyon was carved very slowly by the Colorado River over tens of millions of years. That cultural narrative was taught as fact in public grammar schools in the 1950s. Time is supposed to be a "magic wand" that makes big things happen from everyday little things. Time is championed as "hero of the plot"

<sup>57</sup> Michael Oard (2016) frequently cites Dickinson (2013) when leveling severe criticisms at Hopi Lake. During 2016, the publication year of Oard's book, all kinds of invectives were being hurled by geologists at Hopi Lake. How valid are Oard's criticisms of Hopi Lake?



in the cultural narrative. Yet, for the case of the erosion of the Colorado Plateau and Grand Canyon, time is not the magic wand, and, *certainly not*, as we are now learning, the hero of the plot! Is it possible that mainline geologists are struggling with ideas of geologic time that confuse, not illuminate, understanding the erosion of Grand Canyon? That's a question worth asking. That's a sobering question to be asking, now that geologists have been studying Grand Canyon for 160 years. No other canyon on earth has been so carefully scrutinized by geologists. Therefore, Grand Canyon is the most important landscape on planet earth to be explained by competing creationist and evolutionist paradigms.

Christians should be exploring the far-reaching implications as we rethink Grand Canyon erosion. Christians need to be ready to counter the Grand Canyon cultural narrative with the Creation, Fall, Flood and post-Flood framework of Scripture. We need to remember now that the Apostle Peter warned in these last days that many will be led astray. What is the error that will lead many astray according to the Apostle Peter? Allegiance to "...all things are continuing as they were from the beginning of creation" (2 Peter 3:4) is "...the error of lawless people" (2 Peter 3:17). "All things are continuing" summarizes what many geologists have supposed about Grand Canyon erosion.

### What We Learned

What is the lesson to be learned through the 50 years that creationists have been exploring Grand Canyon erosion? What can we "take home" from the last 30 years since that Grand Canyon rim lecture at No Name Point? New, even outrageous, ways of thinking can lead to unlikely discoveries. Creationist geologists first considered an improbable possibility of the large post-Flood Hopi Lake in northeastern Arizona. It seemed like a long shot. Then, they learned that strong geologic evidence exists for that early-post-Flood lake. Creationist thoughts were next directed toward spillover. The hypothesis fits nicely into Grand Canyon's regional story, helping to explain even the lower Colorado River with its chain of lakes and its marine delta. We conclude that creationist and catastrophist thoughts about spillover have fostered groundbreaking contributions to scholarly science and discovery. Those thoughts, so far, have survived careful scientific scrutiny. That is what would be expected if all Scripture is "inspired by God" so that even geologists can be "completely equipped" for every good work (2 Timothy 3:16).

We also learned that time is not a magic wand that solves the Grand Canyon erosion problem. Catastrophist methodology works just fine! Creationists have an excellent way of thinking and need to pay attention to detail as they continue to

develop a great explanation for the whole Southwest. This new thinking promises to help understanding of the upper Colorado River, such as a big lake in the Canyonlands area of Utah. We should remember the 50 year history of the breached dam hypothesis for erosion of Grand Canyon. Whether the hypothesis is right or wrong, it has certainly directed geologists to think in a much different direction. Sometimes big things have small beginnings.

### References

- Austin, Steven A. 1984. "Rapid Erosion at Mount St. Helens." *Origins* 11, vol.2 (June 1): 90–98. <https://grisda.org/assets/public/publications/origins/11090.pdf>
- Austin, Steven A. 1986. "Mount St. Helens and Catastrophism." In *Proceedings of the First International Conference on Creationism*. Vol.1. Edited by Robert E. Walsh, 3–9. Pittsburgh, Pennsylvania: Creation Science Fellowship.
- Austin, Steven A. ed. 1988. *Grand Canyon Field Study Tour Guidebook, April 9–16, 1988*. Santee, California: Institute for Creation Research.
- Austin, Steven A. ed. 1989. *Grand Canyon Field Study Tour Guidebook, April 8–16, 1989*, 24. Santee, California: Institute for Creation Research.
- Austin, Steven A. 1994. "How was Grand Canyon Eroded?" In *Grand Canyon—Monument to Catastrophe*. Edited by Steven A. Austin, 83–110. Santee, California: Institute for Creation Research.
- Austin, Steven A. 2009. "The Dynamic Landscape on the North Flank of Mount St. Helens." In *Volcanoes to Vineyards: Geologic Field Trips through the Dynamic Landscape of the Pacific Northwest*, 349–356. Edited by Jim E. O'Connor, Rebecca J. Dorsey, and Ian P. Madin. *Geological Society of America Field Guide 15*. Boulder, Colorado: The Geological Society of America.
- Austin, S.A., and J.A. Strelin. 2011. "Megafloods on the Santa Cruz River, Southern Argentina." *Geological Society of America Abstracts with Programs* 43, no. 5: 249.
- Babenroth, Donald L., and Arthur N. Strahler. 1945. "Geomorphology and Structure of the East Kaibab Monocline, Arizona and Utah." *Geological Society of America Bulletin* 56, no. 2 (February): 107–150.
- Billingsley, G.H. 2000. "Geologic Map of the Grand Canyon 30'×60' Quadrangle, Coconino and Mohave Counties, Northwestern Arizona." US Geological Survey, Geological Investigations Series Map I-2688. Scale 1:100,000.
- Blackwelder, Eliot. 1934. "Origin of the Colorado River." *Geological Society of America Bulletin* 45, no. 3 (30 June): 551–566.
- Blackwelder, Eliot, and Elmer William Ellsworth. 1936. "Pleistocene Lakes of the Afton Basin, California." *American Journal of Science* 31, no. 186 (June): 453–463.
- Brand, Leonard, and Arthur Chadwick. 2016. *Faith, Reason, and Earth History: A Paradigm of Earth and Biological Origins by Intelligent Design*. 3rd ed. Berrien Springs, Michigan: Andrews University Press.
- Breed, W.J. 1973. "New Avian Fossils from the Bidahochi Formation (Pliocene), Arizona." In *Cretaceous and Tertiary Rocks of the Southern Colorado Plateau*. Edited by J.E. Fassett, 144–147. *Four Corners Geological Society Memoir*. Durango, Colorado: Four Corners Geological Society.

- Bright, Jordon, Andrew S. Cohen, David L. Dettman, Philip A. Pearthree, Rebecca J. Dorsey, and Mindy B. Homan. 2016. "Did a Catastrophic Lake Spillover Integrate the Late Miocene Early Pliocene Colorado River and the Gulf of California?: Microfaunal and Stable Isotope Evidence from Blythe Basin, California-Arizona, USA." *Palaios* 31, no. 3 (March): 81–91.
- Brown, Walt. 1989. *In the Beginning: Compelling Evidence for Creation and the Flood*. 5th ed. Phoenix, Arizona: Center for Scientific Creation.
- Brown, Walt. 2008. *In the Beginning: Compelling Evidence for Creation and the Flood*. 8th ed. Phoenix, Arizona: Center for Scientific Creation.
- Brown, Walt. 2019. *In the Beginning: Compelling Evidence for Creation and the Flood*. 9th ed. Phoenix, Arizona: Center for Scientific Creation.
- Burdick, Clifford. 1974. *Canyon of Canyons*. Caldwell, Idaho: Bible-Science Association.
- Chen, Christine Y., and Adam C. Maloof. 2017. "Revisiting the Deformed High Shoreline of Lake Bonneville." *Quaternary Science Reviews* 159 (1 March): 169–189.
- Clarey, Tim. 2018. "Grand Canyon Carved by Flood Runoff." *Acts & Facts* 47, no. 12 (November 30): 10–13.
- Clarey, Timothy. 2020. *Carved in Stone, Geological Evidence of the Worldwide Flood*. Dallas, Texas: Institute for Creation Research.
- Clark, Harold W. 1946. *The New Diluvialism*. Angwin, California: Science Publications.
- Clark, Harold W. 1968. *Fossils, Flood, and Fire*. Escondido, California: Outdoor Pictures.
- Crow, Ryan, Karl Karlstrom, Andrew Darling, Laura Crossey, Victor Polyak, Darryl Granger, Yemane Asmerom and Brandon Schmandt. 2014. "Steady Incision of Grand Canyon at the Million Year Timeframe: A Case for Mantle-Driven Differential Uplift." *Earth and Planetary Sciences Letters* 397 (1 July): 159–173.
- Dallegge, Todd A., Michael H. Ort, and William C. McIntosh. 2003. "Mio-Pliocene Chronostratigraphy, Basin Morphology and Paleodrainage Relations Derived from the Bidahochi Formation, Hopi and Navajo Nations, Northeastern Arizona." *Mountain Geologist* 40, no. 3 (July 1): 55–82.
- Davis, George H., and Alex P. Bump. 2009. "Structural Geologic Evolution of the Colorado Plateau." In *Backbone of the Americas: Shallow Subduction, Plateau Uplift, and Ridge and Terrane Collision*. Edited by Suzanne Mahlburg Kay, Victor A. Ramos, and William R. Dickinson, 99–124. Geological Society of America Memoir 204. Boulder, Colorado: The Geological Society of America.
- Davis, William Morris. 1901. "An Excursion to the Grand Canyon of the Colorado." *Bulletin of the Museum of Comparative Zoology at Harvard College* 39, no. 4 (May): 107–201.
- Dickinson, William R. 2013. "Rejection of the Lake Spillover Model for Initial Incision of the Grand Canyon, and Discussion of Alternatives." *Geosphere* 9, no. 1 (February 1): 1–20.
- Dorsey, Rebecca J., Brennan O'Connell, Kristin McDougall, and Mindy B. Homan. 2018. "Punctuated Sediment Discharge during Early Pliocene Birth of the Colorado River: Evidence from Regional Stratigraphy, Sedimentology, and Paleontology." *Sedimentary Geology* 363 (January): 1–33.
- Douglass, John, Norman Meek, Ronald I. Dorn, and Mark W. Schmeekle. 2009. "A Criteria-Based Methodology for Determining the Mechanism of Transverse Drainage Development, with Application to the Southwestern United States." *Geological Society of America Bulletin* 121, nos. 3–4 (March): 586–598.
- Douglass, J. 2011. "One Grand Canyon but Four Mechanisms—Was It Antecedence, Superimposition, Overflow, or Piracy?" In *CRevolution 2—Origin and Evolution of the Colorado River System*, Workshop Abstracts. Edited by L. Sue Beard, Karl E. Karlstrom, Richard A. Young, and George H. Billingsley, 93–98. U.S. Geological Survey Open-File Report 2011-1210.
- Douglass, John C., Brian F. Gootee, Todd Dallegge, A. Jeong, Y.B. Seong, and Y.B. Yu. 2020. "Evidence for the Overflow Origin of Grand Canyon." *Geomorphology* 369 (pre-proof version July 22, 2020).
- Ellsworth, Elmer William. 1932. "Physiographic History of the Afton Basin." Ph.D. diss. Stanford University. Stanford, California.
- Ellsworth, Scott A. 1999. "Memorial to Elmer William Ellsworth 1907–1997." *Geological Society of America Memorials* 30: 11–14.
- Enzel, Yehouda, Stephen G. Wells, and Nicholas Lancaster. 2003. "Late Pleistocene Lakes Along the Mojave River, Southeast California." In *Paleoenvironments and Paleohydrology of the Mojave and Southern Great Basin Deserts*. Edited by Yehouda Enzel, Stephen G. Wells, and Nicholas Lancaster, 61–77. Boulder, Colorado: Geological Society of America Special Paper 368.
- Falvey, Henry T. 1990. *Cavitation in Chutes and Spillways*. Engineering Monograph 42. Denver, Colorado: Bureau of Reclamation.
- Felton, Alisa, Paul W. Jewell, Marjorie Chan, and Donald Currey. 2006. "Controls of Tufa Development in Pleistocene Lake Bonneville, Utah." *The Journal of Geology* 114, no. 3 (May): 377–389.
- Gould, Stephen Jay. 1985. *Ontogeny and Phylogeny*. Cambridge, Massachusetts: Harvard University Press.
- Gross, Essa L., P. Jonathan Patchett, Todd A. Dallegge, and Jon E. Spencer. 2001. "The Colorado River System and Neogene Sedimentary Formations Along Its Course: Apparent Sr Isotopic Connections." *The Journal of Geology* 109, no. 4 (July): 449–461.
- Hansen, Wallace R. 1965. "The Black Canyon of the Gunnison: Today and Yesterday." *United States Geological Survey Bulletin* 1191: 76.
- Harris, Raymond C., Robert S. Leighty, Robert B. Scarborough, and Jon E. Spencer. 1998. "Uranium Levels and Radon Potential in Selected Areas North of Phoenix, in the St. Johns Area, and North of Tuba City, Arizona." Arizona Geological Survey Open-File Report 98-10. Tucson, Arizona: Arizona Geological Survey.
- Helble, Tim, and Carol Hill. 2016. "Carving of the Grand Canyon: A Lot of Time and a Little Water, a Lot of Water and a Little Time (or Something Else?)." In *The Grand Canyon, Monument to an Ancient Earth: Can Noah's Flood Explain the Grand Canyon?* Edited by Carol Hill, Gregg Davidson, Tim Helble, and Wayne Ranney, 170. Grand Rapids, Michigan: Kregel Publications.
- Hereford, Richard, L. Sue Beard, William R. Dickinson, Karl E. Karlstrom, Matthew T. Heizler, Laura J. Crossey, Lee Amarosa, P. Kyle House, and Mark Pecha. 2016. "Reevaluation of the Crooked Ridge River—Early

- Pleistocene (ca.2Ma) Age and Origin of the White Mesa Alluvium, Northeastern Arizona." *Geosphere* 12, no. 3 (June 1): 768–789.
- Hilgendorf, Zach, Greta Wells, Phillip H. Larson, Jason Millett, and Melissa Kohout. 2020. "From Basins to Rivers: Understanding the Revitalization and Significance of Top-Down Drainage Integration Mechanisms in Drainage Basin Evolution." *Geomorphology* 352 (March): 1–17.
- Holm, R. F. 2001. "Pliocene-Pleistocene Incision on the Mogollon Slope, Northern Arizona: Response to the Developing Grand Canyon." In *Colorado River Origin and Evolution*. Monograph 12. Edited by Richard A. Young, and Earle E. Spamer, 59–63. Grand Canyon, Arizona, Grand Canyon Association.
- Holroyd, Edmond W. 1987. "Missing Talus." *Creation Research Society Quarterly* 24, no. 1 (June 1987): 1–5.
- Holroyd, Edmond W. 1990a. "Some Simulations of the Possible Role of Cavitation in Catastrophic Floods." *Creation Research Society Quarterly* 27, no. 2 (September): 49–55.
- Holroyd, Edmond W. 1990b. "Missing Talus on the Colorado Plateau." In *Proceedings of the Second International Conference on Creationism*. Vol. 2. Edited by Robert E. Walsh and Christopher L. Brooks, 115–128. Pittsburgh, Pennsylvania: Creation Science Fellowship.
- Holroyd, Edmond W. 1994. "A Remote Sensing Search for Extinct Lake Shore Lines on the Colorado Plateau." In *Proceedings of the Third International Conference on Creationism*. Edited by Robert E. Walsh, 243–254. Pittsburgh, Pennsylvania: Creation Science Fellowship.
- House, P. Kyle. 2008. "The Birth of the Lower Colorado River: Are Spilling Lakes All It Takes?" <https://www.slideshare.net/drpkhouse/birth-of-the-lower-colorado-river-presentation>.
- House, P. Kyle, Philip A. Pearthree, and Michael E. Perkins. 2008. "Stratigraphic Evidence for the Role of Lake Spillover in the Inception of the Lower Colorado River in Southern Nevada and Western Arizona." In *Late Cenozoic Drainage History of the Southwestern Great Basin and Lower Colorado River Region: Geologic and Biotic Perspectives*. Edited by Marith C. Reheis, Robert Hershler, and David M. Miller, 335–353. Geological Society of America Special Paper 439.
- Karlstrom, Karl E., John P. Lee, Shari A. Kelley, Ryan S. Crow, Laura J. Crosse, R. A. Young, Greg Lazear, L. Sue Beard, Jason W. Ricketts, Matthew Fox, and David L. Shuster. 2014. "Formation of the Grand Canyon 5 to 6 Million Years Ago Through Integration of Older Palaeocanyons." *Nature Geoscience* 7, no. 1 (26 January): 239–244.
- Karlstrom, K. E., L. J. Crosse, E. Embid, R. Crow, M. Heizler, R. Hereford, L. S. Beard, J. W. Ricketts, S. Cather, and S. Kelley. 2017. "Cenozoic Incision History of the Little Colorado River: Its Role in Carving Grand Canyon and Onset of Rapid Incision in the Past ca. 2Ma in the Colorado River System." *Geosphere* 13, no. 1 (February 1): 49–81.
- Larson, Phillip H., Ronald I. Dorn, R. Evan Palmer, Zack Bowles, Emma Harrison, Scott Kelley, Mark W. Schmeeckle, and John Douglass. 2014. "Pediment Response to Drainage Basin Evolution in South-Central Arizona." *Physical Geography* 35, no. 5 (September): 369–389.
- Larson, Phillip H., Norman Meek, John Douglass, Ronald I. Dorn, and Yeong Bae Seong. 2017. "How Rivers Get Across Mountains: Transverse Drainages." *Annals of the American Association of Geographers* 107, no. 2 (March): 274–283.
- Longwell, Chester R. 1928. "Geology of the Muddy Mountains Nevada; With a Section Through the Virgin Range to the Grand Wash Cliffs, Arizona." *U.S. Geological Survey Bulletin* 798: 91–97.
- Lucchitta, Ivo, Richard F. Holm, and Baerbel K. Lucchitta, 2011, "A Miocene River in Northern Arizona and Its Implications for the Colorado River and Grand Canyon." *GSA Today* 21, no. 10 (October): 4–10.
- Lucchitta, Ivo, Richard F. Holm, and Baerbel K. Lucchitta. 2013. "Implications of the Miocene(?) Crooked Ridge River of Northern Arizona for the Evolution of the Colorado River in Grand Canyon." *Geosphere* 9, no. 6 (December 1): 1417–1433.
- Meek, Norman. 1989. "Geomorphic and Hydrologic Implications of the Rapid Incision of Afton Canyon, Mojave Desert, California." *Geology* 17, no. 1 (January 1): 7–10.
- Meek, Norman. 1990. *Late Quaternary Geochronology and Geomorphology of the Manix Basin, San Bernardino County, California*. Ph.D. diss. University of California, Los Angeles, California.
- Meek, Norman, and John Douglass. 2001. "Lake Overflow: An Alternative Hypothesis for Grand Canyon Incision and Development of the Colorado River." In *Colorado River: Origin and Evolution*. Edited by R. A. Young and E. E. Spamer, 199–204. Grand Canyon Association Monograph 12. Grand Canyon, Arizona: Grand Canyon Association.
- Meek, Norman. 2002. "Ponding and Overflow: The Forgotten Transverse Drainage Hypothesis." *98th Annual Meeting Abstracts*. Los Angeles, California: Association of American Geographers.
- Meek, Norman. 2019. "Episodic Forward Prolongation of Trunk Channels in the Western United States." *Geomorphology* 340 (1 September): 172–183.
- Mitchell, Elizabeth. 2013. "What Carved the Grand Canyon?" *Answers in Genesis News to Know*, January 5. <https://answersingenesis.org/geology/grand-canyon-facts/what-carved-grand-canyon/>.
- Nelson, Byron C. 1931. *The Deluge Story in Stone: A History of the Flood Theory of Geology*. Minneapolis, Minnesota: Augsburg Publishing House.
- Newberry, J.S. 1861. *Report Upon the Colorado River of the West*. Part III. Washington, DC: U.S. Government Printing Office.
- Newberry, J.S. 1862. "Colorado River of the West" *American Journal of Science* 33:376–403.
- Northrup, Bernard E. 2004. "The Grand Canyon and Biblical Catastrophes." *Chafer Theological Seminary Journal* 10, no. 2 (Fall): 1–27.
- Oard, Michael J. 1993. "Comments on the Breached Dam Theory for the Formation of the Grand Canyon." *Creation Research Society Quarterly* 30, no. 1 (June): 39–46.
- Oard, Michael J. 2010. "The Origin of Grand Canyon, Part II: Fatal Problems with the Dam-Breach Hypothesis." *Creation Research Society Quarterly* 46, no. 4 (Spring): 290–307.
- Oard, Michael J. 2016. *A Grand Origin for Grand Canyon*. Chino Valley, Arizona: Creation Research Society Books.
- Oskin, Becky. 2012. Grand Canyon Carved by Flood? Geologist Says No. *Yahoo News LiveScience.com*, December 19. <https://www.yahoo.com/news/grand-canyon-carved-flood-geologist-says-no-202017292.html>.
- Paiva, C.A. 1988. *Cavitation in Macro-Fluvial Processes and the Implications for Geologic Time*. Master's thesis. Santee, California: Institute for Creation Research.

- Parmley, Dennis, and David L. Peck. 2002. "Amphibians and Reptiles of the Late Hemphillian White Cone Local Fauna, Navajo County, Arizona." *Journal of Vertebrate Paleontology* 22, no. 1 (March): 175–178.
- Powell, J.W. 1875. *Exploration of the Colorado River of the West and Its Tributaries*. Washington, D.C.: U.S. Government Printing Office.
- Ranney, Wayne. 2012. *Carving Grand Canyon: Evidence, Theories, and Mystery*. 2nd ed. Grand Canyon, Arizona: Grand Canyon Association.
- Reheis, Marith C., and Joanna L. Redwine. 2008. "Lake Manix Shorelines and Afton Canyon Terraces: Implications for Incision of Afton Canyon." In *Late Cenozoic Drainage History of the Southwestern Great Basin and Lower Colorado River Region: Geologic and Biotic Perspectives*. Edited by Marith C. Reheis, Robert Hershler, and David M. Miller, 227–259. Geological Society of America Special Paper 439.
- Reheis, Marith C., Joanna R. Redwine, Elmira Wan, John P. McGeehin, and D. Paco VanSistine. 2014. "Surficial Geology and Stratigraphy of Pleistocene Lake Manix, San Bernardino County, California." *U.S. Geological Survey Scientific Investigations Map 3312*.
- Scarborough, R.B. 1989. "Cenozoic Erosion and Sedimentation in Arizona." In *Geologic Evolution of Arizona*. Edited by J.P. Jenney, and S.J. Reynolds, 515–537. Arizona Geological Society Digest, Vol. 17.
- Scarborough, R. 2001. "Neogene Development of Little Colorado River Valley and Eastern Grand Canyon: Field Evidence for an Overtopping Hypothesis." In *Colorado River: Origin and Evolution*. Edited by R.A. Young, and E.E. Spamer. Grand Canyon Association Monograph 12. Grand Canyon, Arizona: Grand Canyon Association.
- Scheele, Peter. 2010. "A Receding Flood Scenario for the Origin of the Grand Canyon." *Journal of Creation* 24, no. 3 (December): 106–116.
- Smith, Gerald R., Robert E. Reynolds, and Joseph D. Stewart. 2013. "Hydrographic Significance of Fishes from the Early Pliocene White Narrows Beds, Clark County, Nevada." In *Raising Questions in the Central Mojave Desert*. Edited by Robert E. Reynolds, 171–180. Zzyzx, California: California State University, Desert Studies Center.
- Snelling, Andrew A. and Tom Vail. 2009. "When and How Did the Grand Canyon Form?" In *The New Answers Book 3*. Edited by Ken Ham, 173–186. Green Forest, Arkansas. Master Books.
- Spencer, J.E., and P.A. Pearthree. 2001. "Headward Erosion Versus Closed-Basin Spillover as Alternative Causes of Neogene Capture of the Ancestral Colorado River by the Gulf of California." In *Colorado River: Origin and Evolution*. Edited by R.A. Young and E.E. Spamer, 215–222. Grand Canyon Association Monograph 12. Grand Canyon, Arizona: Grand Canyon Association.
- Spencer, J.E., P.A. Pearthree, and P.K. House. 2008. "An Evaluation of the Evolution of the Latest Miocene to Earliest Pliocene Bouse Lake System in the Lower Colorado River Valley, Southwestern USA." In *Late Cenozoic Drainage History of the Southwestern Great Basin and Lower Colorado River Region: Geologic and Biotic Perspectives*. Edited by Marith C. Reheis, Robert Hershler, and David M. Miller, 375–390. Geological Society of America Special Paper 439.
- Spencer, Jon E., P. Jonathan Patchett, Philip A. Pearthree, P. Kyle. House, Andrei M. Sarna-Wojcicki, Elmira Wan, Jennifer A. Roskowski, and James E. Faulds. 2013. "Review and Analysis of the Age and Origin of the Pliocene Bouse Formation, Lower Colorado River Valley, Southwestern USA." *Geosphere* 9, no. 3 (June 1): 444–459.
- Stirton, R.A. 1936. "A New Beaver from the Pliocene of Arizona, with Notes on the Species of Dipoides." *Journal of Mammalogy* 17, no. 3 (14 August): 279–281.
- Taylor, Dwight W. 1957. "Pliocene Fresh-Water Mollusks from Navajo County, Arizona." *Journal of Paleontology* 31, no. 3 (May): 654–661.
- Uyeno, Teruya, and Robert R. Miller. 1965. "Middle Pliocene Cyprinid Fishes from the Bidahochi Formation, Arizona." *Copeia* 1965, no. 1 (March): 28–41.
- Vail, Tom. 2003. *Grand Canyon: A Different View*. Green Forest, Arkansas: New Leaf Publishing.
- Walcott, C.D. 1890. "Study of a Line of Displacement in the Grand Canyon of the Colorado, in Northern Arizona." *Geological Society of America Bulletin* 1, no. 1: 49–64.
- Wells, S.G., and Y. Enzel. 1994. "Fluvial Geomorphology of the Mojave River in the Afton Canyon Area, Eastern California: Implications for the Geomorphic Evolution of Afton Canyon." In *Geological Investigations of an Active Margin*. Edited by Sally F. McGill, and Timothy M. Ross, 177–188. Cordilleran Section Guidebook of the Geological Society of America. Boulder, Colorado: Geological Society of America.
- Wells, Stephen G., William J. Brown, Yehouda Enzel, Roger Y. Anderson, and Leslie D. McFadden. 2003. "Late Quaternary Geology and Paleohydrology of Pluvial Lake Mojave, Southern California." In *Paleoenvironments and Paleohydrology of the Mojave and Southern Great Basin Deserts*. Edited by Yehouda Enzel, Stephen G. Wells, and Nicholas Lancaster, 79–114. Geological Society of America Special Paper 368. Boulder, Colorado: Geological Society of America.
- White, J.D.L. 1990. "Depositional Architecture of a Maar-Pitted Playa: Sedimentation in the Hopi Buttes Volcanic Field, Northeastern Arizona, U.S.A." *Sedimentary Geology* 67, nos. 1–2 (April): 55–84.
- Whitcomb, John C. and Henry M. Morris. 1961. *The Genesis Flood: The Biblical Record and Its Scientific Implications*. Phillipsburg, New Jersey: Presbyterian and Reformed Publishing Co.
- Whitmore, J.H. 1985. "The Origin of the Colorado River Across the Kaibab Upwarp: Four Models" (Unpublished). Submitted to Dr. Austin's Geology 501 class, October 7, 1985. Institute for Creation Research Graduate School, Geology Department.
- Williams, Emmett L., John R. Meyer, and Glen W. Wolfrom. 1992. "Erosion of the Grand Canyon of the Colorado River: Part III—Review of the Possible Formation of Basins and Lakes on the Colorado Plateau and Different Climatic Conditions in the Past." *Creation Research Society Quarterly* 29, no. 1 (June): 18–24.
- Williams, Howel. 1936. "Pliocene Volcanoes of the Navajo-Hopi Country." *Bulletin of the Geological Society of America* 47 (January 31): 111–172.
- Zelawski, Mallory. 2011. "Marginal Deposits of Maar Volcanoes in the First Mesa Area, Hopi Buttes Volcanic Field, Navajo Nation, Arizona." M.S. Thesis, Northern Arizona University. Flagstaff, Arizona.