

On the Complexity of Maslov's Class \overline{K} (Extended Abstract)

Oskar Fiuk¹, Emanuel Kieroński¹ and Vincent Michielini²

¹University of Wrocław, Institute of Computer Science, Wrocław, Poland

²University of Warsaw, Faculty of Mathematics, Informatics, and Mechanics, Warsaw, Poland

Abstract

Maslov's class \overline{K} is an expressive fragment of First-Order Logic that embeds modal logic and many description logics, including \mathcal{ALC} . It is known to have decidable satisfiability problem, whose exact complexity, however, has not been established so far. We show that \overline{K} has the exponential-sized model property, and hence its satisfiability problem is NEXPTIME-complete. Additionally, we get new complexity results on related fragments studied in the literature, and propose a new decidable extension of the uniform one-dimensional fragment (without equality). Our approach involves a use of satisfiability games tailored to \overline{K} and a novel application of paradoxical tournament graphs.

Keywords

satisfiability problem, finite model property, Maslov's class K, paradoxical tournaments

1. Introduction

Basic modal logic and many standard description logics (DLs), including \mathcal{ALC} , embed into First-Order Logic, FO, via the so-called *standard translation*. In contrast to robust decidability of satisfiability of DLs, the satisfiability problem for full FO is undecidable. Therefore, a considerable effort has been made to identify fragments of FO which still embed standard DLs, but have decidable satisfiability. Studying such fragments may help us to understand good computational and model-theoretic properties of DLs, and find their attractive extensions, e.g., admitting relations of arity greater than two.

The list of known decidable fragments that appeared in this line of research includes the two-variable fragment, FO² [1, 2], the guarded fragment, GF [3, 4], the unary negation fragment, UNFO [5], the guarded negation fragment, GNFO [6], the fluted fragment, FF [7, 8], its generalisation the adjacent fragment, AF [9], and the uniform one-dimensional fragment, UF₁ [10, 11, 12]. A survey [13] puts in this context also Maslov's class \overline{K} [14].¹


In contrast to all the other fragments mentioned above, satisfiability of \overline{K} is not yet fully understood. Maslov proved the decidability of the validity problem for K, which is equivalent to the satisfiability problem for \overline{K} , using his own approach, which he called the *inverse method*. There were a few subsequent works [16, 17, 18], whose authors reproved this result by means of the resolution method; all of them work directly with \overline{K} . None of these works, however, studied the complexity of its satisfiability. While a closer inspection of the resolution-based procedure in [17] seems to allow one to extract some elementary upper bound on the complexity of the satisfiability problem for \overline{K} , it would not be easy to get anything better than doubly exponential. This would still leave a gap, as the best lower bound inherited from the fragments embeddable in \overline{K} is NEXPTIME-hardness.

In our paper we add the main missing brick to the understanding of \overline{K} by showing that its satisfiability problem is NEXPTIME-complete. We will do it by showing:

 DL 2024: 37th International Workshop on Description Logics, June 18–21, 2024, Bergen, Norway

 307023@uwr.edu.pl (O. Fiuk); emanuel.kieronski@cs.uni.wroc.pl (E. Kieroński); michielini@mimuw.edu.pl (V. Michielini)

 0009-0006-1312-4899 (O. Fiuk); 0000-0002-8538-8221 (E. Kieroński); 0000-0002-1413-9316 (V. Michielini)

 © 2024 Copyright for this paper by its authors. Use permitted under Creative Commons License Attribution 4.0 International (CC BY 4.0).

¹The reader should not confuse the class \overline{K} with another class, called just the *Maslov class* (see, e.g. [15]).

Theorem 1. *Every satisfiable formula φ in $\overline{\mathcal{K}}$ admits a finite model of size $2^{\mathcal{O}(|\varphi| \cdot \log |\varphi|)}$. Hence, the satisfiability problem for $\overline{\mathcal{K}}$ is NEXPTIME-complete.*

For an extended version of this work see [19].

2. Maslov's Class $\overline{\mathcal{K}}$ and its relation to other logics

We consider signatures with relational symbols of arbitrary arity and constants, but no function symbols of arity greater than 0. Equalities are not allowed as they lead to undecidability [20]. Let φ be a sentence in negation normal form, and let $R(\bar{y})$ be one of its atoms. The φ -prefix of $R(\bar{y})$ is the sequence of quantifiers in φ binding the variables of $R(\bar{y})$. For instance, if φ is the sentence $\exists x. \forall y. \exists z. R(x, y) \wedge T(c, y, z, y)$, with c being a constant symbol, then the φ -prefix of the atom $R(x, y)$ is the sequence “ $\exists x. \forall y$ ”, while that of the atom $T(c, y, z, y)$ is “ $\forall y. \exists z$ ”. An atom without variables (e.g. stating only about constants) has the empty φ -prefix.

The class $\overline{\mathcal{K}}$ consists of the sentences φ which are in negation normal form and in which there exist universally quantified variables x_1, \dots, x_K , called *special variables*, none of which lying within the scope of any existential quantifier, such that each atom of φ has a φ -prefix of one of the following shapes: (i) a φ -prefix of length at most 1, (ii) a φ -prefix ending with an existential quantifier, (iii) or exactly the sequence “ $\forall x_1 \dots \forall x_K$ ”.

While we require the formula φ to be in negation normal form, we allow ourselves to use implications, provided that their left-hand sides do not contain any quantifiers. With this convention, the following formula $\varphi_{\text{co_authors}}$ is in $\overline{\mathcal{K}}$:

$$\begin{aligned} & \forall s_1, s_2, s_3. [\text{scientist}(s_1) \wedge \text{scientist}(s_2) \wedge \text{scientist}(s_3) \wedge \text{co_authors}(s_1, s_2, s_3)] \\ & \rightarrow \exists a. \text{article}(a) \wedge \text{written_by}(a, s_1, s_2, s_3). \end{aligned}$$

Indeed, the $\varphi_{\text{co_authors}}$ -prefixes of the different atoms are: the singleton sequences “ $\forall s_1$ ”, “ $\forall s_2$ ”, “ $\forall s_3$ ” and “ $\exists a$ ”; the sequence “ $\forall s_1. \forall s_2. \forall s_3. \exists a$ ”, which ends with an existential quantifier; and the universal sequence “ $\forall s_1. \forall s_2. \forall s_3$ ”. There is no existential quantifier binding the s_i , so all the conditions are met, with s_1, s_2, s_3 being the special variables.

Another example $\varphi_{\text{marriage}}$ demonstrates the possibility of using quantifier alternation:

$$\begin{aligned} & \forall h, w. \text{husband_and_wife}(h, w) \rightarrow \exists p. \text{problem}(p) \wedge \forall d. \text{date}(d) \rightarrow \\ & \exists d'. \text{date}(d') \wedge \text{later_than}(d', d) \wedge \text{occurs_to_at}(p, h, w, d'). \end{aligned}$$

On the contrary, an example of a property not expressible in $\overline{\mathcal{K}}$ is transitivity. In particular, in the formula $\forall x, y, z. [T(x, y) \wedge T(y, z)] \rightarrow T(x, z)$, one cannot find a subset of variables that could work as the set of special variables.

Turning to relation between $\overline{\mathcal{K}}$ and DLs, we need to remark, that actually the standard translation of, say, \mathcal{ALC} into FO does not land directly in $\overline{\mathcal{K}}$. To embed \mathcal{ALC} in $\overline{\mathcal{K}}$, we first translate \mathcal{ALC} into FO^2 , and then write the FO^2 formula in its *Scott normal form*: $\forall x, y. \varphi_0(x, y) \wedge \bigwedge_i \forall x. \exists y. \varphi_i(x, y)$. One can verify that after renaming the variables in the $\forall\exists$ -conjuncts we indeed get sentences in $\overline{\mathcal{K}}$.

Additional DL features that translate to $\overline{\mathcal{K}}$ this way include Boolean combinations of roles, inversions, role restrictions and positive occurrences of role compositions [13].

In addition to DLs, the class $\overline{\mathcal{K}}$ embeds, either syntactically or via simple reductions preserving satisfiability, many known decidable fragments of First-Order Logic, including the monadic class [21], the Ackermann fragment [22] and its generalised version [23], the Gödel class [24], the two-variable fragment [1] and Class 2.4 from [25] (which we will call $\overline{\mathcal{K}}$ -Skolem class). Even more formalisms, for example the uniform one-dimensional fragment [10] or its variation with alternation of quantifiers in blocks [26], are captured by the class $\overline{\text{DK}}$ of conjunctions of $\overline{\mathcal{K}}$ -sentences, also known to be decidable [17].

3. Our results

Our main result is Theorem 1: the class $\overline{\mathbf{K}}$ possesses the exponential-sized model property, and hence its satisfiability problem is NEXPTIME-complete. In addition, it establishes NEXPTIME-completeness of two subfragments of $\overline{\mathbf{K}}$ studied in the literature, whose complexity has not been known so far: the Generalised Ackermann Class (without equality) and the $\overline{\mathbf{K}}$ -Skolem class, the latter being the intersection of $\overline{\mathbf{K}}$ and the prefix-class $\forall^*\exists^*$ (the Skolem class). The matching NEXPTIME-hardness lower bound holds already for the subclass $\forall\forall\exists$ of the latter [27]. Our remaining results are as follows:

- (A) We show that our upper bound on the size of minimal models (and hence also on the complexity) for $\overline{\mathbf{K}}$ transfers to $\overline{\mathbf{DK}}$, the class of conjunctions of $\overline{\mathbf{K}}$ -sentences.
- (B) We show that this upper bound is essentially optimal by supplying a family of tight examples, contained already in $\overline{\mathbf{K}}$ -Skolem: we construct a sequence $(\varphi_n)_{n \in \mathbb{N}}$ of satisfiable sentences such that each φ_n has size linear in n and its models have size at least $2^{\Omega(n \cdot \log n)}$. We can construct such sentences even without constants and with just one existential quantifier.
- (C) We show that satisfiable sentences φ in $\overline{\mathbf{K}}$ have models of size $2^{O(|\varphi|)}$, under the assumption that the number of universal quantifiers is bounded; this in particular applies to the Gödel class admitting two universal quantifiers.
- (D) We propose a novel generalisation of the uniform one-dimensional fragment of First-Order Logic [10]: the \forall -uniform fragment. We show an efficient satisfiability preserving translation to $\overline{\mathbf{DK}}$, and this way we obtain the exponential-sized model property and NEXPTIME-completeness of the new fragment.

4. Proof strategy

In contrast to the previous works on $\overline{\mathbf{K}}$, which approached the problem *syntactically*, we do it *semantically*. Let us give a taste of our ideas here; the details can be found in [19].

In our small model construction, a crucial role is played by paradoxical tournament graphs. Let us recall that a directed graph without self-loops is a *tournament* if it contains exactly one directed arc between every pair of vertices, and a tournament is *k-paradoxical* if, for every subset of vertices A of cardinality at most k , there is a vertex b *dominating* A , that is sending arcs to all elements of A . It is a classical result by Erdős that such tournaments of size $\mathcal{O}(k^2 \cdot 2^k)$ exist [28]. Inspired by this classical result, we introduce a variant of tournament graphs with colours of vertices and of arcs.

Let us now explain how we employ such tournaments. Suppose that we want to check the satisfiability of a $\overline{\mathbf{K}}$ -sentence $\varphi: \forall x_1, x_2. \exists y. \psi_1 \wedge \psi_2 \wedge \psi_3$, where ψ_1 is $[\neg R(x_1, x_2) \vee \neg R(x_2, x_1)] \wedge [\neg R(x_1, x_2) \vee \neg S(x_1, x_2)]$, saying that R is antisymmetric (thus also irreflexive) and disjoint from S ; ψ_2 is $R(x_1, x_2) \rightarrow [R(y, x_1) \wedge S(y, x_2)]$, stating that each two elements x_1, x_2 , connected via R , share a common predecessor y via R for x_1 and via S for x_2 ; and ψ_3 is $\neg U(x_1) \leftrightarrow U(y)$, requiring that x_1 and y disagree on U .

We consider a tournament $\mathcal{T}=(V, E)$, where arcs are coloured by $\lambda: E \rightarrow \{x_1, x_2\}$ and vertices are coloured by $\mu: V \rightarrow \{U, \neg U\}$. We construct a model \mathfrak{A} over the domain V :

- (i) For each $a \in V$, if its colour via μ is U , we set $\mathfrak{A} \models U(a)$.
- (ii) For each $a, b \in V$, if (a, b) is an arc in E with colour x_1 via λ , we set $\mathfrak{A} \models R(a, b)$.
- (iii) For each $a, b \in V$, if (a, b) is an arc in E with colour x_2 via λ , we set $\mathfrak{A} \models S(a, b)$.
- (iv) Every other atom of \mathfrak{A} is set to be false.

Now, imagine that our tournament \mathcal{T} has the following paradoxical-like property: *for any different v_1, v_2 in V and any colour c in $\{U, \neg U\}$, there exists another vertex w in V such that its colour via μ is c , and there are arcs $(w, v_1), (w, v_2)$ in E with x_1, x_2 being their respective colours via λ .* In this case, \mathfrak{A} indeed becomes a model of φ . In particular, $\mathfrak{A} \models$

$\exists y. R(y, v_1) \wedge S(y, v_2) \wedge (\neg U(v_1) \leftrightarrow U(y))$ for any $v_1 \neq v_2$, as the described property applied to v_1, v_2 with $c = \neg U$ if $\mu(v_1) = U$ and $c = U$ otherwise, gives us an appropriate witness w . The existence of \mathcal{T} satisfying this property follows from our work.

In the general case, we propose a game-theoretic view for the problem: with every \overline{K} -sentence φ we associate a *satisfiability game*, between two players, named Eloisa and Abelard. It resembles a classical *verification game*, but with the structure not being explicitly given; instead players construct it during the play. We show that Eloisa has a winning strategy iff φ is satisfiable. In establishing the “only if” direction, a crucial role is played by the aforementioned paradoxical tournaments: colours of arcs are variables of φ , and colours of vertices are certain positions from the game, corresponding to partial structures constructed by players. The core of the model is the tournament, and the atoms are specified in accordance with the colours, similarly as in our example above.

Acknowledgments. The first and second authors were supported by Polish National Science Center grant No. 2021/41/B/ST6/00996. The third author was supported by the ERC grant INFSYS, agreement no. 950398. The authors thank Warren Goldfarb, Ullrich Hustadt and Harry R. Lewis for some helpful comments on their work.

References

- [1] D. Scott, A decision method for validity of sentences in two variables, *Journal Symbolic Logic* 27 (1962) 477.
- [2] E. Grädel, P. Kolaitis, M. Y. Vardi, On the decision problem for two-variable first-order logic, *Bulletin of Symbolic Logic* 3 (1997) 53–69. doi:10.2307/421196.
- [3] H. Andréka, J. van Benthem, I. Németi, Modal languages and bounded fragments of predicate logic, *Journal of Philosophical Logic* 27 (1998) 217–274. doi:10.1023/A:1004275029985.
- [4] E. Grädel, On the restraining power of guards, *J. Symb. Log.* 64 (1999) 1719–1742. doi:10.2307/2586808.
- [5] B. ten Cate, L. Segoufin, Unary negation, *Logical Methods in Comp. Sc.* 9 (2013). doi:10.2168/LMCS-9(3:25)2013.
- [6] V. Bárány, B. ten Cate, L. Segoufin, Guarded negation, *J. ACM* 62 (2015) 22. doi:10.1145/2701414.
- [7] W. V. Quine, On the limits of decision, in: *Proceedings of the 14th International Congress of Philosophy*, volume III, 1969, pp. 57–62.
- [8] I. Pratt-Hartmann, W. Szwaig, L. Tendera, The fluted fragment revisited, *J. Symb. Log.* 84 (2019) 1020–1048. doi:10.1017/JSL.2019.33.
- [9] B. Bednarczyk, D. Kojelis, I. Pratt-Hartmann, On the limits of decision: the adjacent fragment of first-order logic, in: *50th International Colloquium on Automata, Languages, and Programming, ICALP 2023*, volume 261 of *LIPICs*, Schloss Dagstuhl - Leibniz-Zentrum für Informatik, 2023, pp. 111:1–111:21. doi:10.4230/LIPICs.ICALP.2023.111.
- [10] L. Hella, A. Kuusisto, One-dimensional fragment of first-order logic, in: *Proceedings of Advances in Modal Logic*, 2014, 2014, pp. 274–293.
- [11] E. Kieronski, A. Kuusisto, Complexity and expressivity of uniform one-dimensional fragment with equality, in: *Mathematical Foundations of Computer Science 2014 - 39th International Symposium, MFCS 2014, Proceedings, Part I*, volume 8634 of *Lecture Notes in Computer Science*, 2014, pp. 365–376. doi:10.1007/978-3-662-44522-8_31.
- [12] A. Kuusisto, On the uniform one-dimensional fragment, in: *Proceedings of the 29th International Workshop on Description Logics*, 2016.
- [13] U. Hustadt, R. A. Schmidt, L. Georgieva, A survey of decidable first-order fragments and description logics, *Journal of Relational Methods in Computer Science* 1 (2004) 2004.
- [14] S. J. Maslov, The inverse method for establishing deducibility for logical calculi, *The Calculi of Symbolic Logic I: Proceedings of the Steklov Institute of Mathematics* 98 (1971).

- [15] E. Börger, E. Grädel, Y. Gurevich, The classical decision problem, *Perspectives in Mathematical Logic*, Springer, 1997.
- [16] C. G. Fermüller, A. Leitsch, T. Tammet, N. K. Zamov, Resolution Methods for the Decision Problem, volume 679 of *Lecture Notes in Computer Science*, Springer, 1993. doi:10.1007/3-540-56732-1.
- [17] U. Hustadt, R. A. Schmidt, Maslov’s class K revisited, in: Automated Deduction - CADE-16, 16th International Conference on Automated Deduction 1999, Proceedings, volume 1632 of *Lecture Notes in Computer Science*, Springer, 1999, pp. 172–186. doi:10.1007/3-540-48660-7_12.
- [18] C. G. Fermüller, A. Leitsch, U. Hustadt, T. Tammet, Resolution decision procedures, in: J. A. Robinson, A. Voronkov (Eds.), *Handbook of Automated Reasoning* (in 2 volumes), Elsevier and MIT Press, 2001, pp. 1791–1849. doi:10.1016/B978-044450813-3/50027-8.
- [19] O. Fiuk, E. Kieronski, V. Michielini, On the complexity of Maslov class K, in: 39th Annual ACM/IEEE Symposium on Logic in Computer Science, LICS 2024, 2024, pp. 1–14. doi:10.1145/3661814.3662097.
- [20] W. D. Goldfarb, The unsolvability of the Gödel class with identity, *J. Symb. Logic* 49 (1984) 1237–1252. doi:10.2307/2274274.
- [21] L. Löwenheim, Über möglichkeiten im relativkalkül, *Mathematische Annalen* 76 (1915) 447–470.
- [22] W. Ackermann, Über die erfüllbarkeit gewisser zähl ausdrücke, *Mathematische Annalen* 100 (1928) 638–649. doi:10.1007/BF01448869.
- [23] M. Voigt, Decidable fragments of first-order logic and of first-order linear arithmetic with uninterpreted predicates, Ph.D. thesis, Universität des Saarlandes, Saarbrücken, Germany, 2019.
- [24] K. Gödel, Zum entscheidungsproblem des logischen funktionenkalküls, *Monatshefte für Mathematik und Physik* 40 (1933) 433–443.
- [25] B. Draben, W. D. Goldfarb, *The Decision Problem: Solvable Classes of Quantificational Formulas*, Addison-Wesley, 1979.
- [26] E. Kieronski, A uniform one-dimensional fragment with alternation of quantifiers, in: Proceedings of the Fourteenth International Symposium on Games, Automata, Logics, and Formal Verification, GandALF 2023, volume 390 of *EPTCS*, 2023, pp. 1–15. doi:10.4204/EPTCS.390.1.
- [27] M. Fürer, The computational complexity of the unconstrained limited domino problem (with implications for logical decision problems), in: *Logic and Machines: Decision Problems and Complexity*, Proceedings of the Symposium “Rekursive Kombinatorik” 1983, volume 171 of *Lecture Notes in Computer Science*, Springer, 1983, pp. 312–319. doi:10.1007/3-540-13331-3_48.
- [28] P. Erdős, On a problem in graph theory, *The Mathematical Gazette* 47 (1963) 220 – 223.