



Gary Mendese, MD
Stephen Moyer, DO
Robert Murgia, DO
Daniel O'Connor, MD
Todd Wechter, MD
Neera Nathan, MD

Molly Plovovich, MD
Alaina Iannazzi, PA-C
Rachel York, PA-C
Nathan Hand, PA-C
Michelle Ryder, PA-C
Heather Bigos, PA-C

Stephanie Grant, NP
Jane Tallent, NP
Christine Horrigan, NP
Lindsey Ramsey, RN
Brooke Comire, RN

LIQUID NITROGEN WOUND CARE

Treating an area of skin with liquid nitrogen freezes the skin, which kills the tissue in the top layer of skin. The skin then regenerates: if the frozen area is left undisturbed, a superficial treatment will not scar.

Here is what to expect from a treatment with liquid nitrogen, and how to care for the site.

WHAT TO EXPECT

- **Discomfort at the site:** If this was a light freeze on an area of skin other than the palm or sole, there will be some stinging for 10 or 15 minutes, and possibly some itching. If the area frozen was on the palm or the sole, it may be quite painful, and that pain will often last one or two days.
- **Blistering:** Sometimes a blister will form. This may be a clear blister or a blood blister.
- **Scabbing:** After two or three days, sometimes as long as a week, the area will feel like a scab -- DO NOT PICK AT IT! (Picking at it could cause a scar!)
- **Swelling:** Treatment to areas on the forehead, nose or eyelids can result in painless swelling of the eyelids, especially noted in the morning. This will subside in 2-7 days.
- **Discoloration:** The area frozen may turn darker brown or even black.
- **Peeling:** After 7-10 days the area will start to peel, or, if it forms a scab, dark parts of the scab will start to crumble off.
- **Redness:** After peeling, the area is likely to be slightly pink; this will fade after several days or weeks, depending on your skin type.

--> **STRICT SUN PROTECTION DURING THE HEALING PROCESS IS ESSENTIAL**

CARE OF THE SITE

- DO NOT PICK AT THIS AREA. Picking can damage the skin and cause possible scarring. It can also cause prolonged healing time.
- You may cover it with a bandaid if you desire, but this is not necessary unless the blister breaks.
- If it forms a large blister that gets in the way, you may pop the blister with a sterile needle or razor. If you do this, keep the area covered with antibiotic ointment and a bandaid.
- Call our office if discomfort in the area becomes more noticeable after 1 or 2 days.

Dermskinhealth.com

784 Central Avenue
Dover, NH 03820

2299 Woodbury Ave., Ste. 3
Newington, NH 03801

50 Michels Way, Suite 202
Londonderry, NH 03053

160 South River Road
Bedford, NH 03110

23 Centennial Drive
Peabody, MA 01960

P: 603-742-5556 F: 603-742-8668