## **Minoxidil Dosing in Children with Hair Loss**

(Minoxidil use is off label and should be used only in conjunction with a specialist. Doses shown here are maximum doses)

AGE	Type of Minoxidil	Dose
	(2 % or 5 %)	
0-2 years		Minoxidil not recommended by Dr. Donovan
3	2 %	3 drops once daily for 1 month and if tolerated then
		3 drops twice daily
4	2 %	4 drops once daily for 1 month and if tolerated then
		4 drops twice daily
5	2 %	5 drops once daily for 1 month and if tolerated then
		5 drops twice daily
6	2 %	6 drops once daily for 1 month and if tolerated then
		6 drops twice daily
7	2 %	7 drops once daily for 1 month and if tolerated then
		7 drops twice daily
8	2 % then 5 %	Start with 2 % and use 8 drops once daily for 1 month and if tolerated then
		8 drops twice daily. After 2 more months may use 5 % if tolerating well.
9	2 % then 5 %	Start with 2 % and use 9 drops once daily for 1 month and if tolerated then
		9 drops twice daily. After 2 more months may use 5 % if tolerating well.
10	2 % then 5 %	Start with 2 % and use 10 drops once daily for 1 month and if tolerated then
		10 drops twice daily. After 2 more months may use 5 % if tolerating well.
11	2 % then 5 %	Start with 2 % and use 11 drops once daily for 1 month and if tolerated then
		11 drops twice daily. After 2 more months may use 5 % if tolerating well.
12	2 % then 5 %	Start with 2 % and use 15 drops once daily for 2 months and if tolerated then
		15 drops twice daily. After 2 more months may use 5 % if tolerating well.
13	2 % then 5 %	Start with 2 % and use 15 drops once daily for 2 months and if tolerated then
		15 drops twice daily. After 2 more months may use 5 % if tolerating well.
14	2 % then 5 %	Start with 2 % and use 15 drops once daily for 2 months and if tolerated then
		15 drops twice daily. After 2 more months may use 5 % if tolerating well.
15	2 % then 5 %	Start with 2 % and use 16 drops once daily for 2 months and if tolerated then
		15 drops twice daily. After 2 more months may use 5 % if tolerating well.
16	2 % then 5 %	Start with 2 % and use 18 drops once daily for 2 months and if tolerated then
		18 drops twice daily. After 2 more months may use 5 % if tolerating well.
17	5%	20 drops once daily of 5 % for females. Males may use twice daily. May use $\frac{1}{2}$
		cap of the Minoxidil foam (once daily for females and twice daily for males).
		However, there is no need to switch if lotion is tolerated well.
18	5%	Standard adult dosing