



Jeff Donovan MD PhD FRCPC

Dermatologist, specializing in hair loss

750 West Broadway Suite 905 - Vancouver BC V5Z 1H8

Phone: 604.283.9299 Fax: 604.648.9003

Email: vancouveroffice@donovanmedical.com Web: www.donovanmedical.com

INFORMATION ON ORAL ITRACONAZOLE

What is Itraconazole?

- Itraconazole is an anti-fungal medication
- It belongs to a class of drugs known as the 'azole antifungals'
- It is known by generic names such as SporonoX

What is the dose or oral Itraconazole?

- A variety of doses may be prescribed.
- Commonly Dr. Donovan uses 200 mg daily for 1 week and then 200 mg for the first two days of every month thereafter for THREE additional months.
- 75-85 % of patients with seborrheic dermatitis have an improvement with this course of treatment.

What the side effects of Itraconazole?

- Nausea
- Vomiting
- Headaches
- Diarrhea
- Upset stomach, constipation, bloating
- Fatigue
- Dizziness, sensitivity to light
- Increased triglycerides in the blood
- Rarely liver disease
- Rarely a serious rash
- High blood pressure
- Rarely heart failure



Jeff Donovan MD PhD FRCPC

Dermatologist, specializing in hair loss

750 West Broadway Suite 905 - Vancouver BC V5Z 1H8

Phone: 604.283.9299 Fax: 604.648.9003

Email: vancouveroffice@donovanmedical.com Web: www.donovanmedical.com

INFORMATION ON ORAL ITRACONAZOLE – PAGE 2

How should I take oral Itraconazole?

- The medication is typically taken with food
- Antacids will decrease the absorption of this medication. If you use antacids, take the drug 2 hours before you take your antacid or 1 hour after.
- Dr. Donovan make recommend that you take the drug with an acidic beverage in some cases (like cola).

Can Itraconazole be used during pregnancy?

- No

What medications interact with Itraconazole and therefore may or may not be allowed together?

If you take ANY medications at all, please inform Dr. Donovan. Itraconazole tends to interact with many medications. These include:

1. Statins (cholesterol medications)
2. Pimozide
3. Warfarin
4. Glyberide
5. Clarithromycin, Erythromycin
6. Tacrolimus, Sirolimus, Cyclosporine
7. Proton pump inhibitors
8. Antacids
9. Calcium channel blockers
10. Alcohol and marijuana (can make patients more dizzy)

INFORMATION ON ORAL ITRACONAZOLE – PAGE 3

11. If you take any of these following medications, Dr. Donovan must be informed
- Isoniazid
 - Rifampins
 - Phenytoin
 - HIV medications (such as efavirenz, nevirapine)

Can I drink grapefruit juice if I use Itraconazole?

- No

Who should not use Itraconazole?

- Women who are pregnancy
- Individuals with heart failure
- Individuals who use medications listed above
- Individuals allergic to ‘azole’ medications
- Individuals with kidney problems

Itraconazole rarely causes serious problems. However, if you develop nausea or vomiting that continues for more than 24 hours or have loss of appetite, stomach pain, abdominal pain, rapid heartbeat, yellowing of the skin or eyes or dark urine you must stop the drug right away and advise Dr. Donovan immediately as well as seek medical attention.

Similarly, individuals with blurred vision, confusion, fever, chills, joint pain, pins and needles type feeling must stop the drug right away and advise Dr. Donovan immediately as well as seek medical attention.

ADDITIONAL MEDICATIONS WITH ITRACONAZOLE INTERACTIONS

| | |
|--|--|
| abiraterone | metronidazole |
| aliskiren | mifepristone |
| alpha blockers (e.g., alfuzosin, doxazosin, silodosin, tamsulosin) | mirtazapine |
| amiodarone | modafinil |
| antacids (e.g., aluminum hydroxide, calcium carbonate, magnesium hydroxide) | montelukast |
| anti-cancer medications (e.g., cabazitaxel, docetaxel, doxorubicin, etoposide, ifosfamide, irinotecan, trastuzumab, vincristine) | nadolol |
| anti-psychotics (e.g., chlorpromazine, clozapine, haloperidol, olanzapine, quetiapine, risperidone) | narcotic pain relievers (e.g., codeine, fentanyl, morphine, oxycodone) |
| apixaban | nateglinide |
| aprepitant | nefazodone |
| ariPIPrazole | nitrates (e.g., isosorbide dinitrate, isosorbide mononitrate) |
| “azole” antifungals (e.g., itraconazole, ketoconazole, voriconazole) | ondansetron |
| bedaquiline | oxycodone |
| benzodiazepines (e.g., alprazolam, chlordiazepoxide, clonazepam, clorazepate, diazepam, flurazepam, midazolam) | pentamidine |
| bicalutamide | perampanel |
| birth control pills | phenobarbital |
| boceprevir | phosphodiesterase-5 inhibitors (e.g., sildenafil, tadalafil, vardenafil) |
| bosentan | pimozide |
| buprenorphine | prasugrel |
| calcitriol | praziquantel |
| | procainamide |
| | progestins (e.g., dienogest, levonorgestrel, medroxyprogesterone, norethindrone) |
| | proton pump inhibitors (e.g., omeprazole, esomeprazole, lansoprazole) |

| | |
|---|--|
| calcium channel blockers (e.g., amlodipine, diltiazem, nifedipine, verapamil) | quinidine |
| carvedilol | quinine |
| cetirizine | quinolone antibiotics (e.g., ciprofloxacin, norfloxacin, ofloxacin) |
| chloroquine | repaglinide |
| cilostazol | rifabutin |
| cisapride | rifampin |
| colchicine | rifapentine |
| conivaptan | rivaroxaban |
| inhaled corticosteroids (e.g., budesonide, ciclesonide, fluticasone) | romidepsin |
| nasal corticosteroids (e.g., budesonide, fluticasone) | St. John's wort |
| oral corticosteroids (e.g., dexamethasone, hydrocortisone, prednisone) | salmeterol |
| cyclosporine | seizure medications (e.g., clobazam, ethosuximide, felbamate, levetiracetam, phenobarbital, phenytoin, primidone, topiramate, valproic acid, zonisamide) |
| dabigatran | selective serotonin reuptake inhibitors (SSRIs; e.g., citalopram, duloxetine, fluoxetine, paroxetine, sertraline) |
| dantrolene | serotonin antagonists (anti-emetic medications; e.g., granisetron, ondansetron, palonosetron) |
| dapsone | sirolimus |
| degarelix | sotalol |
| didanosine | "statin" anti-cholesterol medications (e.g., atorvastatin, lovastatin, simvastatin) |
| digoxin | sulcrafate |
| disopyramide | tacrolimus |
| dofetilide | telaprevir |
| domperidone | teniposide |
| dronedarone | terfenidine |
| eplerenone | |
| ergot alkaloids (e.g., dihydroergotamine, ergonovine, ergotamine) | |

| | |
|--|--|
| estrogens (e.g., conjugated estrogen, estradiol, ethinyl estradiol) | tetrabenazine |
| everolimus | tetracycline |
| flecainide | theophylline |
| flutamide | ticagrelor |
| fusidic acid | ticlopidine |
| "gliptin" diabetes medications (e.g., linagliptin, saxagliptin, sitagliptin) | tolterodine |
| grapefruit juice | tolvaptan |
| guanfacine | trabectedin |
| H2-receptor antagonists (e.g., cimetidine, famotidine, nizatidine, ranitidine) | tramadol |
| HIV non-nucleoside reverse transcriptase inhibitors (NNRTIs; e.g., delavirdine, efavirenz, etravirine, nevirapine) | trazodone |
| HIV protease inhibitors (e.g., atazanavir, indinavir, ritonavir, saquinavir) | tricyclic antidepressants (e.g., amitriptyline, clomipramine, desipramine, trimipramine) |
| isoniazid | trimethoprim |
| ivermectin | "triptan" migraine medications (e.g., eletriptan, sumatriptan) |
| lidocaine | tyrosine kinase inhibitors (e.g., bosutinib, dasatinib, imatinib, nilotinib) |
| loperamide | ulipristal |
| losartan | venlafaxine |
| macrolide antibiotics (e.g., clarithromycin, erythromycin) | warfarin |
| maraviroc | zolpidem |
| mefloquine | zopiclone |
| mestranol methadone | |

