

**TUSNEL DIABETIC- dextromethorphan, guaifenesin liquid**  
**LLORENS PHARMACEUTICALS INTERNATIONAL DIVISION**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**DRUG FACTS**

<b>Active Ingredients ( in each 5 mL tsp.)</b>	<b>Purpose</b>
Dextromethorphan HBr USP..... 10 mg .....	Cough Suppressant
Guaifenesin USP ..... 100 mg .....	Expectorant

**Purpose**

☐Cough Suppressant  
Expectorant☐

**Warnings**

**Do not use** if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or if 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

**Ask a doctor before use if you have**

- a cough that occurs with too much phlegm (mucus)
- cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Stop use and ask a doctor if** cough last more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. A persistent cough could b a sign of a serious condition.

**Keep out of the reach of children.** In case of overdose, get medical help or contact a Poison Center right away.

**If pregnant or breast-feeding,** ask a health professional before use.

**Directions**

Do not exceed more than 6 doses in any 24-hour period or as directed by a doctor.

Age	Dose
Adults and children 12 years of age and over	2 teaspoonfuls (10 mL) every 4 hours
Under 2 years of age	Do not use

**Uses**

Temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold  
Helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

**Inactive Ingredients:** anhydrous citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate dihydrate and sucralose.

**Questions or Comments? 1-866-595-5598**

**Distributed By: Llorens International Division, Miami FL 33165**

**Drug Facts**

**Active ingredient (in each 5 mL)** **Purpose**  
 Dextromethorphan HBr, USP 10 mg.....Cough Suppressant  
 Guaifenesin, USP 100 mg.....Expectorant

**Uses** • temporarily relieves cough due to minor throat and bronchial irritations as may occur with a cold •helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

**Warnings** Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use** if you have •cough that occurs with too much phlegm (mucus) •cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis, or emphysema

**Stop use and ask a doctor** if cough lasts for more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. A persistent cough may be a sign of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.



NDC 54859-505-16

# TUSNEL<sup>®</sup> DIABETIC

• NO SUGAR • NO SACCHARINE

• NO ALCOHOL • NO DYE

**Cough Suppressant/  
Expectorant**



**APPLE BANANA**

www.llorenspharm.com

**CONTENTS: 16 FL OZ (473 ML)**

**Drug Facts (Continued)**

**Directions:** Do not exceed more than 6 doses in any 24-hour period.

Age	Dose
Adults and children 12 years and over	2 teaspoonfuls (10 mL) every 4 hours
Children under 12 years	Ask a doctor

**Other information:** • store at controlled room temperature 15-30°C (59-86°F) • Do not use if foil over bottle opening is torn, broken, or missing. • Pharmacist- Preserve and dispense in tight light-resistant container with a child resistant cap as defined in the USP.

**Inactive ingredients** anhydrous citric acid, flavor, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate dihydrate, and sucralose.

**Questions or comments?** 1-866-595-5598

THIS IS A BULK CONTAINER NOT INTENDED FOR DISPENSING

Code#L-48 Rev.: 07/17

Lot#

Exp. Date



Manufactured by:  
**LLORENS**  
 PHARMACEUTICAL  
 INTERNATIONAL DIVISION  
 MIAMI, FL 33147  
 www.llorenspharm.com  
 MADE IN USA

## TUSNEL DIABETIC

dextromethorphan, guaifenesin liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:54859-505
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 5 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

### Inactive Ingredients

Ingredient Name	Strength
SUCRALOSE (UNII: 96K6UQ3ZD4)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	

**Product Characteristics**

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	GRAPE (grape flavor)	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54859-505-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2003	
2	NDC:54859-505-16	472 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2003	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	12/01/2003	

**Labeler** - LLORENS PHARMACEUTICALS INTERNATIONAL DIVISION (037342305)

Revised: 11/2017

LLORENS PHARMACEUTICALS INTERNATIONAL DIVISION