



HAL
open science

IL-6 -174 Genotype Associated with A.actinomycescomitans in Indians

Luigi Nibali, Isobel Madden, Fernando Franch Chillida, Lisa Heitz-Mayfield,
Peter M Brett, Nikos Donos

► **To cite this version:**

Luigi Nibali, Isobel Madden, Fernando Franch Chillida, Lisa Heitz-Mayfield, Peter M Brett, et al..
IL-6 -174 Genotype Associated with A.actinomycescomitans in Indians. Oral Diseases, 2010, 17
(2), pp.232. 10.1111/j.1601-0825.2010.01731.x . hal-00599893

HAL Id: hal-00599893

<https://hal.science/hal-00599893v1>

Submitted on 11 Jun 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ORAL DISEASES

IL-6 -174 Genotype Associated with A.actinomycetemcomitans in Indians

Journal:	<i>Oral Diseases</i>
Manuscript ID:	ODI-04-10-OM-1648.R1
Manuscript Type:	Original Manuscript
Date Submitted by the Author:	18-May-2010
Complete List of Authors:	Nibali, Luigi; UCL Eastman Dental Institute, Periodontology Madden, Isobel; 2 UHI School of Oral Health Science, Centre for Health Science Franch Chillida, Fernando; UCL Eastman Dental Institute, Periodontology Heitz-Mayfield, Lisa; The University of Western Australia, Centre for Rural and Remote Oral Health Brett, Peter; UCL Eastman Dental Institute, Periodontology Donos, Nikos; UCL Eastman Dental Institute, Periodontology
Keywords:	Genetics, Periodontitis, Interleukin 6, Bacteria, A.actinomycetemcomitans



1
2
3
4
5
6 **IL-6 -174 Genotype Associated with *A.actinomycetemcomitans* in**
7 **Indians**
8

9 Running title: IL6 genotypes and *A.actinomycetemcomitans*

10
11 Nibali L.¹, Madden I.^{1,2}, Franch-Chillida F.¹, Heitz-Mayfield L.J.A.³, Brett P.M.¹, Donos N.¹
12
13

14
15 ¹ Periodontology Unit and Division of Clinical Research, Eastman Dental Hospital, University
16 College London (UCL), London, UK.

17
18 ² UHI School of Oral Health Science, Centre for Health Science, Inverness, Scotland

19
20 ³ The University of Western Australia. Centre for Rural and Remote Oral Health. Crawley, Western
21 Australia, Australia

22 Running title: IL-6 polymorphisms and periodontal bacteria

23
24 Keywords: Periodontitis, Interleukin-6, Genetic, Bacteria, *A.actinomycetemcomitans*
25

26 Date of submission: 22/04/2010
27
28
29

30 Corresponding author:

31 Luigi Nibali

32 Periodontology Unit

33 Eastman Dental Institute

34 256 Gray's Inn Road

35 London WC1X 8LD

36 Tel:00442079151086

37 Fax 00442079151137

38 l.nibali@eastman.ucl.ac.uk
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

AIM: Genetic factors have recently been associated with presence of *Aggregatibacter actinomycetemcomitans* subgingivally in populations living in industrialised countries. The aim of this study was to analyze associations between Interleukin-6 (*IL6*) single nucleotide polymorphisms and presence and levels of *A.actinomycetemcomitans* and other subgingival microbes in a rural Indian population.

SUBJECTS AND METHODS: 251 individuals from a rural village in India with a periodontal phenotype ranging from healthy to severe periodontitis were included. Checkerboard DNA-DNA analysis was performed to detect 40 periodontal taxa in subgingival plaque samples. Genomic DNA was extracted to genotype five polymorphisms in the *IL6* promoter region.

RESULTS: The *IL6* -174 GG genotype was associated with high (above median) counts of *A.actinomycetemcomitans* (both in all subjects and in periodontally healthy only) and with presence and counts of *Capnocytophaga sputigena*. Differences in detection of several other bacteria were noted between periodontitis and healthy subjects.

CONCLUSION: These findings support the influence of genetic factors on the subgingival microbiota.

INTRODUCTION

The oral cavity may contain as many as 19,000 bacterial phylotypes (Keijser et al., 2008). We have recently introduced the concept of periodontal infectogenomics, which studies the effect host genetic factors have on the composition of the subgingival microbiota of each individual. This is based on the assumption that genetic factors in the host seem to play a major role in deciding which bacteria (commensal and pathogenic) are able to colonise the host (Nibali et al., 2009a).

Most studies on subgingival microbes with a putative influence on periodontally-relevant immune responses have been carried out in industrialised communities with routine access to dental care (Hardie et al., 1975; Papapanou et al., 1993; Umeda et al., 1998). Analysis of rural populations not subjected to dental care allows the study of possible effects of host genetic factors in the composition of the subgingival microbiota in almost pristine conditions, undisturbed by routine tooth scaling and use of antibiotics.

We have recently shown an association between *IL6* genetic factors and clinical status in non smokers living in a rural Indian village (Franch et al., 2010). The main aim of this analysis was to study the association between *IL6* genetic variants and the presence and levels (counts) of *A.actinomycetemcomitans* in subgingival plaque samples from the same population. Secondary aims were to explore the association between *IL6* genetic factors and the detection and counts of other bacteria, and the association between periodontal status and the detection and levels of other bacteria.

MATERIALS AND METHODS

Subject Selection

The population reported here has also been described elsewhere (Franch et al., 2010). The study took place in the village of Dokur, Andhra Pradesh, where the Institute for Rural Health Studies (IRHS), a local Non-Governmental Organization (NGO), runs a rural clinic. All adult inhabitants between the ages of 18 and 70 were invited to attend a screening visit by means of local advertising within the village. A group of 500 villagers attended and were initially screened based on the following exclusion criteria: i) presence of serious systemic conditions (malaria, tuberculosis or cardiovascular disease), ii) presence of infectious or parasitic diseases, iii) pregnancy or breastfeeding, iv) antibiotic or anti-inflammatory medications within the last 3 months, v) fewer than 16 standing teeth.

Two hundred and fifty one suitable subjects (10% of the total village population) were identified and included in the study.

Ethical Considerations

The study was conducted in full accordance with ethics principles, including the World Medical Association Declaration of Helsinki (version 2002). Ethical approval for the study was obtained from the Medical Board of IRHS and conformed to the Indian Council of Medical Research Guidelines on Medical Ethics 2000 (Indian Council of Medical Research, Ansari Nager, New Delhi). All subjects were native to the village and voluntary informed consent to participation in the study was obtained from each by means of a written consent form, translated into Telugu and also read aloud to each subject by a fluent Telugu speaker. Demographic data and self-reported data on

1
2 education, social status, type of toothbrushing, alcohol consumption and smoking habit were
3
4 recorded (Franch et al., 2010).

7 8 **Clinical Data Collection**

9
10 Full mouth periodontal examination of probing pocket depth (PPD) and clinical attachment level
11 (CAL) was carried out for each subject with an EN-15 probe on 6 sites per tooth. The clinical
12 examination was performed by one single operator (IM), calibrated to an exact kappa level of 0.85
13 for PPD. Patients were divided into 2 groups (periodontitis/ healthy) based on their periodontal
14 status. Both definitions suggested by the European Workshop on Periodontitis (EWP) (Tonetti and
15 Claffey, 2005) were used:
16
17
18
19
20

- 21 • Presence of proximal attachment loss of $\geq 3\text{mm}$ in ≥ 2 non-adjacent teeth (EWP1)
 - 22 • Presence of proximal attachment loss of $\geq 5\text{mm}$ in $\geq 30\%$ of teeth present (EWP2)
- 23
24
25
26
27

28 29 **Microbiological analysis**

30
31 Dental plaque samples were taken from the mesio-buccal aspects of the upper right and lower left
32 first molar teeth, unless a deeper pocket had been found at another site, in which case this site was
33 used in preference. Subgingival plaque was removed from each site isolated with cotton rolls,
34 following removal of supragingival plaque, using sterile 13/14 double-ended Columbia curettes.
35
36 The scraped mass was transferred from the curette to an Eppendorf tube containing 150 μl of sterile
37 TE buffer (10mM Tris HCl, 1.0mM EDTA, pH 7.6) and subsequently transferred to the University
38 of Berne. The presence of 40 periodontal pathogens was evaluated using checkerboard DNA-DNA
39
40
41
42
43
44 hybridisation technique (Socransky et al., 1994).
45
46

47 48 **Genetic analysis**

49
50
51
52
53
54
55
56
57
58
59
60

1
2 Samples of buccal mucosal cells were taken from all subjects using wire cytology brushes (Medical
3 Wire and Equipment Company, UK) (Franch et al., 2010). Genomic DNA was extracted from
4 these samples and blindly genotyped for polymorphisms at positions -174 (CCTTTAGCAT[C-
5 G]GCAAGAC, rs 1800795), -572 (CAACAGCC[C-G]CTCACAG, rs 1800796), -1363
6 (CACTGTTTTATC[G-T]GATCTTG, rs 2069827) and -6106 (TCTCTACA[A-
7 T]TAAGAAATAC) and -1480 (ACCGTCTCT[C-G]TGTTTAC) in the *IL6* gene by real time PCR
8 as previously described (Nibali et al., 2008b).
9
10
11
12
13
14
15
16
17

18 **Statistical Analysis**

19
20 Comparisons of continuous and categorical data between groups (periodontitis and healthy) were
21 analysed with ANOVA and Chi-square test, respectively. The alpha value was set at 0.05.
22
23
24
25

26 The approach to evaluate the association between genetic factors, clinical diagnosis and
27 microbiological outcomes was performed using the SPSS 12.0 package. The main outcome was
28 association between *IL6* -174 GG genotype and the presence (at detection threshold 1×10^4) and
29 levels (total counts) of *A.actinomycetemcomitans*. An exploratory analysis was performed to
30 investigate associations between all studied *IL6* genetic variables (and clinical diagnosis) and other
31 taxa and bacterial complexes. The α value for this exploratory analysis was lowered to 0.01 in order
32 to adjust for multiple testing. Microbiological data were available from two sites for all subjects.
33 Data on detection or not of all 40 studied taxa are reported. For analysis of bacterial counts,
34 negative (no detection) reactions were given an arbitrary value of 0.99×10^4 , just below detection
35 limit. The mean individual count for each species was obtained as the averaged between the 2 sites
36 in each patient. A priori, median values for each bacterium were calculated and two clusters of
37 patients for each bacterium were identified: high (above median) and low (below median) counts.
38 Therefore, the microbiological outcomes were: i) detection of each bacterium (at 1×10^4 detection
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2 value); ii) high or low levels (counts) of each bacterium (both categorical data) and iii) absolute
3 counts of each bacterium (continuous data). Subgroup analyses were performed in subjects divided
4 by their periodontal status (healthy/periodontitis). Multiple logistic regression analysis adjusting for
5 confounders (age, smoking, gender and type of toothbrushing) was performed to investigate these
6 associations (categorical variables). Linear logistic regression adjusting for confounders (age,
7 smoking, gender and type of toothbrushing) was used to detect differences in bacterial counts
8 (continuous variables) between genotypes and by clinical diagnosis.

Deleted: Mann-Whitney test

Deleted: as previously described
(continuous variables) (Socransky et al.,
2000)

18 RESULTS

19
20 A total of 251 subjects took part in the study. Their demographic and clinical characteristics have
21 been reported elsewhere (Franch et al., 2010) (table 1). Forty-two per cent and 19% of subjects
22 were diagnosed with periodontitis, according to EWP1 and EWP2 definitions respectively (Tonetti
23 and Claffey, 2005). Thirteen subjects resulted *IL6* -174 C homozygous, 46 CG and 185 GG, while 7
24 could not be scored. The distribution of genetic polymorphisms has been reported elsewhere and
25 satisfied the chi-squared analysis for Hardy-Weinberg equilibrium on all subjects (Franch et al.,
26 2010).

27
28 The detection of the 40 studied taxa ranged from 98% of subjects for *Prevotella intermedia* to 51%
29 for *Capnocytophaga sputigena*. In particular, 93% of subjects were positive for *A.*
30 *actinomycetemcomitans*, 93% for *Porphyromonas gingivalis*, 90% for *Tannerella forsythia*, 70%
31 for *Treponema denticola* and 60% for *Campylobacter rectus*. The mean counts ranged from $11.2 \times$
32 10^5 for *Neisseria mucosa* to 0.2×10^5 for *Propionibacterium acnes*. Average counts for *A.*
33 *actinomycetemcomitans*, *P. gingivalis* and *T. forsythia* for all subjects were respectively 5.1×10^5 ,
34 12.0×10^5 and 6.4×10^5 .

Association between *IL6* genetic factors and bacterial detection

IL6 -174 GG genotype subjects exhibited high (above median) levels of *A.actinomycetemcomitans* compared to CC and CG individuals, irrespective of clinical diagnosis (adjusted $p=0.043$, O.R. = 1.89, 95% C.I. = 1.02- 3.50) (figure 1). This association was confirmed among the subgroup of non-periodontitis individuals (according to EWP2) (adjusted $p=0.037$, O.R. = 2.11, 95% C.I. = 1.0-4.2).

Deleted: periodontally healthy

Among other bacteria, *IL6* -174 GG genotype subjects showed increased detection, and high (above median) levels of *C.sputigena* (adjusted $p= 0.003$, O.R. = 2.60, 95% C.I. = 1.38- 4.90 and $p=0.009$, O.R. = 2.29, 95% C.I. = 1.23- 4.26 respectively). No statistically significant differences in percentage of bacterial complexes (Socransky *et al.*, 1998) were found according to *IL6* -174 genotypes (figure 2).

Associations between clinical diagnosis and bacterial detection

Figure 3 shows the frequency of detection of the 40 studied bacteria in periodontitis and healthy individuals according to EWP1 (Tonetti and Claffey, 2005). Statistically significant associations were found between diagnosis of periodontitis (EWP1) and presence of *N.mucosa* ($p=0.008$) and *Capnocytophaga ochracea* ($p=0.001$), low (below-median) counts of *N.mucosa* ($p<0.001$), and high (above-median) counts of *Parvimonas micra* (previously *Peptostreptococcus micros*) ($p=0.002$).

Deleted: 8

Periodontitis patients also exhibited lower absolute counts of *N.mucosa* ($p=0.003$).

Deleted:), *C.ochracea* ($p<0.001$), and *Selenomonas noxia* ($p<0.001$)

EWP2 diagnosis of periodontitis was associated with absence (no detection) of *C.ochracea* ($p=0.005$) and *S.noxia* ($p<0.001$) and higher absolute counts of *P.micra* ($p=0.025$). When bacteria were grouped in complexes, periodontal status (EWP1) was associated with percentage of bacteria undefined in complexes ('others') ($p=0.012$).

Deleted: os

Deleted: <0.001

Deleted: <0.001 and $p=0.001$ for EWP1 and EWP2 respectively

Deleted: EWP2 definition was also associated with percentage of green complex bacteria ($p=0.002$).

Deleted: ¶
-----Page Break-----

DISCUSSION

We have previously shown an association between *IL6* -174 genotypes and presence of periodontitis in non-smokers living in a rural village in India (Franch et al., 2010). The present study describes associations between *IL6* -174 GG genotype and presence of *A.actinomycescomitans* in periodontal pockets in these subjects (with a periodontal phenotype ranging from healthy to severe periodontitis), confirming association previously observed in subjects living in the United Kingdom (Nibali et al., 2007, Nibali et al., 2008a). Among these Indian villagers living in Andhra Pradesh, *A.actinomycescomitans* was almost ubiquitous subgingivally, as it was harboured by 93% of participating individuals. However, *IL6* -174 GG individuals had higher counts of this bacterium.

Deleted: subgingival

The *IL6* -174 GG has been associated with presence of periodontitis (Trevilatto et al., 2003, Nibali et al., 2009b). The study reported here confirms the previously reported association between *IL6* genotypes and *A.actinomycescomitans* (Nibali et al., 2007, Nibali et al., 2008a), despite being conducted on a different population of different ethnicity and socio-economic status and not subjected to routine dental care. Furthermore, the microbiological analysis consisted of a checkerboard DNA-DNA hybridisation analysis (Socransky et al., 1994), as opposed to previously reported culture (Nibali et al., 2007) and PCR (Nibali et al., 2008a) analyses. This association was seen in all subjects, independent from clinical diagnosis. In addition to this, this paper for the first time shows the association between *IL6* genotypes and *A.actinomycescomitans* also in non-periodontitis individuals (according to EWP2 definition). As previously hypothesized, this association may be due to the increased local inflammatory response of -174 GG individuals (Fishman et al., 1998, Bennermo et al., 2004), which favours the overgrowth of components of the microbiota that grow well in inflamed areas, such as *A.actinomycescomitans*. This study brings preliminary evidence that other bacteria belonging to the green complex, in particular *C.sputigena*,

Deleted: periodontally healthy

Deleted: subgingival

1
2 may also be associated with increased counts in periodontal pockets of *IL6* -174 GG individuals. In
3
4 other terms, the effect of genetic factors may not be limited to the presence or levels of one or two
5
6 potentially pathogenic bacteria, but may extend to bacterial complexes or in general to the whole
7
8 composition of the subgingival biofilm. This is in agreement with the concept of infectogenomics,
9
10 suggesting that genetic factors in the host seem to play a major role in deciding which bacteria
11
12 (commensal and pathogenic) are able to colonise the host (Kellam and Weiss, 2006, Nibali et al.,
13
14 2009a).

15
16
17
18 The microbiological analysis of the current subject group showed a population presenting with a
19
20 complex oral flora, with a breakdown in bacterial complexes quite different from that observed in
21
22 industrialized populations (Haffajee et al., 1998). Individuals diagnosed with periodontitis had a
23
24 tendency towards higher percentage of green complex bacteria, and lower percentage of bacteria not
25
26 defined in complexes ('others'). Overall, there was a predominance of bacteria belonging to the
27
28 orange complex, which have been associated with established periodontitis in industrialised
29
30 populations, while no significant differences in percentage of orange or red complex bacteria were
31
32 noted between periodontitis and non-periodontitis individuals. This was consistent with studies in
33
34 other untreated populations, which have shown ubiquitous presence of certain bacterial species and
35
36 a tendency towards a stable, mixed microflora (Timmerman et al., 1998, Dowsett et al., 2002,
37
38 Papapanou et al., 2002, Colombo et al., 2002). *N.mucosa*, a Gram negative commensal bacterium,
39
40 was the bacterium found in highest counts, followed by periodontopathogenic bacteria *P.gingivalis*,
41
42 *T.forsythia* and *A.actinomycetemcomitans*. This shows a considerable shift from the bacterial profile
43
44 shown in a study with checkerboard DNA-DNA hybridization on a large United States population
45
46 (Haffajee et al., 2006). The difference in prevalence and counts of *A.actinomycetemcomitans*
47
48 between the study described here and the one by Haffajee and coworkers is particularly striking. It
49
50 needs to be highlighted that only two of the deepest pockets per subject were sampled in the present

1
2 study. Therefore, the average bacterial counts in these cases in not representative of the whole
3
4 subgingival microbiota. However, even when the sampled site was used as unit for analysis, no
5
6 differences were detected between bacterial complexes between deep (>6mm) and shallow (<4mm)
7
8 pockets (data not presented).
9

10
11
12
13 Among the drawbacks of this study, we have to acknowledge the limitation of having only two
14
15 plaque samples sites per patient, the risk for false positive results associated with DNA-DNA
16
17 checkerboard analysis (van Steenberg et al., 1996) and the difficulty of defining periodontitis in
18
19 this population (Preshaw 2009, Franch et al. 2010). The results reported here show an association
20
21 between *IL6* -174 polymorphism and *A.actinomycetemcomitans* in subjects living in a rural Indian
22
23 village, bringing further evidence, consistent with previous studies, of the importance of
24
25 infectogenomics in periodontitis (Nibali et al., 2009a). Further studies should analyze the possible
26
27 effect of these and other genetic factors (in both periodontitis and healthy populations) on a wider
28
29 range of components of the oral microbiota, on different serotypes and clones of
30
31 *A.actinomycetemcomitans*, and should investigate possible functional mechanisms for the observed
32
33 associations.
34
35
36
37
38
39

Deleted: (which might to some extent affect the overall bacterial counts)

40 ACKNOWLEDGEMENTS

41 This work was undertaken at UCLH/UCL who received a proportion of funding from the
42 Department of Health's NIHR Biomedical Research Centres funding scheme. This study was also
43 supported by the Periodontal Research Fund of the Eastman Dental Institute. The help of Professor
44 Gareth Griffiths, Professor Hubert Newman, Dr Bidinger and the Institute for Rural Health Studies,
45 Hyderabad, Andhra Pradesh, are gratefully acknowledged.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

- 1
2
3
4
5
6 Bennermo M, Held C, Stemme S, Ericsson CG, Silveira A, Green F et al. (2004). Genetic
7 predisposition of the interleukin-6 response to inflammation: Implications for a variety of major
8 diseases? *Clinical Chemistry* **50**: 2136-2140.
- 9
10 Fishman D, Faulds G, Jeffery R, Mohammed-Ali V, Humphries S, Woo P (1998). An interleukin-6
11 promoter polymorphism that influences gene transcription, plasma IL-6 levels and is associated
12 with systemic-onset juvenile chronic arthritis. *European Cytokine Network* **9**: 364.
- 13
14 Kellam P, Weiss RA (2006). Infectogenomics: Insights from the host genome into infectious
15 diseases. *Cell* **124**: 695-697.
- 16
17 Keijsers B, Zaura E, Huse SM, van der Vossen JMBM, Schuren F, Montijn RC et al. (2008).
18 Pyrosequencing analysis of the Oral Microflora of healthy adults. *Journal of Dental Research* **87**:
19 1016-1020.
- 20
21 Colombo APV, Teles RP, Torres MC, Souto R, Rosalem W, Mendes MCS et al. (2002).
22 Subgingival microbiota of Brazilian subjects with untreated chronic periodontitis. *Journal of*
23 *Periodontology* **73**: 360-369.
- 24
25 Dowsett SA, Kowolik MJ, Archila LA, Eckert GJ, LeBlanc DJ (2002). Subgingival microbiota of
26 indigenous Indians of Central America. *Journal of Clinical Periodontology* **29**:159-167.
- 27
28 Franch Chillida F, Nibali L, Madden I, Donos N, Brett PM (2009). Association between
29 Interleukin-6 polymorphisms and periodontitis in a rural Indian population. *Journal of Clinical*
30 *Periodontology* **37**: 137-144.
- 31
32 Hardie JM, Bowden GH (1975). Bacterial-Flora of Dental Plaque. *British Medical Bulletin* **31**:131-
33 136.
- 34
35 Haffajee AD, Cugini MA, Tanner A, Pollack RP, Smith C, Kent RL et al. (1998). Subgingival
36 microbiota in healthy, well-maintained elder and periodontitis subjects. *Journal of Clinical*
37 *Periodontology* **25**:346-353.
- 38
39 Haffajee AD, Teles RP, Socransky SS (2006). Association of *Eubacterium nodatum* and *Treponema*
40 *denticola* with human periodontitis lesions. *Oral Microbiology and Immunology* **21**:269-282.
- 41
42 Nibali L, Ready DR, Parkar M, Brett PM, Wilson M, Tonetti MS et al. (2007). Gene
43 polymorphisms and the prevalence of key periodontal pathogens. *Journal of Dental Research*
44 **86**:416-420.
- 45
46 Nibali L, Tonetti MS, Ready D, Parkar M, Brett PM, Donos N et al. (2008a). Interleukin-6
47 polymorphisms are associated with pathogenic bacteria in subjects with periodontitis. *Journal of*
48 *Periodontology* **79**:677-683.
- 49
50 Nibali L, Griffiths GS, Donos N, Parkar M, D'Aiuto F, Tonetti MS et al. (2008b). Association
51 between interleukin-6 promoter haplotypes and aggressive periodontitis. *Journal of Clinical*
52 *Periodontology* **35**:193-198.
- 53
54
55
56
57
58
59
60

- 1
2 Nibali L, Donos N, Henderson B (2009a). Periodontal infectogenomics. *Journal of Medical*
3 *Microbiology* **58**:1269-1274.
4
5 Nibali L, D'Aiuto F, Donos N, Griffiths GS, Parkar M, Tonetti MS et al. (2009b). Association
6 between periodontitis and common variants in the promoter of the interleukin-6 gene. *Cytokine*
7 **45**:50-54.
8
9 Papapanou PN, Sellen A, Wennstrom JL, Dahlen G (1993). An Analysis of the Subgingival
10 Microflora in Randomly Selected Subjects. *Oral Microbiology and Immunology* **8**:24-29.
11
12 Papapanou PN, Teanpaisan R, Obiechina NS, Pithpornchaiyakul W, Pongpaisal S, Pisuthanakan S
13 et al. (2002). Periodontal microbiota and clinical periodontal status in a rural sample in southern
14 Thailand. *European Journal of Oral Sciences* **110**:345-352.
15
16 Preshaw PM (2009). Definitions of periodontal disease in research. *Journal of Clinical*
17 *Periodontology* **36**:1-2.
18
19 Socransky S, Smith AC, Martin L, Paster BJ, Dewhurst FE, Levin AE (1994). Checkerboard
20 DNA-DNA hybridisation. *Biotechniques* **17**: 788 – 792.
21
22 Socransky SS, Haffajee AD, Cugini MA, Smith C, Kent RL (1998). Microbial complexes in
23 subgingival plaque. *Journal of Clinical Periodontology* **25**:134-144.
24
25 Socransky SS, Haffajee AD, Smith C, Duff GW (2000). Microbiological parameters associated
26 with IL-1 gene polymorphisms in periodontitis patients. *Journal of Clinical Periodontology* **27**:810-
27 818.
28
29 Timmerman MF, Van der Weijden GA, Abbas F, Arief EM, Armand S, Winkel EG et al. (2000).
30 Untreated periodontal disease in Indonesian adolescents - Longitudinal clinical data and prospective
31 clinical and microbiological risk assessment. *Journal of Clinical Periodontology* **27**:932-942.
32
33 Tonetti MS, Claffey N. European Workshop in Periodontology group C. (2003) Advances in the
34 progression of periodontitis and proposal of definitions of a periodontitis case and disease
35 progression for use in risk factor research. Group C consensus report of the 5th European Workshop
36 in Periodontology. *Journal of Clinical Periodontology* **32** Suppl.: 210-3
37
38 vanSteenbergen TJM, Timmerman MF, Mikx FHM, deQuincey G, vanderWeijden GA,
39 vanderVelden U et al. (1996). Discrepancy between culture and DNA probe analysis for the
40 detection of periodontal bacteria. *Journal of Clinical Periodontology* **23**:955-959.
41
42 Umeda M, Chen C, Bakker I, Contreras A, Morrison JL, Slots J (1998). Risk indicators for
43 harboring periodontal pathogens. *Journal of Periodontology* **69**:1111-1118.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

FIGURE LEGENDS

Figure 1. Representation of number of subjects with *A.actinomycetecomitans* above or below median according to their IL-6 -174 genotypes. GG vs CG/CC p=0.043, O.R.= 1.89, 95% C.I.=1.02- 3.50.

Figure 2. Pie charts with distribution of bacterial complexes between IL6 -174 GG and CG/CC individuals. % DNA probe counts for each bacterium was determined at each site, and averaged per patient. The % of the DNA probe count for each species in each complex (according to Socransky *et al.*, 1998) was summed and the proportions of each complex were calculated.

Figure 3. Counts of 40 studied taxa divided in complexes in healthy and periodontitis subjects (according to EWP1 definition).

* *P.micra*: p<0.01 for differences in high (above median) counts between healthy and periodontitis

Deleted: os

** *C.ochracea*: p<0.01 for differences in detection and high (above median) counts between healthy and periodontitis

Deleted: * * *S.noxia*: p<0.01 for differences in total counts between healthy and periodontitis

Deleted: *


*** *N.mucosa*: p<0.01 for differences in detection, high (above median) counts and absolute counts between healthy and periodontitis.

Deleted: *

		All subjects (n=251)	EWP1		EWP2	
			Periodontitis (n=105)	Non- periodontitis (n=146)	Periodontitis (n=47)	Non-periodontitis (n=204)
Age			45.8 ± 10.5	37.5± 11.1	49.6 ± 10.3	39.0± 10.9
Gender (female)		162 (64.5%)	62 (59.0%)	100 (68.5%)	28 (59.6%)	134 (65.7%)
Smoking	Current	44 (17.5%)	26 (24.8%)	18 (12.3%)	33(70.2%) 14 (29.8%)	174(85.3%)
	Never	207 (82.5%)	79 (75.2%)	128(87.7%)		30(14.7%)
Clinical data	Number of sites PPD 4-6mm	16.1 ± 22.1	38.5 ± 17.4	0.1± 0.4	45.5± 15.9	8.7±15.6
	Number of sites PPD >6mm	3.3 ± 8.1	7.9 ± 11.1	0.0± 0.2	15.1± 13.1	0.6±1.8
	Number of sites CAL >4mm	16.6 ± 22.1	39.5± 16.3	0.1± 0.6	47.5± 15.5	9.5±16.6
	Number of sites CAL >6mm	6.1± 14.0	14.5± 18.5	0.0± 0.4	27.5± 21.1	1.1± 2.7


Table 1. Demographic and clinical characteristics of the participants, as divided by their periodontal status.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49




1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49



IL6 -174 CC (n=13)




IL6 -174 CG (n=46)



IL6 -174 GG (n=185)





HEALTHY (n=146)

PERIODONTITIS (n=105)