

AperTO - Archivio Istituzionale Open Access dell'Università di Torino

**The impact of the hydroxyl radical photochemical sources on the rivastigmine drug transformation in mimic and natural waters**

**This is the author's manuscript**

*Original Citation:*

*Availability:*

This version is available <http://hdl.handle.net/2318/1701128> since 2019-05-07T17:46:10Z

*Published version:*

DOI:10.1016/j.watres.2013.06.024

*Terms of use:*

Open Access

Anyone can freely access the full text of works made available as "Open Access". Works made available under a Creative Commons license can be used according to the terms and conditions of said license. Use of all other works requires consent of the right holder (author or publisher) if not exempted from copyright protection by the applicable law.

(Article begins on next page)

# The impact of the hydroxyl radical photochemical sources on the rivastigmine drug transformation in mimic and natural waters

Monica Passananti<sup>a,b,c</sup>, Fabio Temussi<sup>c\*</sup>, Maria Rosaria Iesce<sup>c</sup>, Gilles Mailhot<sup>a,b</sup>,  
Marcello Brigante<sup>a,b\*</sup>

<sup>a</sup> Clermont Université, Université Blaise Pascal, Institut de Chimie de Clermont-Ferrand (ICCF)- ENSCCF, BP 10448, F-63000 Clermont-Ferrand, France.

<sup>b</sup> CNRS, UMR 6296, ICCF, F-63171 Aubière, France

<sup>c</sup> UdR Napoli 4 INCA, Dipartimento di Scienze Chimiche, Università di Napoli Federico II, Complesso Universitario Monte S. Angelo, Via Cintia, I-80126 Napoli, Italy.

\* Address correspondence to either author. E-mail: [fabio.temussi@unina.it](mailto:fabio.temussi@unina.it); [marcello.brigante@univ-bpclermont.fr](mailto:marcello.brigante@univ-bpclermont.fr)

## Abstract

In this paper we investigated the degradation of the rivastigmine drug induced by hydroxyl radical in synthetic and natural waters focusing on both reactivity and photoproduct identification. The hydroxyl radical formation rate was quantified by using terephthalic acid as trapping molecule and it was related with the rivastigmine degradation rate. The second order rate constant between hydroxyl radical and rivastigmine was estimated to be  $\sim 5.8 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ . Irradiation of rivastigmine in three natural waters (rain, lake and river) and comparison with degradation rates observed in synthetic solutions using nitrite, nitrate and hydrogen peroxide suggest that, in addition to hydroxyl radical, also nitroderived radicals ( $\text{NO}/\text{NO}_2$ ) are responsible for the pollutant degradation in natural media. In fact, the evaluated degradation rates in three natural waters are greatly higher than those estimated considering only the reactivity with photogenerated hydroxyl radical. Using nitrites and nitrates as photochemical  $\bullet\text{OH}$  source, the rivastigmine degradation cannot be described considering

28 only the hydroxyl radical reactivity suggesting that NO and NO<sub>2</sub> radicals could play a key  
29 role during indirect degradation. Moreover main degradation products have been identified by  
30 means of HPLC-MS. Hydroxylation of the aromatic ring as well as carbamate and amino  
31 chain oxidation were suggested as main reaction mechanisms, but also nitroderived  
32 compounds were characterized.

33 Finally polychromatic irradiations of three rivastigmine doped natural waters (rain, river and  
34 lake) underlined the role of the indirect degradation that needs to be considered when direct  
35 degradation of selected pollutants is negligible under environmental-like conditions.

36

37 **Keywords:** Photoinduced degradation; environmental fate; radical-induced transformation;  
38 photooxidation; hydroxyl radical

39

## 40 **1. Introduction**

41 In recent years, the environmental pollution by emerging pollutants has become a major issue  
42 and it is now well established that human pharmaceuticals effects might potentially play a  
43 major role in other (non-mammalian) eco-organisms once reached the natural media (deep  
44 and surface waters, soils ,...) (Khetan and Collins 2007). On the other hand, the impact and  
45 fate of xenobiotics need to be investigated in order to asses the environmental potential risk.  
46 Furthermore, the investigation of the environmental fate of different kinds of pharmaceuticals  
47 is one of the main goals of recent European projects (Pharmas and Cytothreat). As stated in  
48 the summary of the Pharmas project a classification system needs to be developed with the  
49 intention of enabling all EU citizens to make their own informed decisions about the risk  
50 posed by human pharmaceuticals to their health and to the health of the environment. The  
51 results of this investigation will able EU regulators and policy makers to make better

52 informed decisions on the issue of pharmaceuticals in the environment (RIFERIMENTO\_  
53 link? <http://www.pharmas-eu.org/>).

54 In addition to priority pollutants many organic compounds used for human and veterinary  
55 health, known as pharmaceuticals and personal care products (PPCPs), are regularly  
56 introduced into the environment (McClellan and Halden 2010, Richardson and Ternes 2005,  
57 Yu and Chu 2009). Many of these compounds, after application, are excreted by urine and  
58 feces and are transported into sewage (Farré et al. 2008, Kümmerer et al. 1997). The sewage  
59 treatment plants (STPs) are often unable to completely remove PPCPs that in this way enter  
60 the environment, where they are frequently found (Andreozzi et al. 2003, Ellis 2006, Murray  
61 et al. 2010). In the environment the pollutants may undergo biotic and/or abiotic  
62 transformations (Cermola et al. 2007, Della Greca et al. 2008, DellaGreca et al. 2006, Jjemba  
63 2006). In the case of surface waters, photochemical processes are the most important among  
64 abiotic transformations. In such medium the indirect photolysis involving oxidant species  
65 (singlet oxygen, hydroxyl radical, photoexcited organic matter) could be relevant for the  
66 complete transformation of such compounds. The transformation products may be more  
67 persistent and exhibit higher toxicity than parent compounds (DellaGreca et al. 2004). Drugs  
68 are compounds made to have a biological activity and in the environment, even at small  
69 concentration, they can interact with aquatic or non aquatic organisms with effects, often  
70 unknown, on the ecosystem (Daughton and Ternes 1999, Halling-Sørensen et al. 1998,  
71 Jørgensen and Halling-Sørensen 2000, Vaal et al. 1997). PPCPs include many classes of  
72 heterogeneous compounds. More than 3000 pharmaceutical ingredients are registered  
73 (Richardson et al. 2005, Ternes et al. 2004) and it is hard to predict the environmental fate and  
74 the critical concentration for most of these compounds.

75 Many studies in the literature are about direct photodegradation of pollutants in organic  
76 solvents and in water, identification of their photoproducts and the related formation

77 mechanisms (Burrows et al. 2002, de Bertrand and Barceló 1991, Doll and Frimmel 2003,  
78 Sanz-Asensio et al. 1999). In order to predict the environmental fate of a pollutant it is  
79 essential to study indirect photolysis as well, which is often the most important process. In  
80 fact, in natural waters the pollutants may react with oxidant species more rapidly than with  
81 direct light. Only few studies determine the values of second order kinetic constants with  $\bullet\text{OH}$   
82 (Charbouillot et al. 2012, Razavi et al. 2011), one of the most important oxidants in natural  
83 waters.

84 Rivastigmine (Riv) is a drug used to treat patients with Alzheimer's disease and dementia  
85 associated with Parkinson's disease (Mustazza et al. 2002). Rivastigmine hydrogen tartrate is  
86 one of the few drugs approved in 2000 to treat Alzheimer's disease. This drug is a  
87 cholinesterase inhibitor and acts on the central nervous system. One active site of the  
88 molecule is the carbamate function (Mustazza et al. 2002, Yao and Li 2008), a function often  
89 present in drugs and pesticides (insecticides and herbicides) (Gupta and Ramesh 2006).  
90 Compounds with a carbamate function may undergo hydrolysis in the environment, but many  
91 of these are found in ground and surface waters (Campbell et al. 2004, García de Llasera and  
92 Bernal-González 2001).

93 Recently we investigated the direct degradation of rivastigmine under mono- and  
94 polychromatic wavelengths underlining the phototransformation under environmental-like  
95 conditions (Temussi et al. 2012).

96 In this work the transformation of rivastigmine toward hydroxyl radical generated using  
97 different natural sources ( $\text{H}_2\text{O}_2$ ,  $\text{NO}_3^-$ ,  $\text{NO}_2^-$ ) was investigated showing the possible enhanced  
98 transformation attributed to the  $\text{NO}_2\bullet$  and  $\text{NO}\bullet$  reactivity. Moreover three irradiations were  
99 performed in natural waters (lake, river and rain) in order to compare the rivastigmine  
100 degradation with previously obtained results in synthetic waters. A particular attention was

101 given to the degradation products showing the synergic effect of hydroxyl and NO<sub>x</sub> radicals  
102 on the rivastigmine transformation in waters.  
103

## 104 **2. Materials and Methods**

### 105 **2.1 Chemicals**

106 Rivastigmine hydrogen tartrate (99 %) was purchased by Kemprotec. Hydrogen peroxide (30  
107 %), sodium carbonate (99.5 %) and sodium nitrate (99 %) were purchased from Fluka;  
108 sodium sulphate anhydrous (99.5 %) and sodium nitrite (98%) were purchased from Prolabo;  
109 phosphoric acid (85% in water), sodium chloride (98 %), terephthalic acid (98 %) and 2-  
110 hydroxyterephthalic acid (97 %) were purchased from Sigma Aldrich without additional  
111 purification. All solvents were of HPLC grade and were purchased from Sigma-Aldrich.  
112 Fresh solutions doped with nitrate, nitrite or H<sub>2</sub>O<sub>2</sub> were prepared before each experiment. The  
113 concentration of the stock solution of H<sub>2</sub>O<sub>2</sub> in milli-Q water was determined using a molar  
114 absorption coefficient of  $38.1 \pm 1.4 \text{ M}^{-1} \text{ cm}^{-1}$  at 240 nm (Miller and Kester 2002).

115

### 116 **2.2 Irradiation experiments**

117 Aqueous solutions were irradiated in a thermostated cylindrical reactor. The 40 mL reactor,  
118 cooled by water circulation at a temperature of  $15 \pm 2 \text{ }^\circ\text{C}$  in order to limit thermal reactions,  
119 was located at one focal point of the lamp in order to maintain a constant irradiation of the  
120 whole sample and it was equipped on the top with a pyrex filter removing the wavelengths  
121 lower than ~285 nm. Samples were continuously stirred with a magnetic stirrer and a Teflon  
122 bar to ensure homogeneity.

123 The emission spectrum of the Xenon lamp (see Figure 1) was recorded using a fiber optics  
124 coupled with a CCD spectrophotometer (Ocean Optics USD 2000+UV-VIS). A reference  
125 lamp (DH-2000-CAL, Ocean Optics) was used for calibration. The emission spectrum

126 reaching the reactor surface was calculated to be  $4.31 \times 10^{19}$  photons  $\text{cm}^{-2} \text{s}^{-1}$  over the  
127 wavelength range 290-400 nm.

128

### 129 **2.3 Detection of hydroxyl radicals**

130 The hydroxyl radical formation rate ( $R_{OH}^f$ ) was determined by using terephthalic acid (TA) as  
131 trapping molecule. TA reacts with hydroxyl radical leading to the formation of 2-  
132 hydroxyterephthalic acid (TAOH) with a yield ( $Y_{TAOH}$ ) determined as the ratio between the  
133 initial formation rate of TAOH ( $R_{TAOH}^f$ ) and the initial degradation rate of TA ( $R_{TA}^d$ ).  $Y_{TAOH}$   
134 ranged between 12 and 30% depending on the pH and temperature of aqueous media  
135 (Charbouillot et al. 2011). This method represents a simple and fast detection of  
136 photogenerated  $\cdot\text{OH}$  with a high sensitivity that allows detection of less the  $10^{-18}$  M of  
137 hydroxyl radical steady state concentration.

138 In order to calculate the formation rate of  $\cdot\text{OH}$  as a function of different aqueous media  
139 composition, different solutions were irradiated with the same setup used for rivastigmine  
140 photochemical experiments. An aliquot of solution (3 mL) was withdrawn and put in a  
141 fluorescence cuvette at fixed times. The cuvette was transferred into a Perkin-Elmer MPF 3 L  
142 spectrofluorimeter and TAOH was quantified by using a calibration curve previously made  
143 with standard solutions of TAOH. The concentration of TA used during all experiments was  
144 included between 500  $\mu\text{M}$  and 1 mM in order to trap all photogenerated  $\cdot\text{OH}$ .

145

### 146 **2.4 Natural waters**

147 Lake, river and rain waters, sampled on 2011, were collected in order to perform hydroxyl  
148 radical measurement and rivastigmine degradation studies. River water and lake water come  
149 from Artiere river and Chambon lake, respectively, both located in the Puy de Dôme region,  
150 France. Artiere river passes through Clermont-Ferrand agglomeration, Chambon lake is in the

151 massif of Dore mountains at 877 m above sea level. Rain water was collected on the 1<sup>st</sup>  
152 November 2011 at the "Cézeaux campus" (south of Clermont-Ferrand) located at 394 m of  
153 altitude on the south of Clermont-Ferrand city.

154 Natural waters samples, collected using a glass bottle, were filtered on 0.20 µm membranes  
155 (Minisart<sup>®</sup>, Sartorium Stedim) and stored under refrigeration. Irradiation experiments were  
156 performed the day after, while, for analysis by ion chromatography (IC) inorganic and organic  
157 anions concentrations were determined no more than 24 hours after sampling.

158

## 159 **2.5 Photoinduced degradation and data analysis.**

160 The experiments were performed either in Milli-Q water or in natural waters. The solutions in  
161 Milli-Q water were prepared by mixing different concentrations of hydrogen peroxide (188  
162 µM, 376 µM, 550 µM 752 µM), or nitrate (200 µM, 400 µM, 5 mM) or nitrite (5.4 µM, 10  
163 µM, 50 µM, 100 µM) to a constant concentration of rivastigmine (45 µM). In order to analyze  
164 the time evolution of rivastigmine an aliquot (200 µL) was taken each time, stored in the dark  
165 and analysed by HPLC-UV. An HPLC system (Waters Alliance) equipped with a diode array  
166 detector was used for rivastigmine analysis. An Eclipse XDB-C18 column (Agilent, 4.6 x 150  
167 mm, 5 µm) and a gradient elution at a flow rate of 1.0 mL min<sup>-1</sup> were used. The gradient was:  
168 at initial time 10 % acetonitrile and 90 % water acidified with 3‰ phosphoric acid, it was  
169 kept for 7 min, and then followed by a linear gradient to 90 % acetonitrile within 8 min. Then,  
170 the same ratio was maintained constant for 8 min followed by a linear gradient to 10 %  
171 acetonitrile within 2 min. This ratio was maintained constant for 5 min. UV detector was set  
172 at 210 nm. To prevent reactions after sampling, amber glass vials were used. Retention time  
173 of rivastigmine was 11.2 min.

174 The time evolution of rivastigmine in the presence of photochemical sources of •OH could be  
175 fitted with a pseudo-first order equation  $C_0 = C_t \exp(-k t)$  where  $C_0$  was the initial



176 rivastigmine concentration,  $C_t$  the concentration at time  $t$  and  $k$  the pseudo-first order  
177 degradation rate constant. The error bars associated to the rate data represent  $3\sigma$ , derived from  
178 the scattering of the experimental data around the fitting curves (intra-series variability).

179 Ion chromatography analyses were performed using a DIONEX DX-320 instrument equipped  
180 with an IonPac AG11 (guard-column  $4 \times 50$  mm) and an IonPac AS11 (analytical column  $5 \times$   
181  $250$  mm) for anions. The elution step was performed using the following KOH concentration  
182 gradient: at initial time  $[\text{KOH}] = 0.2$  mM increasing up to  $0.43$  mM at  $4.5$  min, followed by a  
183 linear gradient to  $11.7$  mM up to  $18$  min. The  $[\text{KOH}]$  was then linearly increased up to  $33.5$   
184 mM within  $3$  min. The flow rate was  $1$  mL  $\text{min}^{-1}$ .

185 Pure rivastigmine solution was obtained by dissolving the tartrate salt in a saturated  $\text{Na}_2\text{CO}_3$   
186 aqueous solution. The solution was then extracted using ethyl acetate ( $3 \times 10$  mL) and the  
187 combined organic extracts were washed with brine and dried over anhydrous  $\text{Na}_2\text{SO}_4$ , before  
188 filtration and evaporation under vacuum. Absence of tartrate was confirmed by NMR  
189 spectroscopy and pure rivastigmine was stored in the dark at  $5^\circ\text{C}$ . Riv stability was checked  
190 by comparison with UV-vis spectrum (recorded from  $250$  to  $450$  nm) of fresh extracted  
191 solution (see Figure 1). pH values were measured with a Metrohm combined glass electrode,  
192 connected to a Metrohm 713 pH-meter.

193 Hydrogen peroxide concentration was determined by spectrophotometric absorbance  
194 detection of a solution in which *N,N*-diethyl-*p*-phenylenediamine (DPD) was involved in a  
195 reaction catalysed by a peroxidase (POD). In the presence of POD,  $\text{H}_2\text{O}_2$  oxidizes two  
196 molecules of DPD to cationic radicals  $\text{DPD}^{*\cdot}$  (Bader et al. 1988), which are stable and absorb  
197 light at  $551$  nm ( $\epsilon = 21\,000$   $\text{M}^{-1}$   $\text{cm}^{-1}$ ). With a  $10$  cm optical path length cell, the detection  
198 limit was  $0.1$   $\mu\text{M}$  calculated as  $3$  times the standard deviation of field blanks. Each  
199 measurement took less than  $5$  minutes.

200 Total organic carbon (TOC) and  $\text{HCO}_3^-/\text{CO}_3^{2-}$  concentrations in natural waters were  
201 quantified by TOC-5050A Analyzer (Shimadzu) using calibration curves previously  
202 performed on standard solutions.

203

## 204 **2.6 Phototransformation products identification**

205 Degradation products identification was performed by means of an LC-MS system (Agilent  
206 1100 Series, binary pump) equipped with an ESI ion source (MSD VL). The adopted column  
207 was a Sphere Clone C18 column (Phenomenex, 4.6 x 250 mm, 5  $\mu\text{m}$ ) and the following  
208 gradient elution was used: at initial time 30 % acetonitrile and 70 % water acidified with 1%  
209 formic acid, followed by a linear gradient to 75 % acetonitrile within 55 min. Then, the same  
210 ratio was maintained constant for 20 min. The flow rate was 0.4  $\text{mL min}^{-1}$  and the UV  
211 detector was settled at 254 nm.

212

## 213 **3. Results and discussion**


214 The goal of this work is to assess the reactivity of rivastigmine toward a wide range of  
215 photochemical precursors of hydroxyl radical in the aquatic environment upon irradiation of  
216 laboratory-made and natural solutions.

217

### 218 **3.1 Second order rate constant of rivastigmine with hydroxyl radical**

219 The second-order rate constant between Riv and hydroxyl radical was calculated by  
220 competition kinetics with isopropanol ( $(\text{CH}_3)_2\text{CHOH}$ , 2Pr) added to the solution at different  
221 concentrations. In Figure 2 we report the degradation rate of Riv obtained by irradiation of a  
222 solution containing 45  $\mu\text{M}$  of Riv, 380  $\mu\text{M}$  of  $\text{H}_2\text{O}_2$  and with addition of different isopropanol  
223 concentrations ranging from 0 up to 1 mM.  $\text{H}_2\text{O}_2$  was used as source of hydroxyl radical via  
224 its photolysis (R1) under polychromatic irradiation.

225 The following reactions were included in the kinetic treatment of data reported in this work:



230 The experimental data were fitted by using the following equation (eq1).

231 
$$\text{R}_{\text{Riv}}^d = \frac{\text{R}_{\cdot\text{OH}}^f k_{\text{Riv},\cdot\text{OH}} [\text{Riv}]}{k_{\text{Riv},\cdot\text{OH}} [\text{Riv}] + k_{\text{H}_2\text{O}_2,\cdot\text{OH}} [\text{H}_2\text{O}_2] + k_{2\text{Pr},\cdot\text{OH}} [2\text{Pr}]} \quad (\text{eq 1})$$

232 The data reported in Figure 2 were fitted with an equation of the type:  $y = \frac{a}{1+bx}$  where  $a$  is

233 the ratio  $\frac{\text{R}_{\cdot\text{OH}}^f k_{\text{Riv},\cdot\text{OH}} [\text{Riv}]}{k_{\text{Riv},\cdot\text{OH}} [\text{Riv}] + k_{\text{H}_2\text{O}_2,\cdot\text{OH}} [\text{H}_2\text{O}_2]}$ ,  $b$  is  $\frac{k_{2\text{Pr},\cdot\text{OH}}}{k_{\text{Riv},\cdot\text{OH}} [\text{Riv}] + k_{\text{H}_2\text{O}_2,\cdot\text{OH}} [\text{H}_2\text{O}_2]}$  and  $x$  the

234 concentration of isopropanol. From the data fit we can obtain the second-order reaction rate

235 constant between hydroxyl radical and rivastigmine:  $k_{\text{Riv},\cdot\text{OH}} = \frac{\frac{k_{2\text{Pr},\cdot\text{OH}}}{b} - k_{\text{H}_2\text{O}_2,\cdot\text{OH}} [\text{H}_2\text{O}_2]}{[\text{Riv}]}$

236 To the best of our knowledge,  $k_{\text{Riv},\cdot\text{OH}}$  was determined for the first time to be  $(5.8 \pm 0.3) \times$

237  $10^9 \text{ M}^{-1} \text{ s}^{-1}$ . Additionally we obtained the formation rate of  $\cdot\text{OH}$  from  $\text{H}_2\text{O}_2$  photolysis under

238 the adopted irradiation condition:  $\text{R}_{\cdot\text{OH}}^f = \frac{a (k_{\text{Riv},\cdot\text{OH}} [\text{Riv}] + k_{\text{H}_2\text{O}_2,\cdot\text{OH}} [\text{H}_2\text{O}_2])}{k_{\text{Riv},\cdot\text{OH}} [\text{Riv}]} = 9.6 \times 10^{-11}$

239  $\text{M s}^{-1}$  in agreement with results obtained via direct quantification using TA.

240

### 241           **3.2 Reactivity toward different hydroxyl radical sources**

242   The stability of rivastigmine in aqueous solution was checked under dark conditions at pH 6.0  
243   and no significant degradation was observed after 1 month, moreover direct photogredation  
244   results to be negligible under our irradiation conditions. Rivastigmine (45  $\mu\text{M}$ ) was then  
245   irradiated in the presence of different amounts of hydrogen peroxide, nitrate and nitrite. First  
246   of all the hydroxyl radical formation rate ( $R_{\text{OH}}^{\text{f}}$ ) was estimated as function of the  
247   photochemical concentration by using 1 mM TA as reported in Table 1.

248   As expected, the degradation rate of rivastigmine ( $R_{\text{Riv}}^{\text{d}}$ ) increases with the increasing of the  
249   concentration of the photochemical source of  $\bullet\text{OH}$ . In Figure 3 we report the initial  
250   degradation rate of rivastigmine  $R_{\text{Riv}}^{\text{d}}$  as function of hydrogen peroxide and nitrite  
251   concentration. In the case of hydrogen peroxide a competition due to the reaction R4 could be  
252   predictable. Moreover using nitrite as  $\bullet\text{OH}$  precursor the source reacts with photogenerated  
253   hydroxyl radicals leading to the formation of nitrogen dioxide ( $\text{NO}_2\bullet$ ) with a second order rate  
254   constant  $k_{\text{NO}_2\bullet, \bullet\text{OH}} = 1.0 \times 10^{10} \text{ M}^{-1} \text{ s}^{-1}$  (Mack and Bolton 1999). The plateau observed in figure  
255   3 using 50  $\mu\text{M}$  of nitrite could be explained taking into account its competitive role at high  
256   concentrations toward photogenerated hydroxyl radical. In fact, considering the initial  
257   concentration of Riv (45  $\mu\text{M}$ ) and nitrite (50  $\mu\text{M}$ ) in water we can argue that 34 % of  
258   hydroxyl radical reacts with Riv while up to 66 % with nitrite. Additionally the percentage  
259   significantly changes using 10  $\mu\text{M}$  of nitrite. Under such conditions the 73 % of  $\bullet\text{OH}$  reacts  
260   with Riv while only 28 % with nitrite. Using nitrite and/or hydrogen peroxide a fraction of  
261   photogenerated hydroxyl radical reacts also with their sources. From this assumption it is then  
262   possible to quantify the fraction of hydroxyl radical reacting with rivastigmine for each  
263   experimental point considering the initial concentrations of all species present in the solution  
264   and related second order rate constants with  $\bullet\text{OH}$ . Therefore the correlation of the hydroxyl  
265   radical formation rate (generated form hydrogen peroxide, nitrite and nitrate photolysis)

266 reacting with Riv??? is reported in Figure 4 showing that, under adopted concentrations, the  
267 dependences are linear as expected. Nevertheless, using nitrite and nitrate, the rivastigmine  
268 degradation increases compared to the value obtained in the presence of hydrogen peroxide.  
269 Such trend could be explained considering the additional reactivity of  $\text{NO}_2^\bullet$  (and probably  
270  $\text{NO}^\bullet$ ) generated via nitrate and nitrite direct photolysis or reaction with hydroxyl radical  
271 leading to the formation of both hydroxylated and nitro-derived compounds (Vione et al.  
272 2001, 2004). Such radicals could directly react with Riv and/or with radical intermediates  
273 generated after H-abstraction driven by photogenerated hydroxyl radical. In any case  
274 nitroderived radicals are generated and their reactivity toward Riv can explain the degradation  
275 enhancement discussed in the following section.

276

### 277 **3.3 Product identification**

278 Three experiments were conducted using a rivastigmine concentration of 4 mM in the  
279 presence of hydrogen peroxide 3.5 mM (experiment A), nitrates 15 mM (experiment B) and  
280 nitrite 50  $\mu\text{M}$  (experiment C). After 48 hours of irradiation the solutions were analyzed using  
281 an HPLC-UV-MS.

282 Three main reaction-type mechanisms were discerned and the structures of the nineteen  
283 identified products are proposed in Figure 5 on the basis of their mass spectra. The main  
284 reaction, as expected in the presence  $^\bullet\text{OH}$  precursors, is the hydroxyl radical addition to the  
285 aromatic ring leading to the formation of product **A1**, while di- and tri- hydroxylated  
286 compounds (**A2** and **A3**) are successively generated. Alcohol (**A4**), ketone (**A5**) and vinyl  
287 (**A6**) derivatives could also be formed from direct photolysis via a homolytic and/or  
288 heterolytic cleavage of the N-C bond as recently reported (Temussi et al. 2012) or via  
289 hydroxyl radical attach as shown during mineralization of sulfamethoxazole (Wang et al.  
290 2011).

291 Hydroxylation of **A4** and **A5** leads to the formation of **A7** and **A8** found only in trace levels  
292 under the adopted irradiation conditions.

293 In the presence of nitrite and nitrate two species with  $m/z [M+H]^+ = 280$  and  $310$  suggest that  
294 nitration of the aromatic ring (**B1** and **B2**) could occur (Nélieu et al. 2004).

295 Few products were also found under all experimental conditions, with  $m/z [M+H]^+ = 121$ ,  
296  $123$ ,  $137$ ,  $166$ ,  $178$  and  $180$  obtained from the carbamate chain cleavage or hydroxylation of  
297 previously generated oxidation products. Moreover, high mass values were found ( $m/z = 411$ ,  
298  $429$ ,  $445$  and  $467$ ) suggesting the formation of dimer molecules produced *via* radical  
299 recombination.

300

### 301 **3.4 Irradiation in Natural waters**

302 Three natural waters (rain, lake and river) were doped with Riv ( $45 \mu\text{M}$ ) in order to  
303 investigate the role of hydroxyl radical sources on the rivastigmine degradation.

304 Inorganic anions, TOC and carbonates concentrations are reported in Table 2. The main  
305 differences are the nitrate and nitrite concentrations found in river water:  $231 \mu\text{M}$  and  $\sim 7 \mu\text{M}$   
306 respectively, while in rain and lake they are present only at  $\mu\text{M}$  levels or under the detection  
307 limit. Also carbonates are more concentrated in river as well as chlorine and sulphate ions,  
308 while similar TOC concentrations are found in the three samples.

309 Hydroxyl radical formation rate ( $R_{\text{OH}}^{\text{f}}$ ) measured in lake and rain water using TA as trapping  
310 probe (Charbouillot et al. 2011) are in agreement with previously reported values in literature  
311 (Albinet et al. 2010, Vione et al. 2010). The calculated formation rate values range from  $2.0 \times$   
312  $10^{-11}$  up to  $6.5 \times 10^{-11} \text{ M s}^{-1}$  and from  $3.2 \times 10^{-12}$  to  $4.0 \times 10^{-11} \text{ M s}^{-1}$  for rain and lake waters,  
313 respectively. In our experiment the  $R_{\text{OH}}^{\text{f}}$  in river water was 6-7 times higher than in lake and  
314 rain. Such value could be explained considering the high nitrate and nitrite concentration in  
315 river samples. Moreover measured  $R_{\text{OH}}^{\text{f}}$  is well estimated showing that nitrite and nitrate

316 contributions represent about 57 and 22 % of the total hydroxyl radical formation in such  
 317 medium. Nevertheless for rain and lake waters about 47 and 98 % of hydroxyl radicals are  
 318 produced by other photochemical sources such as free iron, iron-complexes or chromophoric  
 319 dissolved organic matter (CDOM) via its excited state (CDOM\*). The possible impact of the  
 320 carbonate radicals on the rivastigmine degradation rate founded in river water could be  
 321 neglected. In fact carbonates can act as hydroxyl radical scavengers, generating carbonate  
 322 radical, which could induce organic molecule degradation via electron transfer reaction and/or  
 323 hydrogen abstraction with a quite low second order rate constant ( $10^{-6}$ - $10^{-7}$  M<sup>-1</sup> s<sup>-1</sup>) compared  
 324 to •OH. Concerning the "scavenging role", if we consider the relative low carbonates  
 325 concentrations founded in natural waters used in this work (in river water we have 1 mM  
 326 while for others the concentration not exceed  $1.5 \times 10^{-4}$  M) we can argue, taking into account  
 327 the carbonate (HCO<sub>3</sub><sup>-</sup>/CO<sub>3</sub><sup>2-</sup>), nitrite and Riv initial concentrations and the second order rates  
 328 constants with •OH ( $k_{\text{HCO}_3^-, \bullet\text{OH}} = 8.5 \times 10^6$  M<sup>-1</sup> s<sup>-1</sup> (Buxton et al. 1998)  $k_{\text{NO}_2^-, \bullet\text{OH}} = 1.0 \times 10^{10}$   
 329 M<sup>-1</sup> s<sup>-1</sup> (Mack and Bolton 1999) and  $k_{\text{Riv}, \bullet\text{OH}} = 5.8 \times 10^9$  M<sup>-1</sup> s<sup>-1</sup> (this work)) that, in river  
 330 water, about 0.25% of photogenerated hydroxyl radicals reacts with hydrogenocarbonate (at  
 331 pH 7.9 more then 98% of total carbonate species are hydrogenocarbonate), ~22 % with nitrite  
 332 and up then 78 % with rivastigmine.

333

### 334 **3.5 Radical driven degradation in natural selected waters**

335 In order to estimate the fate of the selected pollutant in natural waters, considering the double  
 336 role of naturally occurring compounds, the hydroxyl radical steady-state concentration  
 337 [ $\bullet\text{OH}$ ]<sub>ss</sub> was measured in three natural waters by using TA (1 mM) as chemical probe  
 338 following equation 2:

$$339 \quad -\frac{d[\text{TA}]}{dt} = k_{\text{TA}, \bullet\text{OH}} [\text{TA}] [\bullet\text{OH}]_{\text{ss}} \quad (\text{eq 2})$$

340 where [TA] and  $-d[TA]/dt$  are the initial concentration and the degradation rate of TA  
 341 respectively,  $k_{TA,\bullet OH}$  the second order rate constant between the probe and photogenerated  
 342 hydroxyl radical ( $3.9 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ ) (Charbouillot et al. 2011). The  $[\bullet OH]_{ss}$  values range from  
 343  $1.4 \times 10^{-18} \text{ M}$  for Lake up to  $1.5 \times 10^{-17} \text{ M}$  for River water (Table 3). Experimental results are  
 344 in agreement with those reported by Brezonik and Fulkerson-Brekken (Brezonik and  
 345 Fulkerson-Brekken 1998) in 5 natural waters (river and lakes) for June mid-day sunlight  
 346 radiation. The authors reported  $[\bullet OH]_{ss}$  ranging from  $2.5 \times 10^{-17}$  up to  $1.0 \times 10^{-15} \text{ M}$  and  $9.0 \times$   
 347  $10^{-19}$  to  $5.5 \times 10^{-17} \text{ M}$  in surface and whole water column respectively.

348 The hydroxyl radical steady-state concentration allows us the estimation of the Riv  
 349 degradation rate considering only the hydroxyl radical reactivity and scavenging effect of  
 350 naturally present compounds such as nitrite, carbonates ( $\text{CO}_3^{2-}/\text{HCO}_3^-$ ) and CDOM using  
 351 equation 3.

$$352 \quad R_{Riv}^{d,\bullet OH} = k_{Riv,\bullet OH} [Riv][\bullet OH]_{ss} - \sum_i k_{i,\bullet OH} [S_i][\bullet OH]_{ss} \quad (\text{eq 3})$$

353 Where  $R_{Riv}^{d,\bullet OH}$  and [Riv] are the degradation rate of rivastigmine due to the  $\bullet OH$  and its  
 354 concentration,  $\sum_i k_{i,\bullet OH} [S_i]$  is calculated as the sum of the contribution of the initial  
 355 concentration of scavengers (nitrite, carbonates and CDOM) with the reaction rate constant  
 356 with hydroxyl radical reported above. For these calculations the second order rate constant  
 357 between CDOM and hydroxyl radical of  $2.3 \times 10^4 \text{ mg of C L}^{-1} \text{ s}^{-1}$  (?ignoranza mia . che  
 358 significa?) (Brezonik and Fulkerson-Brekken 1998) has been used. The  $R_{Riv}^d (\text{M s}^{-1})$   
 359 estimated considering only the role of hydroxyl radical using equation 3 (eq. 3) results to be  
 360 less than those measured during irradiation in natural waters showing that additional  
 361 degradation pathways are responsible for the rivastigmine degradation as observed under  
 362 irradiation in synthetic waters.



363 Using the second order rate constant between hydroxyl radical and rivastigmine ( $5.8 \times 10^9 \text{ M}^{-1}$   
364  $\text{s}^{-1}$ ) founded in this work and highest  $\bullet\text{OH}$  steady-state concentrations calculated in some  
365 natural waters ( $10^{-15}$ - $10^{-16}$  M) (Brezonik and Fulkerson-Brekken 1998, Vione et al. 2010) the  
366  $t_{1/2}$  of rivastigmine using equation 4

$$367 \quad t_{1/2} = \frac{\ln 2}{k_{\text{Riv}, \bullet\text{OH}} [\bullet\text{OH}]_{\text{ss}}} \quad (\text{eq 4})$$

368 could be estimated between 1 day up to 14 years?????. Nevertheless, as reported in the text  
369 the real degradation is expected to be considerably enhanced due to the presence of  
370 nitroderived reactive species.

371

#### 372 4. Conclusions

373 The degradation of rivastigmine drug was investigated toward different photochemical radical  
374 sources found in natural waters. The first correlation of the formation rate of hydroxyl radical  
375 underlines a participation of radicals such as  $\text{NO}_x\bullet$  during polychromatic degradation in  
376 mimic and natural media. The chemical structures of the main products generated *via*  
377 oxidation of Riv have been suggested and hydroxylation, dimethylamino and carbamate  
378 chains degradation as well as nitration have been identified as possible pathways to explain  
379 the degradation products formation. Moreover the irradiation of Riv in three natural waters  
380 underlines a key role of indirect degradation in natural media when direct photolysis is  
381 negligible. The discrepancies between the Riv degradation rates calculated considering only  
382 the effect of hydroxyl radical and those measured in three natural waters underline the  
383 possible enhancement of the pollutant degradation in the presence of naturally occurring  
384 compounds such as nitrate and nitrite. Such reactivity could sensibly accelerate the pollutant  
385 degradation and need to be considered for a realistic half-life time estimation of organic  
386 pollutants in natural media.

387

388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430

## **Bibliography**

- Albinet, A., Minero, C. and Vione, D., 2010. Photochemical generation of reactive species upon irradiation of rainwater: Negligible photoactivity of dissolved organic matter. *Science of the Total Environment* 408 (16), 3367-3373.
- Andreozzi, R., Raffaele, M. and Nicklas, P., 2003. Pharmaceuticals in STP effluents and their solar photodegradation in aquatic environment. *Chemosphere* 50 (10), 1319-1330.
- Bader, H., Sturzenegger, V. and Hoigné, J., 1988. Photometric method for the determination of low concentrations of hydrogen peroxide by the peroxidase catalyzed oxidation of N,N-diethyl-p-phenylenediamine (DPD). *Water Research* 22 (9), 1109-1115.
- Brezonik, P.L. and Fulkerson-Brekken, J., 1998. Nitrate-induced photolysis in natural waters: controls on concentrations of hydroxyl radical photo-intermediates by natural scavenging agents. *Environmental Science & Technology* 32 (19), 3004-3010.
- Burrows, H.D., Canle L, M., Santaballa, J.A. and Steenken, S., 2002. Reaction pathways and mechanisms of photodegradation of pesticides. *Journal of Photochemistry and Photobiology, B: Biology* 67 (2), 71-108.
- Buxton, G.V., Greenstock, C.L., Helman, W.P. and Ross, A.B., 1998. Critical review of rate constants for reactions of hydrated electrons, hydrogen atoms and hydroxyl radicals ( $\cdot\text{OH}/\text{O}^-$ ) in aqueous solution. *Journal of Physical and Chemical Reference Data* 17, 513-886.
- Campbell, S., David, M.D., Woodward, L.A. and Li, Q.X., 2004. Persistence of carbofuran in marine sand and water. *Chemosphere* 54 (8), 1155-1161.
- Cermola, F., DellaGreca, M., Iesce, M.R., Montanaro, S., Previtiera, L., Temussi, F. and Brigante, M., 2007. Irradiation of fluvastatin in water: Structure elucidation of photoproducts. *Journal of Photochemistry and Photobiology A: Chemistry* 189 (2-3), 264-271.
- Charbouillot, T., Brigante, M., Deguillaume, L. and Mailhot, G., 2012. Atmospheric aqueous-phase photoreactivity: correlation between the hydroxyl radical photoformation and pesticide degradation rate in atmospherically relevant waters. *Photochemistry and Photobiology* 88 (1), 32-37.
- Charbouillot, T., Brigante, M., Mailhot, G., Maddigapu, P.R., Minero, C. and Vione, D., 2011. Performance and selectivity of the terephthalic acid probe for  $\cdot\text{OH}$  as a function of temperature, pH and composition of atmospherically relevant aqueous media. *Journal of Photochemistry and Photobiology, A: Chemistry* 222, 70-76.
- Daughton, C.G. and Ternes, T.A., 1999. PPCPs in the environment: agents of subtle change. *Environmental Health Perspectives* 107, 907-944.
- de Bertrand, N. and Barceló, D., 1991. Photodegradation of the carbamate pesticides aldicarb, carbaryl and carbofuran in water. *Analytica Chimica Acta* 254 (1-2), 235-244.

431 Della Greca, M., Pinto, G., Pistillo, P., Pollio, A., Previtiera, L. and Temussi, F., 2008.  
432 Biotransformation of ethinylestradiol by microalgae. *Chemosphere* 70 (11), 2047-2053.  
433 DellaGreca, M., Fiorentino, A., Isidori, M., Lavorgna, M., Previtiera, L., Rubino, M. and  
434 Temussi, F., 2004. Toxicity of prednisolone, dexamethasone and their photochemical  
435 derivatives on aquatic organisms. *Chemosphere* 54 (5), 629-637.  
436 DellaGreca, M., Iesce, M.R., Previtiera, L., Rubino, M., Temussi, F. and Brigante, M., 2006.  
437 Degradation of lansoprazole and omeprazole in the aquatic environment. *Chemosphere* 63  
438 (7), 1087-1093.  
439 Doll, T.E. and Frimmel, F.H., 2003. Fate of pharmaceuticals--photodegradation by simulated  
440 solar UV-light. *Chemosphere* 52 (10), 1757-1769.  
441 Ellis, J.B., 2006. Pharmaceutical and personal care products (PPCPs) in urban receiving  
442 waters. *Environmental Pollution* 144 (1), 184-189.  
443 Farré, M.I., Pérez, S., Kantiani, L. and Barceló, D., 2008. Fate and toxicity of emerging  
444 pollutants, their metabolites and transformation products in the aquatic environment.  
445 *Trends in Analytical Chemistry* 27 (11), 991-1007.  
446 García de Llasera, M.P. and Bernal-González, M., 2001. Presence of carbamate pesticides in  
447 environmental waters from the northwest of Mexico: determination by liquid  
448 chromatography. *Water Research* 35 (8), 1933-1940.  
449 Gupta, R.C. and Ramesh, C.G. 2006 *Toxicology of Organophosphate & Carbamate*  
450 *Compounds*, pp. 5-24, Academic Press, Burlington.  
451 Halling-Sørensen, B., Nielsen, S.N., Lanzky, P.F., Igerslev, F., Lützholt, H.C. and Jørgensen,  
452 S.E., 1998. Occurrence, fate, and effects of pharmaceutical substances in the environment-  
453 a review. *Chemosphere* 36, 357-393.  
454 Jjemba, P.K., 2006. Excretion and ecotoxicity of pharmaceutical and personal care products  
455 in the environment. *Ecotoxicology and Environmental Safety* 63 (1), 113-130.  
456 Jørgensen, S.-E. and Halling-Sørensen, B., 2000. Drugs in the environment. *Chemosphere* 40,  
457 691-699.  
458 Khetan, S.K. and Collins, T.J., 2007. Human pharmaceuticals in the aquatic environment:a  
459 challenge to green chemistry. *Chemical Reviews* 107 (6), 2319-2364.  
460 Kümmerer, K., Steger-Hartmann, T. and Meyer, M., 1997. Biodegradability of the anti-  
461 tumour agent ifosfamide and its occurrence in hospital effluents and communal sewage.  
462 *Water Research* 31 (11), 2705-2710.  
463 Mack, J. and Bolton, J.R., 1999. Photochemistry of nitrite and nitrate in aqueous solution: a  
464 review. *Journal of Photochemistry and Photobiology, A: Chemistry* 128 (1-3), 1-13.  
465 McClellan, K. and Halden, R.U., 2010. Pharmaceuticals and personal care products in  
466 archived U.S. biosolids from the 2001 EPA national sewage sludge survey. *Water*  
467 *Research* 44 (2), 658-668.  
468 Miller, W.L. and Kester, D.R., 2002. Hydrogen peroxide measurement in seawater by (p-  
469 hydroxyphenyl)acetic acid dimerization. *Analytical Chemistry* 60 (24), 2711-2715.  
470 Murray, K.E., Thomas, S.M. and Bodour, A.A., 2010. Prioritizing research for trace  
471 pollutants and emerging contaminants in the freshwater environment. *Environmental*  
472 *Pollution* 158 (12), 3462-3471.  
473 Mustazza, C., Borioni, A., Giudice, M.R.D., Gatta, F., Ferretti, R., Meneguz, A., Volpe, M.T.  
474 and Lorenzini, P., 2002. Synthesis and cholinesterase activity of phenylcarbamates related  
475 to Rivastigmine, a therapeutic agent for Alzheimer's disease. *European Journal of*  
476 *Medicinal Chemistry* 37 (2), 91-109.  
477 Nélieu, S., Kerhoas, L., Sarakha, M. and Einhorn, J., 2004. Nitrite and nitrate induced  
478 photodegradation of monolinuron in aqueous solution. *Environmental Chemistry Letters* 2  
479 (2), 83-87.

- 480 Razavi, B., Ben Abdelmelek, S., Song, W., O'Shea, K.E. and Cooper, W.J., 2011.  
481 Photochemical fate of atorvastatin (lipitor) in simulated natural waters. *Water Research* 45  
482 (2), 625-631.
- 483 Richardson, B.J., Lam, P.K.S. and Martin, M., 2005. Emerging chemicals of concern:  
484 Pharmaceuticals and personal care products (PPCPs) in Asia, with particular reference to  
485 Southern China. *Marine Pollution Bulletin* 50 (9), 913-920.
- 486 Richardson, S.D. and Ternes, T.A., 2005. Water Analysis: Emerging Contaminants and  
487 Current Issues. *Analytical Chemistry* 77, 3807-3838.
- 488 Sanz-Asensio, J., Plaza-Medina, M., Martínez-Soria, M.T. and Pérez-Clavijo, M., 1999.  
489 Study of photodegradation of the pesticide ethiofencarb in aqueous and non-aqueous  
490 media, by gas chromatography-mass spectrometry. *Journal of Chromatography, A* 840 (2),  
491 235-247.
- 492 Temussi, F., Passananti, M., Previtiera, L., Iesce, M.R., Brigante, M., Mailhot, G. and  
493 DellaGreca, M., 2012. Phototransformation of the drug rivastigmine: Photoinduced  
494 cleavage of benzyl-nitrogen sigma bond. *Journal of Photochemistry and Photobiology A*  
495 239 (1), 1-6.
- 496 Ternes, T.A., Joss, A. and Siegrist, H., 2004. Scrutinizing pharmaceuticals and personal care  
497 products in wastewater treatment. *Environmental Science and Technology* 38 (20), 392A-  
498 399A.
- 499 Vaal, M., van der Wal, J.T., Hoekstra, J. and Hermens, J., 1997. Variation in the sensitivity of  
500 aquatic species in relation to the classification of environmental pollutants. *Chemosphere*  
501 35 (6), 1311-1327.
- 502 Vione, D., Bagnus, D., Maurino, V. and Minero, C., 2010. Quantification of singlet oxygen  
503 and hydroxyl radicals upon UV irradiation of surface water. *Environmental Chemistry*  
504 *Letters* 8 (2), 193-198.
- 505 Vione, D., Maurino, V., Minero, C. and Pelizzetti, E., 2001. Phenol photonation upon UV  
506 irradiation of nitrite in aqueous solution II: effects of pH and TiO<sub>2</sub>. *Chemosphere* 45 (6-7),  
507 903-910.
- 508 Vione, D., Maurino, V., Minero, C. and Pelizzetti, E., 2004. Phenol nitration upon oxidation  
509 of nitrite by Mn(III,IV) (hydr)oxides. *Chemosphere* 55 (7), 941-949.
- 510 Wang, A., Li, Y.-Y. and Estrada, A.L., 2011. Mineralization of antibiotic sulfamethoxazole  
511 by photoelectro-Fenton treatment using activated carbon fiber cathode and under UVA  
512 irradiation. *Applied Catalysis B: Environmental* 102 (3-4), 378-386.
- 513 Yao, Y. and Li, Z.-s., 2008. Inhibition Mechanism of Cholinesterases by Carbamate: A  
514 Theoretical Study. *Chemical Research in Chinese Universities* 24 (6), 778-781.
- 515 Yu, C.-P. and Chu, K.-H., 2009. Occurrence of pharmaceuticals and personal care products  
516 along the West Prong Little Pigeon River in east Tennessee, USA. *Chemosphere* 75 (10),  
517 1281-1286.

518  
519

520

521

522

523

524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546


**Figure Caption**

- 1) Emission spectrum reaching the solution and molar absorption coefficients of the aqueous rivastigmine, nitrate, nitrite and H<sub>2</sub>O<sub>2</sub> solutions.
- 2) Initial degradation rate of Riv under polychromatic irradiation of H<sub>2</sub>O<sub>2</sub> 380 μM as a function of the isopropanol concentration. The solid curve shows the fit of experimental data with equation 1 and dashed lines denote the 95% confidence interval of this fit. The errors bars represent the 3σ based on the linear fit of experimental data.
- 3) Correlation between the initial degradation rate of Riv ( $R_{Riv}^d$ ) with hydrogen peroxide and nitrite concentration under polychromatic irradiation. The errors represent the 3σ based on the linear fit of experimental data.
- 4) Correlation between the initial degradation rate of Riv ( $R_{Riv}^d$ ) and the formation rate of hydroxyl radical ( $R_{OH,Riv}^f$ ) reacting with rivastigmine. The photochemical sources

547 are: hydrogen peroxyde (empty circles) nitrites (full circles) and nitrates (empty  
548 triangles). Dashed lines denote the 95% confidence of the linear fit.

549

550 5) Proposed structures and formation mechanisms for the main rivastigmine  
551 degradation products found in the presence of hydrogen peroxide, nitrites and  
552 nitrates as photochemical sources of hydroxyl radical under polychromatic  
553 irradiation.



461

462 **Figure 1**


463

464

465

466

467




468

469 **Figure 2**


470





471  
472 **Figure 3**

473



474

475 **Figure 4**

476

477

478

479

480

481

482

483


484

485

486

487

488



489  
 490  
 491  
 492  
 493  
 494  
 495

Figure 5

	H <sub>2</sub> O <sub>2</sub>	NO <sub>3</sub> <sup>-</sup>	NO <sub>2</sub> <sup>-</sup>
k (s <sup>-1</sup> )	(2.35 ± 0.13) x 10 <sup>-6</sup>	(3.34 ± 0.21) x 10 <sup>-8</sup>	(2.75 ± 0.06) x 10 <sup>-6</sup>

**Table 1:** Rate constants of <sup>•</sup>OH formation from different sources (H<sub>2</sub>O<sub>2</sub>, NO<sub>3</sub><sup>-</sup> or NO<sub>2</sub><sup>-</sup>).

496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521

	Rain water	Lake water	River water
<b>NO<sub>3</sub><sup>-</sup> (M)</b>	7.49 x 10 <sup>-6</sup>	3.53 x 10 <sup>-6</sup>	2.31 x 10 <sup>-4</sup>
<b>NO<sub>2</sub><sup>-</sup> (M)</b>	1.15 x 10 <sup>-6</sup>	/	7.30 x 10 <sup>-6</sup>
<b>SO<sub>4</sub><sup>2-</sup> (M)</b>	3.98 x 10 <sup>-6</sup>	5.09 x 10 <sup>-5</sup>	3.75 x 10 <sup>-4</sup>
<b>Cl<sup>-</sup> (M)</b>	4.99 x 10 <sup>-5</sup>	1.97 x 10 <sup>-4</sup>	3.47 x 10 <sup>-3</sup>
<b>HCO<sub>3</sub><sup>-</sup>/CO<sub>3</sub><sup>2-</sup> (M)</b>	2.0	9.32	61.23
<b>TOC (mg/L)</b>	2.45	2.01	2.70
<b>pH</b>	7.2	6.6	7.9
<b>R<sup>f</sup><sub>OH measured</sub> (M s<sup>-1</sup>)<sup>a</sup></b>	(6.42 ± 0.53) x 10 <sup>-12</sup>	(5.45 ± 0.08) x 10 <sup>-12</sup>	(3.55 ± 0.02) x 10 <sup>-11</sup>
<b>R<sup>f</sup><sub>OH predicted</sub> (M s<sup>-1</sup>)<sup>b</sup></b>	3.42 x 10 <sup>-12</sup>	1.21 x 10 <sup>-13</sup>	2.80 x 10 <sup>-11</sup>
<b>% <sup>•</sup>OH formation from NO<sub>3</sub><sup>-</sup> <sup>c</sup></b>	4.0	2.2	22.4
<b>% <sup>•</sup>OH formation from NO<sub>2</sub><sup>-</sup> <sup>c</sup></b>	49.3	/	56.6
<b>% <sup>•</sup>OH formation from other <sup>d</sup></b>	46.7	97.8	21.2
<b>R<sup>d</sup><sub>Riv</sub> (M s<sup>-1</sup>)</b>	(2.47 ± 1.20) x 10 <sup>-12</sup>	(2.83 ± 1.63) x 10 <sup>-12</sup>	(1.02 ± 0.05) x 10 <sup>-11</sup>

522 **Table 2:** Concentrations of nitrite, nitrate, HCO<sub>3</sub><sup>-</sup>/CO<sub>3</sub><sup>2-</sup> and Total organic carbon (TOC) of  
523 natural waters used during rivastigmine experiments. <sup>a</sup> <sup>•</sup>OH formation rate obtained by using 1mM of  
524 terephthalic acid as trapping molecule. <sup>b</sup> Theoretical <sup>•</sup>OH formation rate estimated by considering only  
525 NO<sub>2</sub><sup>-</sup> and NO<sub>3</sub><sup>-</sup> as photochemical sources. <sup>c</sup> Contribution to the hydroxyl radicals attributed to NO<sub>2</sub><sup>-</sup>  
526 and O<sub>3</sub>. <sup>d</sup> Contribution to the hydroxyl radicals obtained by difference from known sources.

Formatted: Highlight