



A categorial equivalence for semi-Nelson algebras

Juan Manuel Cornejo¹ · Andrés Gallardo¹ · Ignacio Viglizzo¹

Accepted: 23 August 2021 / Published online: 13 September 2021
© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2021

Abstract

We present a category equivalent to that of semi-Nelson algebras. The objects in this category are pairs consisting of a semi-Heyting algebra and one of its filters. The filters must contain all the dense elements of the semi-Heyting algebra and satisfy an additional technical condition. We also show that the category of dually hemimorphic semi-Nelson algebras is equivalent to that of dually hemimorphic semi-Heyting algebras.

1 Introduction

Semi-Heyting algebras were introduced in 1985 by Sankappanavar (1985) as a variety generalizing that of Heyting algebras while keeping many of its good features like being distributive pseudocomplemented lattices and having their congruences determined by filters.

On the other hand, D. Vakarelov provided in Vakarelov (1977) a way of constructing Nelson algebras from Heyting ones, by means of what is now known as twist product, thus extending work by Kalman (1958). This program was continued by Sendlewski (1984), Sendlewski (1990), who gave a full representation of Nelson algebras using Heyting algebras and a boolean congruence on them. Later, L. Monteiro and I. Viglizzo, in Viglizzo (1999) and Monteiro and Viglizzo (2019), used filters instead of congruences to study this representation.

It turned out that another one of the nice features of semi-Heyting algebras is that Vakarelov's construction can be carried out on them as well. The resulting algebras form the variety of semi-Nelson algebras (Cornejo and Viglizzo 2018). Vakarelov's construction gives a categorial equivalence between semi-Heyting algebras and *centered* semi-Nelson algebras (Cornejo and San Martín 2018), but this leaves out many semi-Nelson algebras. In this work, we prove that ordered pairs consisting of a semi-Heyting algebra and a filter in it yield all semi-Nelson algebras. This gives a

good representation that can be seen as a categorial equivalence and permits understanding semi-Nelson algebras in terms of the better-known semi-Heyting algebras.

H. Sankappanavar defined in Sankappanavar (2011) the variety of semi-Heyting algebras with a dual hemimorphism, or dually hemimorphic semi-Heyting algebras, \mathcal{DSH} , as an expansion of semi-Heyting algebras with a unary operator that is a common generalization of both De Morgan and the pseudocomplement operations. J. M. Cornejo and H. San Martín applied Vakarelov's construction to this expansion to obtain the variety of dually hemimorphic semi-Nelson algebras, \mathcal{DSN} (Cornejo and San Martín 2018). We prove here that dually hemimorphic semi-Nelson algebras are centered, and therefore they can be represented by dually hemimorphic semi-Heyting algebras.

The paper is organized as follows: Section 2 introduces the varieties of algebras we will be dealing with, recalls some of the basic results on them that we will be using, and provides background on Vakarelov's construction. Section 3 details the representation of semi-Nelson algebras by pairs consisting of a semi-Heyting algebra and one of its filters containing the dense elements of the algebra and satisfying an additional condition, proving this representation is an equivalence of two categories. The final section deals with the case of dually hemimorphic semi-Nelson algebras.

2 Preliminaries

2.1 Definitions and elementary properties

In this section, we will recall the definitions and some basic properties of the varieties of Heyting and semi-Heyting

✉ Ignacio Viglizzo
viglizzo@gmail.com

¹ Instituto de Matemática (INMABB), Departamento de Matemática, Universidad Nacional del Sur (UNS) - CONICET, Av. Alem 1253, Bahía Blanca, Buenos Aires, Argentina

algebras, Nelson and semi-Nelson algebras, pseudocomplemented lattices, and dually hemimorphic semi-Heyting and semi-Nelson algebras.

Definition 2.1 A *semi-Heyting algebra* is an algebra $\mathbf{H} = \langle H; \wedge, \vee, \Rightarrow, 0, 1 \rangle$ of type $(2, 2, 2, 0, 0)$ such that the following identities are satisfied for all $x, y, z \in A$:

- (SH1) $\langle H; \wedge, \vee, 0, 1 \rangle$ is a bounded lattice,
- (SH2) $x \wedge (x \Rightarrow y) = x \wedge y$,
- (SH3) $x \wedge (y \Rightarrow z) = x \wedge ((x \wedge y) \Rightarrow (x \wedge z))$,
- (SH4) $x \Rightarrow x = 1$.

A semi-Heyting algebra is a *Heyting algebra* if it satisfies the identity:

$$(H) \quad (x \wedge y) \Rightarrow x = 1.$$

On a semi-Heyting algebra \mathbf{H} , one can always define the term $x \Rightarrow_{\mathbf{H}} y = x \Rightarrow (x \wedge y)$. With this operation, the system $\langle H; \wedge, \vee, \Rightarrow_{\mathbf{H}}, 0, 1 \rangle$ is a Heyting algebra (Abad et al. 2013).

Nelson algebras are the algebraic counterpart of D. Nelson’s constructive logic with strong negation. The identities presented here were proved to be a complete and independent axiomatization of Nelson algebras in Monteiro and Monteiro (1996):

Definition 2.2 A *Nelson algebra* is an algebra $\mathbf{A} = \langle A; \wedge, \vee, \rightarrow, \sim, 1 \rangle$ of type $(2, 2, 2, 1, 0)$ such that the following conditions are satisfied for all $x, y, z \in A$:

- (N1) $x \wedge (x \vee y) = x$,
- (N2) $x \wedge (y \vee z) = (z \wedge x) \vee (y \wedge x)$,
- (N3) $\sim \sim x = x$,
- (N4) $\sim (x \wedge y) = \sim x \vee \sim y$,
- (N5) $x \wedge \sim x = (x \wedge \sim x) \wedge (y \vee \sim y)$,
- (N6) $x \rightarrow x = 1$,
- (N7) $x \rightarrow (y \rightarrow z) = (x \wedge y) \rightarrow z$,
- (N8) $x \wedge (x \rightarrow y) = x \wedge (\sim x \vee y)$.

The variety of semi-Nelson algebras was introduced in Cornejo and Viglizzo (2018) as a generalization of Nelson algebras.

Definition 2.3 A *semi-Nelson algebra* is an algebra $\mathbf{A} = \langle A; \wedge, \vee, \rightarrow, \sim, 1 \rangle$ of type $(2, 2, 2, 1, 0)$ such that the following conditions are satisfied for all $x, y, z \in A$:

- (SN1) $x \wedge (x \vee y) = x$,
- (SN2) $x \wedge (y \vee z) = (z \wedge x) \vee (y \wedge x)$,
- (SN3) $\sim \sim x = x$,
- (SN4) $\sim (x \wedge y) = \sim x \vee \sim y$,
- (SN5) $x \wedge \sim x = (x \wedge \sim x) \vee (y \vee \sim y)$,

- (SN6) $x \wedge (x \rightarrow_N y) = x \wedge (\sim x \vee y)$,
- (SN7) $x \rightarrow_N (y \rightarrow_N z) = (x \wedge y) \rightarrow_N z$,
- (SN8) $(x \rightarrow_N y) \rightarrow_N [(y \rightarrow_N x) \rightarrow_N [(x \rightarrow z) \rightarrow_N (y \rightarrow z)]] = 1$,
- (SN9) $(x \rightarrow_N y) \rightarrow_N [(y \rightarrow_N x) \rightarrow_N [(z \rightarrow x) \rightarrow_N (z \rightarrow y)]] = 1$,
- (SN10) $(\sim (x \rightarrow y)) \rightarrow_N (x \wedge \sim y) = 1$,
- (SN11) $(x \wedge \sim y) \rightarrow_N (\sim (x \rightarrow y)) = 1$,

where $x \rightarrow_N y := x \rightarrow (x \wedge y)$.

For a semi-Nelson algebra \mathbf{A} , the system $\langle A; \wedge, \vee, \rightarrow_N, \sim, 1 \rangle$ is a Nelson algebra (Cornejo and Viglizzo 2018).

We will denote by $\mathcal{H}, \mathcal{SH}, \mathcal{N}$ and \mathcal{SN} the varieties of Heyting, semi-Heyting, Nelson and semi-Nelson algebras, respectively.

Heyting algebras, as is well known, are pseudocomplemented (with $x^* = x \rightarrow 0$ as the pseudocomplement of $x \in H$) in the sense of the following definition:

Definition 2.4 An algebra $\mathbf{A} = \langle A; \wedge, \vee, *, 0, 1 \rangle$ of type $(2, 2, 1, 0, 0)$ is a *pseudocomplemented lattice* if the following conditions hold for all $x, y, z \in A$:

- (PS1) $\langle A; \wedge, \vee, 0, 1 \rangle$ is a bounded lattice,
- (PS2) $x \wedge (x \wedge y)^* = x \wedge y^*$,
- (PS3) $0^* = 1$ and $1^* = 0$.

If we define in a semi-Heyting algebra the term $x^c = x \Rightarrow 0$, it turns out that x^c is the pseudocomplement of x in the sense of the previous definition. Moreover, $x \Rightarrow_{\mathbf{H}} 0 = x \Rightarrow (x \wedge 0) = x \Rightarrow 0$, so x^* and x^c coincide in \mathcal{SH} . From now on, we will use indistinctly the notation x^* for both x^c and x^* .

We will use some properties of pseudocomplemented lattices. Their proof can be found in Balbes and Dwinger (1974):

Proposition 2.5 If $\mathbf{A} = \langle A; \wedge, \vee, *, 0, 1 \rangle$ is a pseudocomplemented lattice, then for every $x, y \in A$, if $x \wedge y = 0$, then $x \leq y^*$. It also holds that $(x \vee y)^* = x^* \wedge y^*$.

Definition 2.6 Let $\mathbf{A} = \langle A; \wedge, \vee, *, 0, 1 \rangle$ be a pseudocomplemented lattice. We say that an element $a \in A$ is *dense* if $a^* = 0$. We denote by $\mathcal{D}(A)$ the set of all dense elements of A .

The following characterization of dense elements is going to be useful:

Lemma 2.7 If $\mathbf{H} \in \mathcal{SH}$, then $x \in \mathcal{D}(H)$ if and only if $x = y \vee y^*$ for some $y \in H$.

For the results in section 5, we now define the varieties of dually hemimorphic semi-Heyting and semi-Nelson algebras. The former were introduced by H.P. Sankappanavar in Sankappanavar (2011), and the latter were presented in Cornejo and San Martín (2020).

Definition 2.8 An algebra $\mathbf{A} = \langle A; \wedge, \vee, \rightarrow, \dagger, 0, 1 \rangle$ of type $(2, 2, 2, 1, 0, 0)$ is said to be a *dually hemimorphic semi-Heyting algebra* if $\langle A; \wedge, \vee, \rightarrow, 0, 1 \rangle$ is a semi-Heyting algebra and the following equations are satisfied for all $x, y, z \in A$:

- (DSM1) $0^\dagger = 1$,
- (DSM2) $1^\dagger = 0$,
- (DSM3) $(x \wedge y)^\dagger = x^\dagger \vee y^\dagger$.

We write DSH to denote the variety of dually hemimorphic semi-Heyting algebras.

Definition 2.9 An algebra $\mathbf{A} = \langle A; \wedge, \vee, \rightarrow, \sim, ', 1 \rangle$ of type $(2, 2, 2, 1, 1, 0)$ is said to be a *dually hemimorphic semi-Nelson algebra* if $\langle A; \wedge, \vee, \rightarrow, \sim, 1 \rangle$ is a semi-Nelson algebra and the following equations are satisfied for all $x, y, z \in A$:

- (DSN1) $(\sim 1)' = 1$,
- (DSN2) $1' \rightarrow (\sim 1) = 1$,
- (DSN3) $((x \rightarrow y) \wedge (y \rightarrow x) \wedge x') \rightarrow ((x \rightarrow y) \wedge (y \rightarrow x) \wedge y)' = 1$,
- (DSN4) $\sim x' \rightarrow (\sim x \wedge (x' \rightarrow x)) = 1$,
- (DSN5) $(\sim x \wedge (x' \rightarrow x)) \rightarrow \sim x' = 1$,
- (DSN6) $(x \wedge y)' \rightarrow (x' \vee y') = 1$,
- (DSN7) $(x' \vee y') \rightarrow (x \wedge y)' = 1$.

We use the convention that the unary operation $'$ has higher priority than \sim , so the expression $\sim x'$ means $\sim (x')$.

We write DSN to denote the variety of dually hemimorphic semi-Nelson algebras.

2.2 Vakarelov’s construction

In this section, we review the well-known twist construction that established the connection between Heyting and Nelson algebras, and later was used to define (dually hemimorphic) semi-Nelson algebras as a variety constructed from the one of (dually hemimorphic) semi-Heyting algebras.

Let $\mathbf{A} \in \mathcal{H}$. We denote with $V_k(A)$ the set $\{(a, b) \in A^2 : a \wedge b = 0\}$ and with $\mathbf{V}_k(\mathbf{A})$ the algebra $\langle V_k(A); \sqcap, \sqcup, \rightarrow, \sim, \top \rangle$ where the operations are defined as follows:

- (V1) $(a, b) \sqcap (c, d) = (a \wedge c, b \vee d)$,
- (V2) $(a, b) \sqcup (c, d) = (a \vee c, b \wedge d)$,
- (V3) $(a, b) \rightarrow (c, d) = (a \Rightarrow c, a \wedge d)$,
- (V4) $\sim (a, b) = (b, a)$,
- (V5) $\top = (1, 0)$.

Observation 2.10 From the previous definitions, we can deduce the rule

$$(V6) (a, b) \rightarrow_N (c, d) = (a \Rightarrow_H c, a \wedge d).$$

It was shown in Vakarelov (1977) that $\mathbf{V}_k(\mathbf{A}) \in \mathcal{N}$.

Vakarelov’s construction can be generalized the following way: if $\mathbf{A} \in \mathcal{H}$, and F is a filter of \mathbf{A} , we define the structure

$$N(A, F) := \{(a, b) \in A^2 : a \wedge b = 0, a \vee b \in F\}.$$

It was proved in Viglizzo (1999), Monteiro and Viglizzo (2019) that $N(A, F)$, algebraized with (V1)-(V5), is well defined and it is a Nelson algebra. This algebra will be denoted by $\mathbf{N}(\mathbf{A}, \mathbf{F})$.

A different generalization was to start from a semi-Heyting algebra, which lead to the definition of semi-Nelson algebras (Cornejo and Viglizzo 2018): if \mathbf{A} is a semi-Heyting algebra, then $\mathbf{V}_k(\mathbf{A})$ is a semi-Nelson algebra.

Going in the other direction, we can go from semi-Nelson algebras to semi-Heyting algebras by means of the following quotient:

Let $\mathbf{A} = \langle A; \wedge, \vee, \rightarrow, \sim, 1 \rangle \in \mathcal{SN}$, and define:

$$x \equiv y \text{ if and only if } x \rightarrow y = 1 \text{ and } y \rightarrow x = 1.$$

It can be proved that “ \equiv ” is an equivalence relation and a congruence with respect of the operations \wedge, \vee and \rightarrow . We denote with $\llbracket a \rrbracket$ the equivalence class of an element $a \in A$ under the relation \equiv .

We consider $\mathbf{sH}(\mathbf{A}) = \langle A/\equiv; \cap, \cup, \Rightarrow, \perp, \top \rangle$, where

- $\perp = \llbracket \sim 1 \rrbracket$,
- $\top = \llbracket 1 \rrbracket$,
- $\llbracket x \rrbracket \cap \llbracket y \rrbracket = \llbracket x \wedge y \rrbracket$,
- $\llbracket x \rrbracket \cup \llbracket y \rrbracket = \llbracket x \vee y \rrbracket$,
- $\llbracket x \rrbracket \Rightarrow \llbracket y \rrbracket = \llbracket x \rightarrow y \rrbracket$.

Then $\mathbf{sH}(\mathbf{A}) \in \mathcal{SH}$ (Cornejo and Viglizzo 2018).

We will use the next result in the following sections:

Proposition 2.11 (Cornejo and Viglizzo 2018, Lemma 2.7 (h) and (l)) *Let $\mathbf{A} \in \mathcal{SN}$, and $a, b, c \in A$. Then the following properties hold:*

- (1) $a \rightarrow_N b = b \rightarrow_N a = 1$ if and only if $a \rightarrow b = b \rightarrow a = 1$,
- (2) If $a \rightarrow_N b = 1 = b \rightarrow_N c = 1$, then $a \rightarrow_N c = 1$.

Part (1) of the previous proposition tells us that the congruence \equiv is the same as the one that can be defined if we replace \rightarrow with \rightarrow_N .

The following representation theorem establishes the connection between semi-Heyting and semi-Nelson algebras using Vakarelov’s construction and the quotient by the relation \equiv .

Theorem 2.12 (Cornejo and Viglizzo 2018, Theorem 5.3 and Corollary 5.2) *If $\mathbf{A} \in \mathcal{SH}$, then \mathbf{A} is isomorphic to $\mathbf{sH}(\mathbf{V}_k(\mathbf{A}))$ through the isomorphism $g(a) = \llbracket(a, a^*)\rrbracket$. Also, if $\mathbf{A} \in \mathcal{SN}$, then \mathbf{A} is isomorphic to a subalgebra of $\mathbf{V}_k(\mathbf{sH}(\mathbf{A}))$ through the application $h(a) = (\llbracket a \rrbracket, \llbracket \sim a \rrbracket)$.*

3 Representations

Theorem 2.12 gives a representation of any semi-Nelson algebra \mathbf{A} as a subalgebra of $\mathbf{V}_k(\mathbf{sH}(\mathbf{A}))$, but there can be more than one non-isomorphic semi-Nelson algebra having the same quotient semi-Heyting algebra $\mathbf{sH}(\mathbf{A})$ (up to isomorphism). In order to get a sharper representation, we are going to consider pairs consisting of a semi-Heyting algebra and one of its filters satisfying extra conditions. To be more precise, we obtain a categorial equivalence between the category of semi-Nelson algebras and a category whose objects are pairs (\mathbf{H}, F) where \mathbf{H} is a semi-Heyting algebra and F is a filter of \mathbf{H} that contains all the dense elements of \mathbf{H} and satisfies an extra condition.

Recall the construction $N(A, F) = \{(a, b) \in A^2 : a \wedge b = 0, a \vee b \in F\}$ for some Heyting algebra \mathbf{A} , and F a filter of \mathbf{A} . For any $\mathbf{A} \in \mathcal{H}$, we consider $\mathbf{V}_k(\mathbf{A})$, and a subalgebra \mathbf{S} such that the projection over the first component, π_1 , verifies $\pi_1(S) = A$. It was shown in Viglizzo (1999) that there is a filter $F \subseteq A$ containing the dense elements of A such that $\mathbf{S} = \mathbf{N}(\mathbf{A}, F)$.


To extend these results to the variety \mathcal{SH} , we define:

Definition 3.1 Let $\mathbf{H} \in \mathcal{SH}$. A subset $F \subseteq H$ is an *i*-filter, if the following conditions hold:

- (IF1) F is a lattice filter of \mathbf{H} .
- (IF2) $\mathcal{D}(H) \subseteq F$.
- (IF3) If $z \vee t \in F$, then for all $x \in H$, $(x \Rightarrow z) \vee (x \wedge t) \in F$.

It is immediate from the definition that every *i*-filter is a filter. The converse is not true, as the following example shows:

Example 3.2 Consider the semi-Heyting algebra $\mathbf{A} = \langle A; \wedge, \vee, \Rightarrow, 1 \rangle$, with the operation \Rightarrow indicated in the table:



The subset $F = \{1\}$ of \mathbf{A} is a filter, and it satisfies (IF2) (since the only dense element is 1). But F does not satisfy

(IF3) because $1 \vee 0 = 1 \in F$, but $(0 \Rightarrow 1) \vee (0 \wedge 0) = 0 \vee 0 = 0 \notin F$.

Observation 3.3 *If we consider the structure $\mathbf{N}(\mathbf{H}, F)$, with $\mathbf{H} \in \mathcal{SH}$, an *i*-filter F , and $(a, b), (c, d) \in N(H, F)$, then (1) $c \wedge d = 0$ and (2) $c \vee d \in F$. Notice that*

$$(a \Rightarrow c) \wedge (a \wedge d) = a \wedge (a \Rightarrow c) \wedge d \stackrel{(SH2)}{=} (a \wedge c) \wedge d = a \wedge (c \wedge d) \stackrel{(1)}{=} a \wedge 0 = 0.$$

Also, using (2) and (IF3) we have that $(a \Rightarrow c) \vee (a \wedge d) \in F$, so the pair $(a \Rightarrow c, a \wedge d)$ is in $N(H, F)$ and we can use it to define $(a, b) \rightarrow (c, d) = (a \Rightarrow c, a \wedge d)$.

Lemma 3.4 *If $\mathbf{H} \in \mathcal{SH}$ and F is an *i*-filter of \mathbf{H} , then the system $\langle N(H, F); \sqcap, \sqcup, \rightarrow, \sim, \top \rangle$ with the operations defined as in (V1)-(V5) is a semi-Nelson algebra.*

Proof Using that F is a filter we can prove that the operations \sqcap, \sqcup, \top and \sim are well defined in $N(H, F)$ (see Monteiro and Viglizzo 2019). Also, by Observation 3.3, it follows that the operation \rightarrow is well defined. Furthermore, following the proof of Theorem 4.1 in Cornejo and Viglizzo (2018), $\mathbf{N}(\mathbf{H}, F)$ verifies the axioms (SN1)-(SN11) so it is a semi-Nelson algebra. \square

Lemma 3.5 *Let $\mathbf{H} \in \mathcal{SH}$. If \mathbf{S} is a subalgebra of $\mathbf{V}_k(\mathbf{H})$, such that $\pi_1(S) = H$ (with π_1 the projection over the first component of $\mathbf{V}_k(H)$), then there exists an *i*-filter E of \mathbf{H} such that $S = N(H, E)$.*

Proof Let $E = \{x \in H : x = a \vee b \text{ for some } (a, b) \in S\}$. We are going to prove that E is an *i*-filter.

It was shown in Viglizzo (1999) that if $\mathbf{H} \in \mathcal{H}$, then E is a filter of \mathbf{H} . But that proof can be extended naturally to the variety \mathcal{SH} , because it only uses lattice properties.

Consider a dense element $x \in H$. Since $\pi_1(S) = H$, then there exists $y \in H$ such that $(x, y) \in S$. Besides, since S is a subalgebra of $\mathbf{V}_k(\mathbf{H})$, $(0, 1) \in S$, and therefore $(x, y) \rightarrow (0, 1) = (x \Rightarrow 0, x \wedge 1) = (x^*, x) = (0, x) \in S$. Hence, $x \in E$ because $x = x \vee 0$, and we conclude that E satisfies (IF2).

Suppose now that for $z, t \in H$ we have $z \vee t \in E$,

Since $z \vee t \in E$, there exists a pair $(a, b) \in S$ such that $z \vee t = a \vee b$. But S is a subalgebra of $\mathbf{V}_k(\mathbf{H})$, so $\sim(a, b) = (b, a) \in S$, and therefore $(a, b) \cup (b, a) = (a \vee b, a \wedge b) = (a \vee b, 0) \in S$. This means that $(z \vee t, 0) \in S$, and hence

$$\sim(z \vee t, 0) = (0, z \vee t) \in S. \tag{3.1}$$

Using the hypothesis $\pi_1(S) = H$, and $x, z, t \in H$, it follows that there exist some $x', z', t' \in H$ such that

$(x, x'), (z, z'), (t, t') \in S$. This implies that the next term belongs to S :

$$\begin{aligned} & ((x, x') \rightarrow (z, z')) \cup ((x, x') \cap (t, t')) \\ &= (x \Rightarrow z, x \wedge z') \cup (x \wedge t, x' \vee t') \\ &= ((x \Rightarrow z) \vee (x \wedge t), (x \wedge z') \wedge (x' \vee t')). \end{aligned} \tag{3.2}$$

Since $(x \wedge z') \wedge (x' \vee t') = (x \wedge z' \wedge x') \vee (x \wedge z' \wedge t') = 0 \vee (x \wedge z' \wedge t') = x \wedge z' \wedge t'$, (3.2) yields

$$((x \Rightarrow z) \vee (x \wedge t), x \wedge z' \wedge t') \in S. \tag{3.3}$$

Combining (3.1) with (3.3), we get

$$\begin{aligned} & ((x \Rightarrow z) \vee (x \wedge t), x \wedge z' \wedge t') \cup (0, z \vee t) \\ &= ((x \Rightarrow z) \vee (x \wedge t), x \wedge z' \wedge t' \wedge (z \vee t)) \in S. \end{aligned} \tag{3.4}$$

Since $x \wedge z' \wedge t' \wedge (z \vee t) = (x \wedge z' \wedge t' \wedge z) \wedge (x \wedge z' \wedge t' \wedge t) = (x \wedge t' \wedge (z \wedge z')) \vee (x \wedge z \wedge (t \wedge t')) = 0 \vee 0 = 0$, from (3.4), it follows that

$$((x \Rightarrow z) \vee (x \wedge t), 0) \in S.$$

By the definition of E , it follows that

$$(x \Rightarrow z) \vee (x \wedge t) \vee 0 = (x \Rightarrow z) \vee (x \wedge t) \in E.$$

Hence, E is an i -filter.

Let us prove now that $N(H, E) = S$. It is clear that $S \subseteq N(H, E)$. If we have $(a, b) \in N(H, E)$, then $a \wedge b = 0$, and $a \vee b \in E$, so there exists $(c, d) \in S$ such that $c \vee d = a \vee b$. Then, as before $(c, d) \cup (d, c) = (c \vee d, 0) = (a \vee b, 0) \in S$. On the other hand, since $\pi_1(S) = H$, there exists some $b' \in H$ such that $(b, b') \in S$, and therefore $(b, b') \rightarrow (0, 1) = (b^*, b) \in S$. By properties of the pseudocomplement, $a \wedge b = 0$ implies that $a \leq b^*$ and hence $(a \vee b) \wedge b^* = (a \wedge b^*) \vee (b \wedge b^*) = a \vee 0 = a$, so we can conclude that $(a \vee b, 0) \cap (b^*, b) = ((a \vee b) \wedge b^*, 0 \vee b) = (a, b) \in S$. \square

Theorem 3.6 Every semi-Nelson algebra \mathbf{A} is isomorphic to an algebra of the form $\mathbf{N}(\mathbf{H}, \mathbf{F})$, for some semi-Heyting algebra \mathbf{H} and some i -filter F of \mathbf{H} .

Proof By Theorem 2.12, every semi-Nelson algebra \mathbf{A} can be obtained as a subalgebra of $\mathbf{V}_k(\mathbf{sH}(\mathbf{A}))$, and clearly $\pi_1(\mathbf{V}_k(\mathbf{sH}(\mathbf{A}))) = \mathbf{sH}(\mathbf{A})$, so \mathbf{A} is isomorphic to $\mathbf{N}(\mathbf{sH}(\mathbf{A}), \mathbf{E})$, where E is the i -filter of Lemma 3.5. \square

Much in the same manner as for the case of Heyting algebras, for $\mathbf{H} \in \mathcal{SH}$, the i -filters E such that $\pi_1(N(H, E)) = H$ are exactly the ones that contain the dense elements of H :

Lemma 3.7 If $\mathbf{H} \in \mathcal{SH}$, and F is an i -filter of \mathbf{H} , then $\pi_1(N(H, F)) = H$.

Proof If $x \in H$, by Lemma 2.7, we have that $x \vee x^* \in \mathcal{D}(H) \subseteq F$. Also, $x \wedge x^* = 0$ is valid, so $(x, x^*) \in N(H, F)$ and $x \in \pi_1(N(H, F))$. \square

3.1 Categorical equivalence

We define now the category \mathbf{sHF} , which has as objects pairs (\mathbf{H}, F) , where $\mathbf{H} \in \mathcal{SH}$, and F is an i -filter of H , and as morphisms, functions $f : (\mathbf{H}, F) \rightarrow (\mathbf{H}', F')$ such that $f : H \rightarrow H'$ is a homomorphism of semi-Heyting algebras and $f(F) \subseteq F'$.

Consider now the category \mathbf{sN} of semi-Nelson algebras and their morphisms; our objective now is to establish an equivalence between these categories. For this, we define functors $\alpha : \mathbf{sHF} \rightarrow \mathbf{sN}$ and $\beta : \mathbf{sN} \rightarrow \mathbf{sHF}$.

Proposition 3.8 The application $\alpha : \mathbf{sHF} \rightarrow \mathbf{sN}$ defined on the objects of \mathbf{sHF} by $\alpha((\mathbf{H}, F)) = \mathbf{N}(\mathbf{H}, \mathbf{F})$, and on morphisms by $\alpha(f)(a, b) = (f(a), f(b))$ for all $(a, b) \in N(H, F)$, is a functor from \mathbf{sHF} to \mathbf{sN} .

Proof Let us check that α is well defined. If (\mathbf{H}, F) is an object of \mathbf{sHF} , then $\alpha((\mathbf{H}, F)) = \mathbf{N}(\mathbf{H}, \mathbf{F})$ is a semi-Nelson algebra due to Lemma 3.4.

Now we take a morphism $f : (\mathbf{H}, F) \rightarrow (\mathbf{H}', F')$. Since f is a Heyting homomorphism, we have by straightforward calculations that: $\alpha(f)((a, b) \rightarrow (c, d)) = (f(a \Rightarrow c), f(a \wedge d)) = (f(a), f(b)) \rightarrow ((f(c), f(d))) = \alpha(f)(a, b) \rightarrow \alpha(f)(c, d)$, and similarly for the rest of the operations.

We denote by Id_X the identity morphism for an object X in a category. Then, for all $(a, b) \in N(H, F)$, $(a, b) = (Id_{(H, F)}(a), Id_{(H, F)}(b))$, therefore $\alpha(Id_{(\mathbf{H}, F)}) = Id_{\alpha(\mathbf{H}, F)}$.

Finally, for any two given morphisms $f : (\mathbf{H}', F') \rightarrow (\mathbf{H}'', F'')$, and $g : (\mathbf{H}, F) \rightarrow (\mathbf{H}', F')$, we have that for all $(a, b) \in N(H, F)$, $\alpha(f \circ g)(a, b) = ((f \circ g)(a), (f \circ g)(b)) = (f(g(a)), f(g(b))) = \alpha(f)((g(a), g(b))) = ((\alpha(f)) \circ (\alpha(g)))(a, b)$. \square

Definition 3.9 Let $A \in \mathcal{SN}$. We say that an element $a \in A$ is positive if $\sim a \leq a$. We denote by A^+ the set of positive elements of A .

Observation 3.10 If $A \in \mathcal{SN}$, then $A^+ = \{x \vee \sim x : x \in A\}$. Indeed, if $x \in A^+$, then $\sim x \leq x$, so $x = x \vee \sim x$. On the other hand, if $y = x \vee \sim x$, by (SN4) and (SN5) it follows that $y = x \vee \sim x \geq x \wedge \sim x = \sim(\sim x \vee x) = \sim y$.

We write $\llbracket A^+ \rrbracket$ for the set $\{\llbracket a \rrbracket : a \in A^+\}$.

Lemma 3.11 If $\mathbf{A} \in \mathcal{SN}$, then $\llbracket A^+ \rrbracket$ is an i -filter of $\mathbf{sH}(\mathbf{A})$.

Proof By Theorem 2.12, there exists a subalgebra \mathbf{S} of $\mathbf{V}_k(\mathbf{sH}(\mathbf{A}))$ such that \mathbf{A} is isomorphic to \mathbf{S} , and the isomorphism is given by $h(a) = (\llbracket a \rrbracket, \llbracket \sim a \rrbracket)$ for all $a \in A$.

Therefore, $S = \{\llbracket a \rrbracket, \llbracket \sim a \rrbracket : a \in A\}$. Then, it is immediate that $\pi_1(S) = \mathbf{sH}(A)$.

By Lemma 3.5, $S = N(\mathbf{sH}(A), E)$, where $E = \{\llbracket a \rrbracket \vee \llbracket \sim a \rrbracket : a \in A\}$ is an i -filter of $\mathbf{sH}(A)$. But $E = \llbracket A^+ \rrbracket$, due to observation 3.10, which completes the proof. \square

Proposition 3.12 *The application $\beta : \mathbf{sN} \rightarrow \mathbf{sHF}$ defined on the objects of \mathbf{sN} by $\beta(\mathbf{A}) = (\mathbf{sH}(\mathbf{A}), \llbracket A^+ \rrbracket)$, and on morphisms $f : A \rightarrow A'$ by $\beta(f)(\llbracket a \rrbracket) = \llbracket f(a) \rrbracket$ for all $a \in A$, is a functor from \mathbf{sN} to \mathbf{sHF} .*

Proof Let us check that β is well defined. For an object \mathbf{A} of \mathbf{sN} , $\mathbf{sH}(\mathbf{A})$ is a semi-Heyting algebra, and by Lemma 3.11 we have that $\llbracket A^+ \rrbracket$ is an i -filter of $\mathbf{sH}(\mathbf{A})$. Therefore, $(\mathbf{sH}(\mathbf{A}), \llbracket A^+ \rrbracket)$ is an object in \mathbf{sHF} .

For any semi-Nelson morphism $f : A \rightarrow A'$, we calculate: $\beta(f)(\llbracket a \rightarrow b \rrbracket) = \llbracket f(a \rightarrow b) \rrbracket = \llbracket f(a) \rightarrow f(b) \rrbracket = \llbracket f(a) \rrbracket \Rightarrow \llbracket f(b) \rrbracket = \beta(f)(\llbracket a \rrbracket) \Rightarrow \beta(f)(\llbracket b \rrbracket)$, and similarly for the other operations.

Also, if $\llbracket x \rrbracket \in \llbracket A^+ \rrbracket$, then by observation 3.10 we can write $\llbracket x \rrbracket = \llbracket y \vee \sim y \rrbracket$, for some $y \in A$, and so $\beta(f)(\llbracket x \rrbracket) = \beta(f)(\llbracket y \vee \sim y \rrbracket) = \beta(f)(\llbracket y \rrbracket) \cup \beta(f)(\llbracket \sim y \rrbracket) = \llbracket f(y) \rrbracket \cup \llbracket f(\sim y) \rrbracket = \llbracket f(y) \rrbracket \cup \llbracket \sim f(y) \rrbracket = \llbracket f(y) \vee \sim f(y) \rrbracket$. Since $f(y) \in f(A) \subseteq A'$, it follows that $\beta(f)(\llbracket x \rrbracket) \in \llbracket A'^+ \rrbracket$. This shows that $\beta(f)(\llbracket A^+ \rrbracket) \subseteq \llbracket A'^+ \rrbracket$, so $\beta(f)$ is a morphism in \mathbf{sHF} .

It is straightforward to check that β preserves identities and compositions, so it is indeed a functor. \square

The connection between theorems 3.8 and 3.12 is the following:

Theorem 3.13 *The functors α and β establish an equivalence between the categories \mathbf{sHF} and \mathbf{sN} .*

Proof Notice that $\beta\alpha(\mathbf{A}, F) = \beta(\mathbf{N}(\mathbf{A}, \mathbf{F})) = (\mathbf{sH}(\mathbf{N}(\mathbf{A}, \mathbf{F})), \llbracket N(A, F)^+ \rrbracket)$, with $\mathbf{sH}(\mathbf{N}(\mathbf{A}, \mathbf{F})) = \{\llbracket (x, y) \rrbracket : (x, y) \in N(A, F)\} = \{\llbracket (x, y) \rrbracket : x, y \in A, x \wedge y = 0, x \vee y \in F\}$.

We define now $\eta_A(x) = \llbracket (x, x^*) \rrbracket$, for each object \mathbf{A} of \mathbf{sHF} and for all $x \in A$, and we prove that $\eta = \{\eta_A\}$ is a natural isomorphism from $ids_{\mathbf{sHF}}$ to $\beta\alpha$.

η_A is well defined: Since $x \wedge x^* = 0$ and also $\mathcal{D}(A) \subseteq F$ it follows that $x \vee x^* \in \mathcal{D}(A) \subseteq F$. Hence, $(x, x^*) \in N(A, F)$.

η_A is surjective: If $\llbracket (x, y) \rrbracket \in \beta\alpha(A, F)$, then $\llbracket (x, y) \rrbracket = \llbracket (x, x^*) \rrbracket$. Indeed, since $(x, y) \rightarrow (x, x^*) = (x \Rightarrow x, x \wedge x^*) = (1, 0)$ and $(x, x^*) \rightarrow (x, y) = (x \Rightarrow x, x \wedge y) = (1, 0)$, so by definition we have that $\llbracket (x, y) \rrbracket = \llbracket (x, x^*) \rrbracket$. Therefore, $\eta_A(x) = \llbracket (x, y) \rrbracket$.

η_A is injective: If $\llbracket (x, x^*) \rrbracket = \llbracket (y, y^*) \rrbracket$, then $(x, x^*) \rightarrow (y, y^*) = (x \Rightarrow y, x \wedge y^*) = (1, 0)$. Therefore, from $x \Rightarrow y = 1$, using properties of semi-Heyting algebras we obtain $x \leq y$. Analogously, $(y, y^*) \rightarrow (x, x^*) = (y \Rightarrow x, y \wedge x^*) = (1, 0)$ implies $y \leq x$.

η_A is a \mathbf{sHF} -morphism:

- $\eta_A(x) \cup \eta_A(y) = \llbracket (x, x^*) \rrbracket \cup \llbracket (y, y^*) \rrbracket = \llbracket (x, x^*) \sqcup (y, y^*) \rrbracket = \llbracket (x \vee y, x^* \wedge y^*) \rrbracket = \llbracket (x \vee y, (x \vee y)^*) \rrbracket = \eta_A(x \vee y)$.
- $\eta_A(x) \cap \eta_A(y) = \llbracket (x, x^*) \rrbracket \cap \llbracket (y, y^*) \rrbracket = \llbracket (x, x^*) \sqcap (y, y^*) \rrbracket = \llbracket (x \wedge y, x^* \vee y^*) \rrbracket$. Also, $\eta_A(x \wedge y) = \llbracket (x \wedge y, (x \wedge y)^*) \rrbracket$. Since $(x \wedge y, x^* \vee y^*) \rightarrow (x \wedge y, (x \wedge y)^*) = ((x \wedge y) \Rightarrow (x \wedge y), (x \wedge y) \wedge (x \wedge y)^*) = (1, 0)$ and $(x \wedge y, (x \wedge y)^*) \rightarrow (x \wedge y, x^* \vee y^*) = ((x \wedge y) \Rightarrow (x \wedge y), (x \wedge y) \wedge (x^* \vee y^*)) = (1, (x \wedge y \wedge x^*) \vee (x \wedge y \wedge y^*)) = (1, 0)$, we have that $\eta_A(x) \cap \eta_A(y) = \eta_A(x \wedge y)$.
- $\eta_A(x) \Rightarrow \eta_A(y) = \llbracket (x, x^*) \rrbracket \Rightarrow \llbracket (y, y^*) \rrbracket = \llbracket (x, x^*) \Rightarrow (y, y^*) \rrbracket = \llbracket (x \Rightarrow y, x \wedge y^*) \rrbracket$. Also, $\eta_A(x \Rightarrow y) = \llbracket (x \Rightarrow y, (x \Rightarrow y)^*) \rrbracket$. Since $(x \Rightarrow y, x \wedge y^*) \rightarrow (x \Rightarrow y, (x \Rightarrow y)^*) = ((x \Rightarrow y) \Rightarrow (x \Rightarrow y), (x \Rightarrow y) \wedge (x \Rightarrow y)^*) = (1, 0)$ and $(x \Rightarrow y, (x \Rightarrow y)^*) \rightarrow (x \Rightarrow y, x \wedge y^*) = ((x \Rightarrow y) \Rightarrow (x \Rightarrow y), (x \Rightarrow y) \wedge (x \wedge y^*)) \stackrel{(SH_2)}{=} (1, x \wedge (y \wedge y^*)) = (1, 0)$, we have that $\llbracket (x \Rightarrow y, x \wedge y^*) \rrbracket = \llbracket (x \Rightarrow y, (x \Rightarrow y)^*) \rrbracket$. Therefore, we conclude that $\eta_A(x) \Rightarrow \eta_A(y) = \eta_A(x \Rightarrow y)$.
- $\eta_A(F) \subseteq \llbracket N(A, F)^+ \rrbracket$
If $y \in \eta_A(F)$, then $y = \eta_A(x)$, for some $x \in F$. Hence, $y = \llbracket (x, x^*) \rrbracket$, with $x \in F$.
We observe that for $x \in F$, we have that $(x, 0) \in N(A, F)$. Besides, $\llbracket (x, x^*) \rrbracket = \llbracket (x, 0) \rrbracket$ because $(x, 0) \rightarrow (x, x^*) = (x \Rightarrow x, x \wedge x^*) = (1, 0)$ and $(x, x^*) \rightarrow (x, 0) = (x \Rightarrow x, x \wedge 0) = (1, 0)$.
Therefore, we can write $y = \llbracket (x, 0) \rrbracket$. Now, since $\sim(x, 0) = (0, x) \leq (x, 0)$, it follows that $(x, 0) \in N(A, F)^+$, and hence $y \in \llbracket N(A, F)^+ \rrbracket$.

Finally, we check the naturality of η , that is, the following diagram commutes:

$$\begin{array}{ccc}
 A & \xrightarrow{\eta_A} & \beta\alpha(A) \\
 f \downarrow & & \downarrow \beta\alpha(f) \\
 A' & \xrightarrow{\eta_{A'}} & \beta\alpha(A')
 \end{array}$$

Notice that since η_A is an isomorphism, $\eta_A^{-1}(\llbracket (x, y) \rrbracket) = x$ for every $x \in A$. Then we have

$$\begin{aligned}
 (\eta_{A'}^{-1} \circ \beta\alpha(f) \circ \eta_A)(x) &= \eta_{A'}^{-1}(\beta\alpha(f)(\llbracket (x, 0) \rrbracket)) \\
 &= \eta_{A'}^{-1}(\llbracket \alpha(f)(x, 0) \rrbracket) = \eta_{A'}^{-1}(\llbracket f(x), 0 \rrbracket) = f(x).
 \end{aligned}$$

Thus $f = \eta_{A'}^{-1} \circ \beta\alpha(f) \circ \eta_A$, and therefore $\eta_{A'} \circ f = \beta\alpha(f) \circ \eta_A$.

Now we consider the functor $\alpha\beta$, which transforms every object \mathbf{A} of \mathbf{sN} into $\mathbf{N}(\mathbf{sH}(\mathbf{A}), \llbracket A^+ \rrbracket)$. On a morphism $f : A \rightarrow A'$ it is defined by $\alpha\beta(f)(\llbracket x \rrbracket, \llbracket y \rrbracket) =$

$(\llbracket f(x) \rrbracket, \llbracket f(y) \rrbracket)$. We define for each object \mathbf{A} of \mathbf{sN} , $\delta_A : A \rightarrow \mathbf{N}(\mathbf{sH}(\mathbf{A}), \llbracket \mathbf{A}^+ \rrbracket)$ by $\delta_A(x) = (\llbracket x \rrbracket, \llbracket \sim x \rrbracket)$.

δ_A is well defined: On the one hand, $\llbracket x \rrbracket \cap \llbracket \sim x \rrbracket = \llbracket x \wedge \sim x \rrbracket = \llbracket 0 \rrbracket$, because $(x \wedge \sim x) \rightarrow 0 = (x \wedge \sim x) \rightarrow (x \wedge \sim x \wedge 0) = (x \wedge \sim x) \rightarrow_N 0 = 1$ and $0 \rightarrow_N (x \wedge \sim x) = 1$ by well-known properties of Nelson algebras. On the other hand, using (SN5) and De Morgan properties we have $\sim(x \vee \sim x) = \sim x \wedge x \leq x \vee \sim x$, so $x \vee \sim x \in \mathbf{A}^+$. Therefore $\llbracket x \rrbracket \cup \llbracket \sim x \rrbracket = \llbracket x \vee \sim x \rrbracket \in \llbracket \mathbf{A}^+ \rrbracket$.


δ_A is surjective: If $(\llbracket x \rrbracket, \llbracket y \rrbracket) \in \alpha\beta(A, F)$, then $(\llbracket x \rrbracket, \llbracket y \rrbracket) = (\llbracket x \rrbracket, \llbracket \sim x \rrbracket)$. Indeed, this is a consequence of $(\llbracket x \rrbracket, \llbracket y \rrbracket) \rightarrow (\llbracket x \rrbracket, \llbracket \sim x \rrbracket) = (\llbracket x \rrbracket \Rightarrow \llbracket x \rrbracket, \llbracket x \rrbracket \cap \llbracket \sim x \rrbracket) = (\llbracket 1 \rrbracket, \llbracket x \wedge \sim x \rrbracket) = (\llbracket 1 \rrbracket, \llbracket 0 \rrbracket)$ and $(\llbracket x \rrbracket, \llbracket \sim x \rrbracket) \rightarrow (\llbracket x \rrbracket, \llbracket y \rrbracket) = (\llbracket x \rrbracket \Rightarrow \llbracket x \rrbracket, \llbracket x \rrbracket \cap \llbracket y \rrbracket) = (\llbracket 1 \rrbracket, \llbracket 0 \rrbracket)$. Therefore $\delta_A(x) = (\llbracket x \rrbracket, \llbracket y \rrbracket)$.

δ_A is injective: If $(\llbracket x \rrbracket, \llbracket \sim x \rrbracket) = (\llbracket y \rrbracket, \llbracket \sim y \rrbracket)$, then $\llbracket x \rrbracket = \llbracket y \rrbracket$ and $\llbracket \sim x \rrbracket = \llbracket \sim y \rrbracket$, so we have the following: $x \rightarrow y = 1, y \rightarrow x = 1, \sim x \rightarrow \sim y = 1, \sim y \rightarrow \sim x = 1$. By proposition 2.11 (1), we have that $x \rightarrow_N y = 1, y \rightarrow_N x = 1, \sim x \rightarrow_N \sim y = 1$, and $\sim y \rightarrow_N \sim x = 1$. These four equalities imply that $x = y$ (see Viglizzo 1999, (1.15)).

δ_A is a \mathbf{sN} -morphism:


- $\delta_A(x) \sqcup \delta_A(y) = (\llbracket x \rrbracket, \llbracket \sim x \rrbracket) \sqcup (\llbracket y \rrbracket, \llbracket \sim y \rrbracket) = (\llbracket x \rrbracket \cup \llbracket y \rrbracket, \llbracket \sim x \rrbracket \cap \llbracket \sim y \rrbracket) = (\llbracket x \vee y \rrbracket, \llbracket \sim(x \vee y) \rrbracket) = \delta_A(x \vee y)$. In a similar manner, we can prove that $\delta_A(x) \sqcap \delta_A(y) = \delta_A(x \wedge y)$.
- $\sim \delta_A(x) = \sim (\llbracket x \rrbracket, \llbracket \sim x \rrbracket) = (\llbracket \sim x \rrbracket, \llbracket x \rrbracket) = \delta_A(\sim x)$.
- $\delta_A(x) \rightarrow \delta_A(y) = (\llbracket x \rrbracket, \llbracket \sim x \rrbracket) \rightarrow (\llbracket y \rrbracket, \llbracket \sim y \rrbracket) = (\llbracket x \rrbracket \Rightarrow \llbracket y \rrbracket, \llbracket x \rrbracket \cap \llbracket \sim y \rrbracket) = (\llbracket x \rightarrow y \rrbracket, \llbracket x \wedge \sim y \rrbracket)$. By (SN10) and (SN11), we see that $\llbracket x \wedge \sim y \rrbracket = \llbracket \sim(x \rightarrow y) \rrbracket$. Hence $\delta_A(x) \rightarrow \delta_A(y) = (\llbracket x \rightarrow y \rrbracket, \llbracket x \wedge \sim y \rrbracket) = (\llbracket x \rightarrow y \rrbracket, \llbracket \sim(x \rightarrow y) \rrbracket) = \delta_A(x \rightarrow y)$.

Finally, δ is a natural transformation: to see that this diagram commutes,



we calculate $\alpha\beta(f) \circ \delta_A(x) = \alpha\beta(f)(\llbracket x \rrbracket, \llbracket \sim x \rrbracket) = (\llbracket f(x) \rrbracket, \llbracket f(\sim x) \rrbracket) = (\llbracket f(x) \rrbracket, \llbracket \sim f(x) \rrbracket) = \delta_{A'} \circ f(x)$. \square


Example 3.14 Consider the semi-Nelson algebra \mathbf{A} , with Hasse diagram indicated below, in which the operations \rightarrow and \sim are given by the following tables:



\rightarrow	0	a	b	c	d	e	f	g	1
0	1	1	1	1	g	1	g	1	g
a	1	1	1	1	g	1	g	1	g
b	1	1	1	1	g	1	g	1	g
c	1	1	1	1	g	1	g	1	g
d	e	g	e	g	1	e	1	g	1
e	d	d	f	f	a	1	c	1	g
f	e	g	e	g	1	e	1	g	1
g	d	d	f	f	a	1	c	1	g
1	0	a	b	c	d	e	f	g	1

x	0	a	b	c	d	e	f	g	1
$\sim x$	1	g	f	c	e	d	b	a	0

$S = \{0, a, d, e, g, 1\}$ is the universe of a subalgebra \mathbf{S} of \mathbf{A} . Both \mathbf{S} and \mathbf{A} have the same quotient semi-Nelson algebra \mathbf{H} indicated below:



\Rightarrow	[0]	[d]	[e]	[1]
[0]	[1]	[e]	[1]	[e]
[d]	[e]	[1]	[e]	[1]
[e]	[d]	[0]	[1]	[e]
[1]	[0]	[d]	[e]	[1]

$E = \{[e], [1]\}$ is an i -filter of \mathbf{H} , and $\mathbf{N}(\mathbf{H}, E)$ is isomorphic to \mathbf{S} , while $\mathbf{N}(\mathbf{H}, \mathbf{H}) = \mathbf{V}_k(\mathbf{H})$ is isomorphic to \mathbf{A} .

4 Dually hemimorphic semi-Nelson algebras

In this section, we will show some results about the variety of dually hemimorphic semi-Heyting and semi-Nelson

algebras, which were presented in definitions 2.8 and 2.9, respectively. We omit some proofs of the following results, since they can be found in Cornejo and San Martín (2020).

Lemma 4.1 *Let $\mathbf{A} \in \mathcal{DSN}$. If we consider the equivalence relation \equiv defined in Sect. 2.2, then that relation is compatible with respect of the operation $'$. Hence, \equiv is a congruence in \mathcal{DSN} .*

In view of the previous lemma, we can define

$$\llbracket x \rrbracket^\dagger = \llbracket x' \rrbracket$$

and moreover, it can be proved that $\mathbf{dsH}(A) = \langle A / \equiv; \cap, \cup, \Rightarrow, \dagger, \perp, \top \rangle \in \mathcal{DSH}$.

Definition 4.2 Let $\mathbf{A} = \langle A; \cap, \cup, \Rightarrow, \dagger, \perp, \top \rangle \in \mathcal{DSH}$. For $(a, b) \in V_k(A)$, we define the following unary operation on $V_k(A)$:

$$(a, b)' = (a^\dagger, b \cap (a^\dagger \Rightarrow a)).$$

The operation is well defined. Indeed, using (SH2) we have $a^\dagger \cap b \cap (a^\dagger \Rightarrow a) = b \cap a^\dagger \cap (a^\dagger \Rightarrow a) = b \cap a^\dagger \cap a = 0$.

Theorem 4.3 *If $\mathbf{A} \in \mathcal{DSH}$, then the system $\mathbf{V}_k(\mathbf{A}) = \langle V_k(A); \cap, \cup, \Rightarrow, \sim, ', \top \rangle$ is a dually hemimorphic semi-Nelson algebra.*

In a similar manner as in sect. 2.1, we have the following representations:

Theorem 4.4 Cornejo and San Martín (2020) *If $\mathbf{A} \in \mathcal{DSH}$, then \mathbf{A} is isomorphic to $\mathbf{dsH}(\mathbf{V}_k(\mathbf{A}))$. Also, if $\mathbf{A} \in \mathcal{DSN}$, then \mathbf{A} is isomorphic to a subalgebra of $\mathbf{V}_k(\mathbf{dsH}(\mathbf{A}))$.*

Now we will generalize the results of sect. 3. In Cornejo and San Martín (2018), the authors introduced the category \mathcal{SN}_c of centered semi-Nelson algebras whose objects are algebras $\mathbf{A} = \langle A; \wedge, \vee, \rightarrow, \sim, c, 1 \rangle$ of type $(2, 2, 2, 1, 0, 0)$ such that $c \approx \sim c$, and the morphisms are the algebra homomorphisms. The element c is called a *center* of the algebra A , and it is necessarily unique. We can expand the category \mathcal{SN}_c to \mathcal{DSN}_c of dually hemimorphic-centered semi-Nelson algebras whose objects are algebras $\mathbf{A} = \langle A; \wedge, \vee, \rightarrow, \sim, ', c, 1 \rangle$ of type $(2, 2, 2, 1, 1, 0, 0)$ such that $c \approx \sim c$, and the morphisms are the algebra homomorphisms.

We will prove that the categories \mathcal{DSN} and \mathcal{DSN}_c actually coincide.

Proposition 4.5 *If $\mathbf{A} \in \mathcal{DSN}$, then \mathbf{A} is centered, with center $1'$.*

Proof We want to prove that $\sim 1' = 1'$, or equivalently, the following conditions hold simultaneously:

$$\begin{aligned} &\sim 1' \rightarrow 1' = 1, \quad 1' \rightarrow \sim 1' = 1, \\ &\sim 1' \rightarrow \sim (\sim 1') = 1, \quad \sim (\sim 1') \rightarrow \sim 1' = 1. \end{aligned} \tag{4.1}$$

By Proposition 2.11 (1), (4.1) is equivalent to

$$\begin{aligned} &\sim 1' \rightarrow_N 1' = 1, \quad 1' \rightarrow_N \sim 1' = 1, \\ &\sim 1' \rightarrow_N \sim (\sim 1') = 1, \quad \sim (\sim 1') \rightarrow_N \sim 1' = 1, \end{aligned}$$

which is equivalent to show that

$$\sim 1' \rightarrow_N 1' = 1, \tag{4.2}$$

$$1' \rightarrow_N \sim 1' = 1. \tag{4.3}$$

By (DSN4), (DSN5), and proposition 1, we have that $\sim 1' \rightarrow_N (\sim 1 \wedge (1' \rightarrow 1)) = 1$, that is $(1) \sim 1' \rightarrow_N 0 = 1$. Since in every Nelson algebra $0 \rightarrow_N x = x$ holds for all x , it follows that $(2) 0 \rightarrow_N 1' = 1'$. By (1), (2) and proposition 2.11 (2), we have that $\sim 1' \rightarrow_N 1' = 1$, which proves (4.2).

By (DSN2), we have $(3) 1 = 1' \rightarrow \sim 1 = 1' \rightarrow 0 = 1' \rightarrow (1' \wedge 0) = 1' \rightarrow_N 0$. Using again that $0 \rightarrow_N 1' = 1'$, and proposition 2.11 (2), gives us $\sim 1' \rightarrow_N 1 = 1$, which proves (4.3). \square

Proposition 4.5 shows us that the category \mathcal{DSN} can be viewed as the category \mathcal{DSN}_c with a distinguished element c , but despite the change of the language, both categories are essentially the same.

It was proved in Cornejo and San Martín (2018) that there is a categorial equivalence between semi-Heyting algebras and centered semi-Nelson algebras. Moreover, the function h defined in the proof of Theorem 4.4 is an isomorphism. This gives us the following:

Theorem 4.6 *If $\mathbf{A} \in \mathcal{DSN}$, then \mathbf{A} is isomorphic to $\mathbf{V}_k(\mathbf{dsH}(\mathbf{A}))$.*

Observation 4.7 *Another way of proving Theorem 4.6 is the following: if we consider the structure $\mathbf{N}(\mathbf{H}, \mathbf{E})$ as in Lemma 3.5, and take $\mathbf{H} \in \mathcal{DSH}$, we have no guarantee that this set is going to be closed under the operation \dagger . If instead we consider a subalgebra \mathbf{S} such that $\pi_1(S) = H$, then the proof indicated in the lemma still holds for the variety \mathcal{DSH} , since it does not use the operations $'$ or \dagger . Hence, the sets S and $N(H, E)$ are equal, so $N(H, E)$ inherits the operations from \mathbf{S} . Indeed, due to Theorem 4.4, we know that the function $h : A \rightarrow \mathbf{V}_k(\mathbf{dsH}(A))$ defined by $h(a) = (\llbracket a \rrbracket, \llbracket \sim a \rrbracket)$ is a monomorphism. Hence, $A \cong h(A)$. Let us take then the subalgebra $\mathbf{S} = h(\mathbf{A})$ of $\mathbf{V}_k(\mathbf{dsH}(\mathbf{A}))$. It is immediate that $\pi_1(S) = \mathbf{dsH}(A)$. Therefore, by Lemma 3.5, $h(\mathbf{A}) = \mathbf{N}(\mathbf{dsH}(\mathbf{A}), \mathbf{E})$ for $E = \{\llbracket x \rrbracket \in \mathbf{dsH}(A) : \llbracket x \rrbracket = \llbracket a \rrbracket \vee \llbracket b \rrbracket, (\llbracket a \rrbracket, \llbracket b \rrbracket) \in S\}$.*

Since \mathbf{S} is a subalgebra of $\mathbf{V}_k(\mathbf{dsH}(\mathbf{A}))$, we have that $(\llbracket 1 \rrbracket, \llbracket 0 \rrbracket) \in S$, so the center $(\llbracket 0 \rrbracket, \llbracket 0 \rrbracket) = (\llbracket 1 \rrbracket, \llbracket 0 \rrbracket)'$ is an element of S . Then $\llbracket 0 \rrbracket = \llbracket 0 \rrbracket \vee \llbracket 0 \rrbracket \in E$. Since E is a filter, this shows that $E = \mathbf{dsH}(A)$.

We have proved that $S = N(\mathbf{dsH}(A), \mathbf{dsH}(A)) = \mathbf{V}_k(\mathbf{dsH}(A))$, and therefore h is an isomorphism.

Example 4.8 Consider the semi-Nelson algebras \mathbf{A} and its subalgebra \mathbf{S} from Example 3.14. We know by the results above that \mathbf{S} cannot be expanded by adding a unary operation $'$ in such a way that the system $\mathbf{B} = \langle B; \wedge, \vee, \rightarrow, \sim, ', 1 \rangle$ becomes a dually hemimorphic semi-Nelson algebra, since \mathbf{S} has no center.

On the other hand, any operation $'$ that makes $\mathbf{sH}(\mathbf{A})$ into a dually hemimorphic semi-Heyting algebra yields an operation \dagger that will turn A into a dually hemimorphic semi-Nelson algebra.

Declarations

Conflict of interest The authors declare that they have no conflict of interest

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

Data sharing not applicable to this article as no datasets were generated or analyzed during the current study.

References

- Abad M, Comejo JM, Varela JPD (2013) Semi-Heyting algebras term-equivalent to Godel algebras. *Order*, 30(2): 625-642
- Balbes R, Dwinger P (1974) *Distributive lattices*. University of Missouri Press, Columbia, Mo

- Cornejo JM, San Martín HJ (2018) A categorial equivalence between semi-Heyting algebras and centered semi-Nelson algebras. *Logic J IGPL* 26(4):408–428
- Cornejo JM, San Martín HJ (2020) Dually hemimorphic semi-Nelson algebras. *Log J IGPL* 28(3):316–334
- Cornejo JM, Viglizzo I (2018) Semi-Nelson algebras. *Order* 35(1):23–45
- Kalman JA (1958) Lattices with involution. *Trans Amer Math Soc* 87:485–491
- Monteiro A, Monteiro L (1996) Axiomes indépendants pour les algèbres de Nelson, de Łukasiewicz trivalentes, de De Morgan et de Kleene. In *Unpublished papers, I*, *Notas Lógica Mat* 40:13
- Monteiro A, Monteiro L (1996) Axiomes indépendants pour les algèbres de Nelson, de Łukasiewicz trivalentes, de De Morgan et de Kleene. In *Unpublished papers I*, *Notas Lógica Mat.* 40:13
- Sankappanavar HP (1985) Semi-Heyting algebras. *Amer Math Soc Abstracts*, 13
- Sankappanavar HP (2008) Semi-Heyting algebras: an abstraction from Heyting algebras. In *Proceedings of the 9th “Dr. Antonio A. R. Monteiro” Congress (Spanish)*, *Actas Congr. “Dr. Antonio A. R. Monteiro”*, pages 33–66, Bahía Blanca, 2008. Univ Nac del Sur
- Sankappanavar HP (2011) Expansions of semi-Heyting algebras I: discriminator varieties. *Studia Logica* 98(1–2):27–81
- Sendlewski A (1984) Some investigations of varieties of \mathcal{N} -lattices. *Studia Logica* 43(3):257–280
- Sendlewski A (1990) Nelson algebras through Heyting ones. I *Studia Logica* 49(1):105–126
- Vakarelov D (1977) Notes on \mathcal{N} -lattices and constructive logic with strong negation. *Studia Logica* 36(1–2):109–125
- Viglizzo I (1999) *Álgebras de Nelson*. Instituto de Matemática de Bahía Blanca, Universidad Nacional del Sur, 1999. Magister dissertation in Mathematics, Universidad Nacional del Sur, Bahía Blanca, available at <https://sites.google.com/site/viglizzo/viglizzo99nelson>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.