



Published in final edited form as:

Med Sci Sports Exerc. 2014 December ; 46(12): 2195–2201. doi:10.1249/MSS.0000000000000360.

Cancer-Related Impairments Influence Physical Activity in Uterine Cancer Survivors

Sean M. Hammer¹, Justin C. Brown¹, Saya Segal², Christina S. Chu³, and Kathryn H. Schmitz¹

¹Center for Clinical Epidemiology and Biostatistics, University of Pennsylvania, PA

²Division of Female Pelvic Medicine and Reconstructive Surgery, Department of Obstetrics and Gynecology; Department of Surgery (Urology), University of Medicine and Dentistry of New Jersey, New Brunswick, NJ

³Division of Gynecologic Oncology, Department of Obstetrics and Gynecology, University of Pennsylvania, PA

Abstract

Introduction—The extent to which physical activity (PA) participation among uterine cancer survivors may be limited by physical and functional impairments (PFIs) related to cancer treatment is unknown. We sought to describe PA participation, characterize the prevalence of PFIs, and examine the association between PFI status and PA participation within this population.

Methods—We conducted a study using a mailed survey among uterine cancer survivors who received treatment at a university hospital. We asked about PA and PFIs using validated self-report questionnaires. PA was calculated using metabolic equivalent hours per week (MET-hrs·wk⁻¹). PFI was defined as having one or more of the following symptoms: lower limb lymphedema (LLL), general pain, fatigue, or severe bladder, bowel, or pelvic issues. Ordinal logistic regression was used to quantify the odds ratio (OR) between PA and PFIs.

Results—The response rate to our survey was 43%. Among the 213 study participants, 40%, 13%, 13%, 12%, and 23% reported participating in <3.0, 3.0–8.9, 9.0–17.9, 18.0–26.9, and ≥27.0 MET-hrs·wk⁻¹ of PA, respectively. Walking is the preferred mode of exercise for physically active uterine cancer survivors. 53% of survivors experience at least one PFI. The most common PFI is LLL (36.2%), followed by general pain (22.5%). The OR of PFI decreased as MET-hrs·wk⁻¹ of PA increased (OR = 0.51, 95% CI: 0.31–0.84; *P*=.009).

Conclusion—The majority of uterine cancer survivors experience physical and functional impairments that significantly reduce the likelihood of PA participation. Physical activity recommendations for uterine cancer survivors should take into account treatment-related impairments that can affect PA participation.

Corresponding Author: Kathryn H. Schmitz, University of Pennsylvania School of Medicine, 423 Guardian Drive, 8th Floor, Blockley Hall, Philadelphia, PA 19104, Phone: 215–898–6604, Fax: 215–573–5311, schmitz@mail.med.upenn.edu.

Conflicts of Interest

All other authors declare there are no conflicts of interest.

Keywords

gynecologic cancer; exercise; lymphedema; pain; barriers to exercise

INTRODUCTION

Uterine cancer is the fourth most common cancer among women and the most commonly diagnosed gynecologic cancer in the United States, with 47,130 incident cases expected in 2012 (20). Endometrial cancer is the most common type of uterine cancer, accounting for 95% of all uterine cancer diagnoses (20). Obesity is a major risk factor for developing uterine cancer; approximately half of all cases of uterine cancer in postmenopausal women are attributable to excess weight and obesity (38). Uterine cancer is usually diagnosed at early stages due to the onset of noticeable symptoms, such as abnormal or post-menopausal bleeding (22). The primary treatment modality is surgery, which may be followed by adjuvant radiation and/or chemotherapy depending on stage and risk factors for recurrence (22).

As a consequence of early detection and successful medical intervention, the five year survival rate for early stage uterine cancer patients is >95% (20). However, survivors experience symptoms and adverse effects associated with their cancer and cancer treatment, such as pain, fatigue, pelvic floor symptoms, and activities of daily living (ADL) impairments for 5–7 years after completing treatment (3, 11, 12, 26, 28, 29, 44). Approximately 50% of women with uterine cancer are obese (7). This is important because uterine cancer survivors frequently suffer from obesity-related comorbidities, including diabetes, hypertension, heart disease, pulmonary disease, and musculoskeletal impairments (7, 21). Higher body mass index is associated with decreased survival and lower quality of life (QOL) among uterine cancer survivors (13, 18, 33, 37, 38, 43).

While evidence from observational trials suggest increasing physical activity (PA) among uterine cancer survivors is a viable path to improve QOL and reduce morbidity and mortality, the extent to which cancer-related adverse effects influence PA participation in this population remains unknown (7, 41, 42). Several studies have reported that ~70% of endometrial cancer survivors that do not meet public health guidelines of 150 minutes of moderate-to-vigorous intensity physical activity per week (7, 9, 18, 24, 36). According to a 2011 CDC report, 53.7% of 55–64 year olds and 64.6% of people age 65 years and older met neither the aerobic activity nor muscle-strengthening guidelines (34); this suggests that uterine cancer survivors are less likely to participate in PA than their peers in the general population. High intention to participate in PA, high self-efficacy in completing PA, and favorable attitudes and beliefs regarding PA are factors associated with higher rates of participation in PA among uterine cancer survivors. (24). Conversely, poor health, lack of time, poor weather conditions, injury, and fatigue are factors associated with lower rates of participation in PA among uterine cancer survivors (24). Few studies have detailed PA participation patterns among uterine cancer survivors, including type, frequency, intensity, and duration of PA.

The purpose of the present study was three-fold. First, we sought to describe the characteristics of PA participation among a cohort of uterine cancer survivors. Second, we sought to describe the prevalence of at least one cancer-related adverse effect or other health condition that may impair PA participation among survivors. Third, we sought to investigate the extent to which physical and functional impairments (PFI) associated with PA.

METHODS

Participants and procedures

We conducted a cross-sectional survey of patients with uterine cancer who received care at a large university cancer center. Potential participants included women ≥ 20 years old, with a history of uterine cancer. Potentially eligible uterine cancer survivors were identified using fellow surgical case logs from 2008–2010, and ICD-9 diagnosis codes 179.0 and 182.0–182.8, from 2006–2010. Women identified by their physician or with notes in the surgical logs that were unable to complete a written survey due to illiteracy, lack of English fluency, or cognitive impairment, were excluded. Women identified to meet inclusion criteria by study staff were sent a letter signed by their oncologist explaining the purpose of the study. Potentially eligible survivors were provided with the option to decline participation within one week of receiving the letter from their oncologist. Those who did not decline participation were sent a survey to complete. After two weeks, a second survey was sent to those who did not reply to the first mailed survey (25). This protocol was approved by the University Institutional Review Board and the Clinical Trials Scientific Review and Monitoring Committee of the cancer center. Women who mailed back a completed survey were classified as providing their informed consent.

Physical activity assessment

The Paffenbarger Physical Activity Questionnaire (PPAQ) was used to assess participation in PA (35). The PPAQ has been validated (2, 27), and used previously among cancer survivors (30, 31). Participants were asked to “list any sports, leisure, or recreation activities you have participated in on a regular basis during the past year.” Survivors were also asked to list the average number of sessions per week and the duration of each session for each PA listed. Trained research staff then coded each PA listed with a metabolic equivalent (MET) using the compendium of physical activities (2). For reference, one MET is the energy expended when sitting quietly for one hour, and 3.5 METs is walking for pleasure. For each MET value, we calculated the weekly activity-specific MET-hrs·wk⁻¹, as the product of the MET value, the number of sessions per week, and the number of hours per session. For each participant, we summed the activity-specific MET-hrs·wk⁻¹ to generate an aggregate measure of MET-hrs·wk⁻¹. Consistent with prior analyses (30, 31), we created categories of MET-hrs·wk⁻¹, defined as <3.0, 3.0–8.9, 9.0–17.9, 18.0–26.9, and ≥ 27.0 that correspond to <1.0, 1.0–2.9, 3.0–5.9, 6.0–8.9, and ≥ 9.0 hours per week of moderate-intensity PA.

Defining physical and functional impairments

Participants completed questionnaires to examine symptoms and side effects of cancer diagnosis and treatment that we hypothesized to influence PA. Participants with significant pain, severe fatigue, lower limb lymphedema (LLL), severe bladder/urinary symptoms,

severe bowel symptoms and/or severe pelvic symptoms as defined below, were classified as having PFI. The individual components of the PFI are described below in detail.

Lower limb lymphedema assessment

The Gynecologic Cancer Lymphedema Questionnaire (GCLQ) was used to assess symptoms associated with LLL (14). The GCLQ is a validated self-report measure that assesses seven domains of symptoms in the lower extremities. The seven domains include heaviness, general swelling, limb-related swelling, infection, aching, numbness, and physical function, with one or more symptom questions per domain. Women reporting 5 symptoms of the lower extremities within the seven above-listed domains were classified as having LLL (14). Our study group omitted one of the three questions in the swelling-general domain (question #20). To ensure that our findings were robust, we conducted sensitivity analyses assuming everyone responded “no” or “yes” to this question; our findings were consistent in all analyses with those reported herein.

Pain assessment

The Brief Pain Inventory (BPI) was used to assess pain symptoms (17). The BPI is a validated self-report measure that asks patients to rate their pain at the time of responding to the questionnaire (“pain now”) and also at its “worst”, “least”, and “average” over the last week. Item ratings are scored on a 0–10 scale. Ratings can be combined to generate composite indices of pain severity and pain interference. Participants with either a composite pain severity or pain interference index 5 were classified as having significant pain capable of impairing PA participation (17).

Fatigue assessment

Fatigue was assessed using the Fatigue Symptom Inventory (FSI) (23). The FSI is a valid and reliable measure of fatigue among cancer survivors (19). The FSI is a 14-item measure that assesses three domains of fatigue, including frequency, severity, and interference with activities of daily living. A composite fatigue score was derived by calculating the average across the three fatigue domains. Women with FSI composite scores 80 were classified as having severe fatigue capable of impairing PA participation (19).

Urinary, bowel, and pelvic symptom assessment

The Pelvic Floor Distress Inventory (PFDI-20) and Pelvic Floor Incontinence Questionnaire (PFIQ-7) short forms were used to assess the severity of urinary, bowel, and pelvic symptoms (5). The PFDI is a validated instrument that is both a symptom and bother inventory. It assesses three domains of distress: the Pelvic Organ Prolapse Distress Inventory 6 (POPDI-6), the Colorectal Anal Distress Inventory 8 (CRADI-8), and the Urinary Distress Inventory 6 (UDI-6). Each scale is scored from 0 (least distress) to 100 (greatest distress) and has been shown to demonstrate good test-retest reliability (4, 6). Participants who reported POPDI-6, CRADI-8, or UDI-6 score 75 were classified as having severe pelvic issues, severe bowel issues, or severe bladder issues, respectively. These cutoff scores were identified as the meaningful cut points since they correspond with average responses of a “moderate” bother score on the PFDI (i.e., 3 on the 0–4 scale).

The PFIQ-7 is a validated questionnaire that assesses the extent to which bladder, bowel, or vaginal symptoms affect women's activities, relationships, and feelings (6, 45). Participants rated their level of interference according to symptoms on a 4-point scale ranging from "not at all" to "quite a bit." Responses to the PFIQ are scored according to symptom, yielding scores on a scale from 0–100 for the Urinary Impact Questionnaire (UIQ-7), the Colorectal-Anal Impact Questionnaire (CRAIQ-7), and Pelvic Organ Prolapse Impact Questionnaire (POPIQ-7). Participants who reported UIQ-7, CRAIQ-7, or POPIQ-7 score ≥ 66.7 were classified as having severe bladder symptoms, severe bowel symptoms, or severe pelvic symptoms, respectively. These cutoff scores were identified as the meaningful cut points since they correspond with average responses of being "moderately" affected by bladder or urinary symptoms on the PFIQ-7 (i.e., 2 on the 0–3 scale).

Participants who reported severe bladder, bowel, or pelvic issues on either of the relevant PFDI-20 or PFIQ-7 subscales were classified as having PFI.

Covariates

Information on covariates came from self-report or electronic medical records. Variables collected from self-report included age, marital status, race, education, employment, height, weight, and comorbidities whose symptoms may be similar to our defined PFIs were assessed using the Charlson Comorbidity Index (15, 16). Variables collected from the electronic medical record included pathology type, stage of cancer, time since diagnosis, cancer treatment history, and recurrence.

Statistical analysis

Response rates to our survey were calculated using the American Association for Public Opinion Research; this method which counts complete and partial survey responses as respondents and divides by the number of surveys mailed (25). We performed descriptive statistics and bivariate analyses on all study variables using the Wilcoxon Rank-Sum test for continuous variables and chi square test for categorical variables. The distribution of PFI's was significantly skewed ($P < 0.0001$ for skewness). We created a binary variable (0 PFI vs. 1 PFI) which had a similar proportion of participants (47% vs. 53%), respectively. Ordered logistic regression models estimated the odds ratio (OR) with 95% confidence intervals (95% CI) to estimate the proportional odds of increasing PA given the presence of a PFI. Statistical tests were two-sided and $P < .05$ was the threshold for statistical significance. All statistical analyses were conducted using Stata 12.0 (College Station, TX).

RESULTS

Participant Characteristics

We identified 531 potentially eligible participants using the fellow surgical case logs and ICD-9 codes. Among the 531 mailed letters, we had a 43% response rate. Variables analyzed as potential covariates are shown in Tables 1 and 2. The age of the 213 participants ranged from 29–94 years (Table 1). The majority of participants reported being white, married, high school graduates, and retired or working full time. Employment status (retired vs. full-time) was associated with PFI status ($p = .03$). Participants were commonly

diagnosed with stage I endometrioid adenocarcinoma, and treated with surgery (Table 2). The BMI of study participants ranged from 14–67 kg/m²; 26%, 22%, and 52% reported a BMI of <25.0, 25.0–29.9, and ≥30.0 kg/m², respectively. The median Charlson comorbidity index score was 2 (IQR = 2–4).

Physical activity among Uterine Cancer Survivors

Among the 213 participants, 80 (38%) reported no PA participation on a regular basis in the past year. Seventy-one (33%), 33 (15%), 21 (10%), and 8 (4%) participants reported participating in one, two, three, and four or more weekly PA's, respectively (Table 3). Among participants reporting ≥1 PA, the most common modality of PA was walking.

Characteristics between participants with versus without PFI

Among the 213 participants, 113 (53%) experienced at least one PFI hypothesized to associate with PA participation (Table 4). The most common PFI was LLL; 77 (36%) participants were classified as having LLL. Forty-eight (22.5%) participants reported significant pain, and 12 (5.6%) participants reported severe fatigue. The number of PFI's was not related to BMI as a continuous variable ($r = 0.07$; $p = 0.28$) or time since diagnosis ($r = 0.10$; $p = 0.16$). The number of PFI's was related to BMI as a categorical variable (<25.0, 25.0–30.0, ≥30.0) (spearman rho (ρ) = 0.16; $p = 0.02$), but not time since diagnosis ($\rho = 0.08$; $p = 0.24$). Fatigue and pain were not related to BMI as a continuous variable ($r = 0.11$; $p = 0.09$), and ($r = 0.11$; $p = 0.11$), respectively. Pain was related to BMI as a categorical variable ($\rho = 0.21$; $p = 0.002$), but not fatigue ($\rho = 0.08$; $p = 0.22$). The number of LLL symptoms was not related to BMI as a continuous or categorical variable. The least common PFI's were severe bladder/urinary symptoms (3.8%), severe pelvic symptoms (1.4%), and severe bowel symptoms (0.9%). Among the 113 participants who reported a PFI, 89 (78.8%) experienced only one of the above cancer-related adverse events (Table 4). Meanwhile, 13.3% of these participants reported 2 PFIs and 8.0% reported ≥3 PFIs.

Association between PFI and level of physical activity

Among the 213 study participants, 40%, 13%, 13%, 12%, and 23% reported participating in <3.0, 3.0–8.9, 9.0–17.9, 18.0–26.9, and ≥27.0 MET-hrs·wk⁻¹ of PA, respectively (Table 5). The OR of PFI decreased as MET-hrs·wk⁻¹ of PA increased (OR = 0.51 (95% CI: 0.31–0.84); $P = .009$). This corresponds to a 49% reduced likelihood for uterine cancer survivors with ≥1 PFI to increase their MET-hrs·wk⁻¹ to the next higher level of PA (i.e., reporting 3.0–8.9 MET-hrs·wk⁻¹ instead of <3.0 MET-hrs·wk⁻¹) compared to uterine cancer survivors without a PFI.

DISCUSSION

The major findings of this study are that 53% of uterine cancer survivors experience at least one PFI, and that this is associated with low levels of physical activity. Among uterine cancer survivors the most common PFIs are LLL and pain. Notably, many uterine cancer survivors do not meet recommended guidelines for PA, and 38% self-report being completely sedentary; this is comparable to the general population, in which 28–34% of adults aged 65 to 74 years and 35–44% of adults ages 75 or older are inactive (8). Among

uterine cancer survivors who are physically active, low- and moderate-intensity PAs, such as walking, are preferred. Women who reported a PFI were more frequently retired and generally reported a smaller weekly volume of PA participation. These findings provide evidence that PA participation may be influenced by PFIs in uterine cancer survivors. Our study confirms previous reports that comorbid health problems such as obesity and musculoskeletal impairments may negatively impact participation in PA among uterine cancer survivors (7). Attitudes about PA, such as intention, self-efficacy, and beliefs are also important factors that may be mediated by comorbid health conditions such as obesity, which act to negatively influence PA participation (24). Poor health is the most common barrier to participating in PA among endometrial cancer survivors (24).

The major limitation of this study is its cross-sectional design, which precludes determining causal associations. For example, survivors who are more physically active may have more leakage of urine stemming from their exercise; conversely, those with urinary incontinence may have an aversion to PA. Though we postulate uterine cancer survivors who self-report PFIs may be physically or psychologically unable to engage in PA, it is plausible that individuals who engage in more PA subsequently experience fewer PFIs. Exercise has been shown to have beneficial effects on cancer survivor's health-related QOL, including reduced fatigue and pain (32). Other limitations to our study include possible selection bias; we do not have data to determine if women who experienced PFIs were more likely to reply to the survey. Alternatively, women who were doing poorly or doing extremely well may have not replied to the survey. Information bias is another possible limitation, though we attempted to minimize this bias by using validated questionnaires and by verifying self-reported exposure and outcome data with survivors' medical records. A limitation to self-reported PA questionnaire was that participants provided non-specific descriptions of PA such as "gym-based activities". Although self-reported PA is correlated with objective measures of PA (39), 47% of participants in our study reported meeting physical activity guidelines (i.e., 9.0 MET-hrs-wk⁻¹), whereas only approximately 10% of US adults meet such guidelines (40). Therefore, participants in our study may have over reported PA levels. We also cannot rule out the possibility that our conservative definition of PFIs might exclude other adverse effects capable of influencing PA participation such as arthritis or psychosocial issues (1, 7). Furthermore, some of the PFI's in our study may have not been specific to cancer, rather they be related to obesity or other comorbid health conditions (7, 24). Our study sample was hospital-based, and was not population-based which is a limitation compared to other previously published studies among uterine cancer survivors (7, 9, 24). Thus, our estimates may underestimate the extent to which PFIs influence PA participation in this population.

Consistent with previous studies, our results confirm the prevalence of cancer-related adverse effects as well as obesity-related comorbidities among uterine cancer survivors. More importantly, these findings identify potential barriers to PA participation in this population and highlight the need for PA interventions to consider PFI status as a tailoring variable. Uterine cancer survivors who are sedentary and/or obese are unlikely to spontaneously modify their exercise and diet following cancer diagnosis and treatment (22). There is limited, but growing evidence to suggest that exercise interventions can improve health and QOL outcomes for uterine cancer survivors. A recent randomized controlled study of an interventional lifestyle program demonstrated that uterine cancer survivors can

lose weight and improve their exercise for 6 months following the intervention (41). At 12 months, the intervention group lost 3.5 kg while the control group gained 1.4 kg, suggesting that a lifestyle intervention program among obese uterine cancer survivors is feasible and can result in sustained behavior change and weight loss over a one year period (41). No serious adverse events were reported in this study (41).

To build upon such results, physician and nurse recommendations for PA among this population should account for the impairments as described in this study. Future research should focus on elucidating specific PA recommendations based on PFI status so as to reduce obesity-related comorbidities and improve QOL among uterine cancer survivors. Considering the prevalence of sedentary behavior in this population, qualitative interviews with uterine cancer survivors who report little to no PA participation would be a useful first step towards understanding potential motivating factors and barriers to PA in this population. In addition to PFI's, understanding behavioral determinants such as intention, self-efficacy, and attitudes associated with PA are also of importance to promote PA among uterine cancer survivors (24). Interventions designed to help uterine cancer survivors overcome the many barriers to PA, including but not limited to the PFI's described here, may provide significant benefits in terms of QOL improvements and reductions in mortality attributable to obesity-related comorbidities.

Conclusion

Among uterine cancer survivors, 62% regularly participate in at least one weekly PA. Walking is the preferred mode of exercise for physically active uterine cancer survivors. Meanwhile, 53% of uterine cancer survivors experience at least one PFI. The most common PFI is LLL (36.2%), followed by pain (22.5%). Most importantly, these PFIs are negatively associated with participation in PA. Physicians recommending PA for uterine cancer survivors should take into account treatment-related impairments that can affect PA participation, and health and fitness professionals should receive adequate training to properly prescribe and modify PA interventions in order to improve health and QOL, without sacrificing safety.

Acknowledgments

Source of Funding: The senior author, KH Schmitz, has received funding from the National Institutes of Health unrelated to this work. The results of this study do not constitute endorsement by ACSM. This work was completed without funding.

References

1. Ahlberg K, Ekman T, Wallgren A, Gaston-Johansson F. Fatigue, psychological distress, coping and quality of life in patients with uterine cancer. *J Adv Nurs*. 2004; 45(2):205–13. [PubMed: 14706006]
2. Ainsworth BE, Leon AS, Richardson MT, Jacobs DR, Paffenbarger RS Jr. Accuracy of the College Alumnae Physical Activity Questionnaire. *J Clin Epidemiol*. 1993; 46(12):1403–11. [PubMed: 8263567]
3. Andersen BL. Predicting sexual and psychologic morbidity and improving the quality of life for women with gynecologic cancer. *Cancer*. 1993; 71(4 Suppl):1678–90. [PubMed: 8431906]

4. Barber MD, Kuchibhatla MN, Pieper CF, Bump RC. Psychometric evaluation of 2 comprehensive condition-specific quality of life instruments for women with pelvic floor disorders. *Am J Obstet Gynecol.* 2001; 185(6):1388–95. [PubMed: 11744914]
5. Barber MD, Walters MD, Bump RC. Short forms of two condition-specific quality-of-life questionnaires for women with pelvic floor disorders (PFDI-20 and PFIQ-7). *Am J Obstet Gynecol.* 2005; 193(1):103–13. [PubMed: 16021067]
6. Barber MD, Walters MD, Cundiff GW. PESSRI Trial Group. Responsiveness of the Pelvic Floor Distress Inventory (PFDI) and Pelvic Floor Impact Questionnaire (PFIQ) in women undergoing vaginal surgery and pessary treatment for pelvic organ prolapse. *Am J Obstet Gynecol.* 2006; 194(5):1492–8. [PubMed: 16647933]
7. Basen-Engquist K, Scruggs S, Jhingran A, Bodurka DC, Lu K, Ramondetta L, Hughes D, Carmack Taylor C. Physical activity and obesity in endometrial cancer survivors: associations with pain, fatigue, and physical functioning. *Am J Obstet Gynecol.* 2009; 200(3):288.e1–288.e8. [PubMed: 19110220]
8. Behavioral Risk Factor Surveillance System Survey Data. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2010. [Internet][cited 2013 July 28]. Available from: <http://www.cdc.gov/brfss/>
9. Blanchard CM, Courneya KS, Stein K. American Cancer Society's SCS-II. Cancer survivors' adherence to lifestyle behavior recommendations and associations with health-related quality of life: results from the American Cancer Society's SCS-II. *J Clin Oncol.* 2008; 26(13):2198–204. [PubMed: 18445845]
10. Branca F, Ferrari M. Impact of micronutrient deficiencies on growth: the stunting syndrome. *Ann Nutr Metab.* 2002; 46 (Suppl 1):8–17. [PubMed: 12428076]
11. Butler L, Banfield V, Sveinson T, Allen K. Conceptualizing sexual health in cancer care. *West J Nurs Res.* 1998; 20(6):683, 99. discussion 700–5. [PubMed: 9842287]
12. Bye A, Trope C, Loge JH, Hjermstad M, Kaasa S. Health-related quality of life and occurrence of intestinal side effects after pelvic radiotherapy--evaluation of long-term effects of diagnosis and treatment. *Acta Oncol.* 2000; 39(2):173–80. [PubMed: 10859007]
13. Calle EE, Rodriguez C, Walker-Thurmond K, Thun MJ. Overweight, obesity, and mortality from cancer in a prospectively studied cohort of U.S. adults. *N Engl J Med.* 2003; 348(17):1625–38. [PubMed: 12711737]
14. Carter J, Raviv L, Appollo K, Baser RE, Iasonos A, Barakat RR. A pilot study using the Gynecologic Cancer Lymphedema Questionnaire (GCLQ) as a clinical care tool to identify lower extremity lymphedema in gynecologic cancer survivors. *Gynecol Oncol.* 2010; 117(2):317–23. [PubMed: 20163847]
15. Charlson M, Szatrowski TP, Peterson J, Gold J. Validation of a combined comorbidity index. *J Clin Epidemiol.* 1994; 47(11):1245–51. [PubMed: 7722560]
16. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis.* 1987; 40(5):373–83. [PubMed: 3558716]
17. Cleeland CS, Ryan KM. Pain assessment: global use of the Brief Pain Inventory. *Ann Acad Med Singapore.* 1994; 23(2):129–38. [PubMed: 8080219]
18. Courneya KS, Karvinen KH, Campbell KL, Pearcey RG, Dundas G, Capstick V, Tonkin KS. Associations among exercise, body weight, and quality of life in a population-based sample of endometrial cancer survivors. *Gynecol Oncol.* 2005; 97(2):422–30. [PubMed: 15863140]
19. Donovan KA, Jacobsen PB. The Fatigue Symptom Inventory: a systematic review of its psychometric properties. *Support Care Cancer.* 2010; 19(2):169–85. [PubMed: 20824482]
20. Endometrial cancer [Internet]. American Cancer Society; [cited 2012 August 10]. Available from: <http://www.cancer.org>
21. Fader AN, Arriba LN, Frasure HE, von Gruenigen VE. Endometrial cancer and obesity: epidemiology, biomarkers, prevention and survivorship. *Gynecol Oncol.* 2009; 114(1):121–7. [PubMed: 19406460]
22. Gil KM, von Gruenigen VE. Physical activity and gynecologic cancer survivorship. *Recent Results Cancer Res.* 2011; 186:305–15. [PubMed: 21113770]

23. Hann DM, Jacobsen PB, Azzarello LM, Martin SC, Curran SL, Fields KK, Greenberg H, Lyman G. Measurement of fatigue in cancer patients: development and validation of the Fatigue Symptom Inventory. *Qual Life Res.* 1998; 7(4):301–10. [PubMed: 9610214]
24. Karvinen KH, Courneya KS, Campbell KL, Pearcey RG, Dundas G, Capstick V, Tonkin KS. Correlates of exercise motivation and behavior in a population-based sample of endometrial cancer survivors: an application of the Theory of Planned Behavior. *Int J Behav Nutr Phys Act.* 2007; 4:21. [PubMed: 17537255]
25. Kelly BJ, Frazee TK, Hornik RC. Response rates to a mailed survey of a representative sample of cancer patients randomly drawn from the Pennsylvania Cancer Registry: a randomized trial of incentive and length effects. *BMC Med Res Methodol.* 2010; 10:65, 2288–10–65. [PubMed: 20630101]
26. Klee M, Machin D. Health-related quality of life of patients with endometrial cancer who are disease-free following external irradiation. *Acta Oncol.* 2001; 40(7):816–24. [PubMed: 11859980]
27. Kohl HW, Blair SN, Paffenbarger RS Jr, Macera CA, Kronenfeld JJ. A mail survey of physical activity habits as related to measured physical fitness. *Am J Epidemiol.* 1988; 127(6):1228–39. [PubMed: 3369421]
28. Kullmer U, Stenger K, Milch W, Zygmunt M, Sachsse S, Munstedt K. Self-concept, body image, and use of unconventional therapies in patients with gynaecological malignancies in the state of complete remission and recurrence. *Eur J Obstet Gynecol Reprod Biol.* 1999; 82(1):101–6. [PubMed: 10192496]
29. Li C, Samsioe G, Iosif C. Quality of life in endometrial cancer survivors. *Maturitas.* 1999; 31(3): 227–36. [PubMed: 10340282]
30. Meyerhardt JA, Giovannucci EL, Holmes MD, Chan AT, Chan JA, Colditz GA, Fuchs CS. Physical activity and survival after colorectal cancer diagnosis. *J Clin Oncol.* 2006; 24(22):3527–34. [PubMed: 16822844]
31. Meyerhardt JA, Heseltine D, Niedzwiecki D, Hollis D, Saltz LB, Mayer RJ, Thomas J, Nelson H, Whittom R, Hantel A, Schilsky RL, Fuchs CS. Impact of physical activity on cancer recurrence and survival in patients with stage III colon cancer: findings from CALGB 89803. *J Clin Oncol.* 2006; 24(22):3535–41. [PubMed: 16822843]
32. Mishra SI, Scherer RW, Geigle PM, Berlanstein DR, Topaloglu O, Gotay CC, Snyder C. Exercise interventions on health-related quality of life for cancer survivors. *Cochrane Database Syst Rev.* 2012; 8:CD007566. [PubMed: 22895961]
33. Modesitt SC, Tian C, Kryscio R, Thigpen JT, Randall ME, Gallion HH, Fleming GF. Gynecologic Oncology Group. Impact of body mass index on treatment outcomes in endometrial cancer patients receiving doxorubicin and cisplatin: a Gynecologic Oncology Group study. *Gynecol Oncol.* 2007; 105(1):59–65. [PubMed: 17150247]
34. National Center for Health Statistics. *Health, United States, 2011: With Special Feature on Socioeconomic Status and Health.* Hyattsville, MD: National Center for Health Statistics; 2012. p. 271 Available From
35. Paffenbarger RS Jr, Hyde RT, Wing AL. Physical activity and incidence of cancer in diverse populations: a preliminary report. *Am J Clin Nutr.* 1987; 45(1 Suppl):312–7. [PubMed: 3799521]
36. Pate RR, Pratt M, Blair SN, Haskell WL, Macera CA, Bouchard C, Buchner D, Ettinger W, Heath GW, King AC. Physical activity and public health. A recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *JAMA.* 1995; 273(5):402–7. [PubMed: 7823386]
37. Patel AV, Feigelson HS, Talbot JT, McCullough ML, Rodriguez C, Patel RC, Thun MJ, Calle EE. The role of body weight in the relationship between physical activity and endometrial cancer: results from a large cohort of US women. *Int J Cancer.* 2008; 123(8):1877–82. [PubMed: 18651569]
38. Reeves GK, Pirie K, Beral V, Green J, Spencer E, Bull D. Million Women Study Collaboration. Cancer incidence and mortality in relation to body mass index in the Million Women Study: cohort study. *BMJ.* 2007; 335(7630):1134. [PubMed: 17986716]

39. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc.* 2008; 40(1):181–8. [PubMed: 18091006]
40. Tucker JM, Welk GJ, Beyler NK. Physical activity in U.S.: adults compliance with the Physical Activity Guidelines for Americans. *Am J Prev Med.* 2011; 40(4):454–61. [PubMed: 21406280]
41. von Gruenigen VE, Courneya KS, Gibbons HE, Kavanagh MB, Waggoner SE, Lerner E. Feasibility and effectiveness of a lifestyle intervention program in obese endometrial cancer patients: a randomized trial. *Gynecol Oncol.* 2008; 109(1):19–26. [PubMed: 18243282]
42. von Gruenigen VE, Gil KM, Frasure HE, Jenison EL, Hopkins MP. The impact of obesity and age on quality of life in gynecologic surgery. *Am J Obstet Gynecol.* 2005; 193(4):1369–75. [PubMed: 16202728]
43. von Gruenigen VE, Tian C, Frasure H, Waggoner S, Keys H, Barakat RR. Treatment effects, disease recurrence, and survival in obese women with early endometrial carcinoma : a Gynecologic Oncology Group study. *Cancer.* 2006; 107(12):2786–91. [PubMed: 17096437]
44. Weiss E, Hirnle P, Arnold-Bofinger H, Hess CF, Bamberg M. Therapeutic outcome and relation of acute and late side effects in the adjuvant radiotherapy of endometrial carcinoma stage I and II. *Radiother Oncol.* 1999; 53(1):37–44. [PubMed: 10624851]
45. Wren PA, Janz NK, Brubaker L, Fitzgerald MP, Weber AM, LaPorte FB, Wei JT. Pelvic Floor Disorders Network. Reliability of health-related quality-of-life measures 1 year after surgical procedures for pelvic floor disorders. *Am J Obstet Gynecol.* 2005; 192(3):780–8. [PubMed: 15746672]

Table 1

Demographic characteristics stratified by physical and functional impairment (PFI) status

Variable	Total Sample (n=213)	No PFI (n=100)	1 PFI (n=113)	<i>P</i> ^a
Age — yr	63.5±10.6	62.8±10.6	64.2±10.6	.58
Marital status — no. (%)				
Married	128 (60%)	61 (61%)	67 (59%)	Referent
Widowed	33 (16%)	11 (11%)	22 (19%)	.14
Divorced or separated	31 (15%)	18 (18%)	13 (12%)	.30
Never married	20 (9%)	9 (9%)	11 (10%)	.83
Self-reported race — no. (%)				
White	177 (84%)	85 (86%)	92 (81%)	Referent
Black	28 (13%)	10 (10%)	18 (16%)	.23
Other	7 (3%)	4 (4%)	3 (3%)	.64
Education — no. (%)				
College degree or more	114 (54%)	56 (56%)	58 (52%)	Referent
Some college	51 (24%)	24 (24%)	27 (24%)	.81
High school or less	46 (22%)	19 (19%)	27 (24%)	.37
Employment — no. (%)				
Retired	94 (45%)	40 (40%)	54 (49%)	Referent
Full time	80 (38%)	47 (47%)	33 (30%)	.03
Homemaker	16 (8%)	6 (6%)	10 (9%)	.71
Other	14 (7%)	5 (5%)	9 (8%)	.63
Unemployed	7 (3%)	2 (2%)	5 (5%)	.47

^aBy Wilcoxon rank sum, or chi square test. Values may not sum to 213 or 100% due to rounding error and item non-response.

Table 2

Clinical characteristics stratified by physical and functional impairment (PFI) status

Variable	Total Sample (n=213)	No PFI (n=100)	1 PFI (n=113)	<i>P</i> ^a
Pathology type — no. (%)				
Endometrioid Adenocarcinoma	158 (75%)	75 (76%)	83 (73%)	Referent
Papillary serous or Clear Cell	35 (17%)	15 (15%)	20 (18%)	.62
Sarcoma	8 (4%)	5 (5%)	3 (3%)	.41
Carcinosarcoma	8 (4%)	4 (4%)	4 (4%)	.89
Other (Undifferentiated)	3 (1%)	0 (0%)	3 (3%)	.10
Stage — no. (%)				
1	157 (74%)	75 (75%)	82 (73%)	Referent
2	13 (6%)	6 (6%)	7 (6%)	.91
3	26 (12%)	11 (11%)	15 (13%)	.61
4	5 (2%)	2 (2%)	3 (3%)	.73
Unknown	12 (6%)	5 (5%)	6 (5%)	.88
Treatment Modalities — no. (%)				
Surgery Only	100 (47%)	46 (46%)	54 (48%)	Referent
Surgery, Radiation	37 (17%)	17 (17%)	20 (18%)	.99
Surgery, Radiation, and Chemotherapy	47 (22%)	24 (24%)	23 (20%)	.57
Chemotherapy only	22 (10%)	9 (9%)	13 (12%)	.66
None or Unknown	7 (3%)	4 (4%)	3 (3%)	.57
No. of nodes removed	9.0±10.2	9.3±9.9	8.6±10.5	.31
Time since diagnosis — no. (%)				
0–2 yrs	69 (32%)	32 (32%)	36 (32%)	Referent
3–4 yrs	94 (44%)	48 (48%)	46 (41%)	.62
5–6 yrs	50 (23%)	19 (19%)	30 (27%)	.37
BMI — kg/m ²	31.4±9.8	31.5±10.7	31.2±9.0	.58
Median Charlson Comorbidity score ^b	2 (2–4)	2 (2–3.5)	2 (2–5)	.70

^aBy Wilcoxon rank sum, or chi square test. Values may not sum to 213 or 100% due to rounding error and item non-response.

^bIncludes the following comorbidities: myocardial infarction, congestive heart failure, peripheral vascular disease, cerebrovascular disease, dementia, COPD, connective tissue disease, ulcer disease, diabetes mellitus, moderate to severe chronic kidney disease, hemiplegia, leukemia, malignant lymphoma, solid tumor, liver disease, and AIDS. There were no significant differences in any specific comorbidity ($p > .10$).

Table 3

Physical activity characteristics of uterine cancer survivors

Characteristic	Activity Specific				Total MET-hrs/week (mean ± SD)
	% doing activity	Bouts per week (mean ± SD)	Minutes per bout (mean ± SD)	Total MET-hrs/week (mean ± SD)	
Physical activities reported with 5% prevalence in study sample					
Walking - 3.5mph pace (3.8 METs)	89 (42%)	4.1 ± 2.6	55.0 ± 49.4	30.3 ± 35.4	
Gym based activities* (5.5 METs)	23 (11%)	2.8 ± 1.6	67.9 ± 34.1	43.7 ± 33.0	
Swimming (7.0 METs)	18 (8%)	3.5 ± 2.1	62.9 ± 33.8	43.8 ± 29.5	
Hatha Yoga / Mild stretching (2.5 METs)	17 (8%)	2.4 ± 1.7	65.6 ± 31.1	32.9 ± 46.5	
Bicycling (4.0 METs)	14 (7%)	3.1 ± 1.9	48.2 ± 31.6	43.7 ± 48.8	
Number of physical activities completed per week including activities from list described above					
0	80 (38%)	—	—	—	—
1	71 (33%)	3.5 ± 2.3	59.6 ± 42.6	15.2 ± 12.3	
2	33 (15%)	6.3 ± 3.4	77.3 ± 56.9	36.8 ± 39.8	
3	21 (10%)	8.9 ± 4.2	78.8 ± 60.0	43.5 ± 22.2	
4	8 (4%)	14.8 ± 7.9	73.0 ± 38.5	76.7 ± 56.3	

* Includes aerobic, resistance, and flexibility exercises.

Table 4

Physical and functional impairments (PFI) among uterine cancer survivors

PFI Characteristic	n	% of 213
Lower Limb Lymphedema ^d	77	36.2
Pain ^b	48	22.5
Severe Fatigue ^c	12	5.6
Pelvic floor symptoms		
Severe Bladder/Urinary symptoms ^d	8	3.8
Severe Bowel symptoms ^e	2	0.9
Severe Pelvic symptoms ^f	3	1.4
Number of above PFIs per Participant		
0	100	47.0
1	89	41.8
2	15	7.0
3	6	2.8
4	2	0.9
5	1	0.5
TOTAL	213	100.0

^aLymphedema: 5 self-reported symptoms on GCLQ.^bPain: BPI pain severity or pain interference scale score 5 (scale: 0–10).^cFatigue: FSI composite fatigue score 80 (scale: 0–134).^dBladder/urinary symptoms: Average impact on UDI-6 (PFDI) 75 (scale: 0–100) or average impact on UIQ-7 (PFIQ) 66.7 (scale: 0–100).^eBowel symptoms: Average impact on CRADI-8 (PFDI) 75 (scale: 0–100) or average impact on CRAIQ-7 (PFIQ) 66.7 (scale: 0–100).^fPelvic symptoms: Average impact on POPDI-6 (PFDI) 75 (scale: 0–100) or average impact on POPIQ-7 (PFIQ) 66.7 (scale: 0–100).^gOne or more of above complications, includes: pain, fatigue, lower limb lymphedema, urinary/bladder symptoms, bowel symptoms, pelvic symptoms.

Table 5

Effect of PFI on Levels of Physical Activity

Physical Activity Intensity (MET·hrs·wk ⁻¹)	Total Sample (n=213)	No PFI (n=100)	PFI (n=113)
<3.0	85 (40%)	33 (33%)	52 (46%)
3.0–8.9	27 (13%)	13 (13%)	14 (12%)
9.0–17.9	26 (13%)	10 (10%)	17 (15%)
18.0–26.9	26 (12%)	16 (16%)	10 (9%)
27.0	48 (23%)	28 (28%)	20 (18%)
Odds Ratio (95% CI)^a	—	—	0.51 (0.31–0.84)
P	—	—	.009

^aOrdered logistic regression controlling for employment status