

A new stenochirid lobster
(Crustacea, Decapoda, Stenochiridae)
from the Early Jurassic of France

Sylvain CHARBONNIER & Denis AUDO

DIRECTEUR DE LA PUBLICATION : Bruno David,
Président du Muséum national d'Histoire naturelle

RÉDACTEUR EN CHEF / *EDITOR-IN-CHIEF*: Didier Merle

ASSISTANTS DE RÉDACTION / *ASSISTANT EDITORS*: Emmanuel Côté (geodiv@mnhn.fr)

MISE EN PAGE / *PAGE LAYOUT*: Emmanuel Côté

COMITÉ SCIENTIFIQUE / *SCIENTIFIC BOARD*:

Christine Argot (Muséum national d'Histoire naturelle, Paris)
Beatrix Azanza (Museo Nacional de Ciencias Naturales, Madrid)
Raymond L. Bernor (Howard University, Washington DC)
Alain Blicek (chercheur CNRS retraité, Haubourdin)
Henning Blom (Uppsala University)
Jean Broutin (Sorbonne Université, Paris)
Gaël Clément (Muséum national d'Histoire naturelle, Paris)
Ted Daeschler (Academy of Natural Sciences, Philadelphie)
Bruno David (Muséum national d'Histoire naturelle, Paris)
Gregory D. Edgecombe (The Natural History Museum, Londres)
Ursula Göhlich (Natural History Museum Vienna)
Jin Meng (American Museum of Natural History, New York)
Brigitte Meyer-Berthaud (CIRAD, Montpellier)
Zhu Min (Chinese Academy of Sciences, Pékin)
Isabelle Rouget (Muséum national d'Histoire naturelle, Paris)
Sevket Sen (Muséum national d'Histoire naturelle, Paris)
Stanislav Štamberg (Museum of Eastern Bohemia, Hradec Králové)
Paul Taylor (The Natural History Museum, Londres)

COUVERTURE / *COVER*:

Made from the Figures of the article.

Geodiversitas est indexé dans / *Geodiversitas is indexed in*:

- Science Citation Index Expanded (SciSearch®)
- ISI Alerting Services®
- Current Contents® / Physical, Chemical, and Earth Sciences®
- Scopus®

Geodiversitas est distribué en version électronique par / *Geodiversitas is distributed electronically by*:

- BioOne® (<http://www.bioone.org>)

Les articles ainsi que les nouveautés nomenclaturales publiés dans *Geodiversitas* sont référencés par /
Articles and nomenclatural novelties published in Geodiversitas are referenced by:

- ZooBank® (<http://zoobank.org>)

Geodiversitas est une revue en flux continu publiée par les Publications scientifiques du Muséum, Paris
Geodiversitas is a fast track journal published by the Museum Science Press, Paris

Les Publications scientifiques du Muséum publient aussi / *The Museum Science Press also publish*:

Adansonia, *Zoosystema*, *Anthropozoologica*, *European Journal of Taxonomy*, *Naturae*, *Cryptogamie* sous-sections *Algologie*, *Bryologie*, *Mycologie*.

Diffusion – Publications scientifiques Muséum national d'Histoire naturelle

CP 41 – 57 rue Cuvier F-75231 Paris cedex 05 (France)

Tél.: 33 (0)1 40 79 48 05 / Fax: 33 (0)1 40 79 38 40

diff.pub@mnhn.fr / <http://sciencepress.mnhn.fr>

© Publications scientifiques du Muséum national d'Histoire naturelle, Paris, 2020
ISSN (imprimé / *print*): 1280-9659/ ISSN (électronique / *electronic*): 1638-9395

A new stenochirid lobster (Crustacea, Decapoda, Stenochiridae) from the Early Jurassic of France

Sylvain CHARBONNIER

Centre de Recherche en Paléontologie – Paris (CR2P, UMR 7207),
(CNRS, MNHN, Sorbonne Université), Muséum national d'Histoire naturelle,
case postale 38, 57 rue Cuvier, F-75231 Paris cedex 05 (France)
sylvain.charbonnier@mnhn.fr

Denis AUDO

Yunnan Key Laboratory for Palaeobiology, Yunnan University,
North Cuihu road 2#, 650091 Kunming (China)
and MEC International Joint Laboratory for Palaeobiology and Palaeoenvironment,
Yunnan University, Kunming (China)

Submitted on 16 July 2019 | accepted on 23 July 2019 | published on 26 March 2020

[urn:lsid:zoobank.org:pub:02F13624-0172-44B2-AEE4-46663760B7A4](https://doi.org/10.5252/geodiversitas2020v42a7)

KEY WORDS

France,
Jurassic,
Sinemurian,
Crustacea,
Decapoda,
Astacidea,
Stenochiridae,
new species.

MOTS CLÉS

France,
Jurassique,
Sinémurien,
Crustacea,
Decapoda,
Astacidea,
Stenochiridae,
espèce nouvelle.

Charbonnier S. & Audo D. 2020. — A new stenochirid lobster (Crustacea, Decapoda, Stenochiridae) from the Early Jurassic of France. *Geodiversitas* 42 (7): 93-102. <https://doi.org/10.5252/geodiversitas2020v42a7>. <http://geodiversitas.com/42/7>

ABSTRACT

The marine clawed lobster genus *Pseudastacus* Oppel, 1861, previously known mainly from the Late Jurassic lithographic limestones of Bavaria (Germany), is herein recorded from the Early Jurassic (Sinemurian, *c.* 190-199 Ma) of France. A new species, *Pseudastacus lemovices* n. sp., is introduced. It is the oldest representative of the genus and of the family Stenochiridae Beurlen, 1928.

RÉSUMÉ

Une nouvelle langoustine sténochiride (Crustacea, Decapoda, Stenochiridae) du Jurassique inférieur de France. *Pseudastacus* Oppel, 1861, un genre de langoustine connu jusqu'à présent principalement dans les calcaires lithographiques du Jurassique supérieur de Bavière (Allemagne) est ici découvert dans le Jurassique inférieur (Sinémurien, *c.* 190-199 Ma) de France. Une nouvelle espèce, *Pseudastacus lemovices* n. sp., est décrite. Il s'agit du plus ancien représentant du genre et de la famille des Stenochiridae Beurlen, 1928.




FIG. 1. — *Pseudastacus lemovices* n. sp. from the Sinemurian of Chauffour-sur-Vell, France: **A**, map of France with location of the type locality; **B-C**, slab CN-NMP-130 with the association of five preserved specimens, natural light (**B**) and UV light (**C**). Abbreviations: **1**, holotype CN-NMP-130-1; **2-5**, paratypes CN-NMP-130-2 to 130-5. Photographs: P. Loubry. Scale bars: 10 mm.

INTRODUCTION

Stenochirid lobsters (Astacidea, Stenochiroidea, Stenochiridae) are a group of exclusively fossil marine clawed lobsters which members are relatively rare and limited to some Jurassic and Cretaceous outcrops (Table 1). Following the phylogeny and

systematics of lobsters proposed by Karasawa *et al.* (2013), the historical composition of the family evolved including five genera (*Stenochirus* Opper, 1861a, *Pseudastacus* Opper, 1861b, *Palaeophoberus* Glaessner, 1932, *Tillocheles* Woods, 1957, *Chilenophoberus* Chong & Förster, 1976). The inclusion of *Pseudastacus* within the stenochirids is recent and any discovery

TABLE 1. — List of the members of the family Stenochiridae Beurlen, 1928.

Taxa	Outcrops		Ages		References
<i>Stenochirus</i> Oppel, 1861					
<i>Stenochirus angutus</i> (Münster, 1839)	Solnhofen	Germany	Tithonian	Late Jurassic	Münster (1839); Oppel (1861a, 1862); Garassino & Schweigert (2006)
<i>Stenochirus mayeri</i> Oppel, 1862	Solnhofen	Germany	Tithonian	Late Jurassic	Oppel (1862); Garassino & Schweigert (2006)
<i>Stenochirus vahldieki</i> Schweigert, Garassino & Riou, 2006	La Voulte-sur-France Rhône		Callovian	Middle Jurassic	Schweigert <i>et al.</i> (2006); Charbonnier (2009)
<i>Pseudastacus</i> Oppel, 1861					
<i>Pseudastacus pustulosus</i> (Münster, 1839)	Solnhofen	Germany	Tithonian	Late Jurassic	Münster (1839); Oppel (1861b, 1862); Garassino & Schweigert (2006)
? <i>Pseudastacus pusillus</i> Van Straelen, 1925	May-sur-Orne	France	Bajocian	Middle Jurassic	Van Straelen (1925)
<i>Pseudastacus lemovices</i> n. sp.	Chauffour-sur-Vell	France	Sinemurian	Early Jurassic	this work
<i>Palaeophoberus</i> Glaessner, 1932					
<i>Palaeophoberus portlandicus</i> Roger & Lapparent, 1944	Hannaches	France	Tithonian	Late Jurassic	Roger & Lapparent (1944)
<i>Palaeophoberus suevicus</i> (Quenstedt, 1867)	Reutlingen	Germany	Aalenian	Middle Jurassic	Quenstedt (1867); Glaessner (1932)
<i>Chilenophoberus</i> Chong & Förster, 1976					
<i>Chilenophoberus atacamensis</i> Chong & Förster, 1976	Cordillera de Domeyko	Chile	Oxfordian	Late Jurassic	Chong & Förster (1976)
<i>Tillocheles</i> Woods, 1957					
<i>Tillocheles kaoriae</i> Yokoi & Karasawa, 2000	Hokkaido	Japan	Cenomanian, Turonian	Late Cretaceous	Yokoi & Karasawa (2000)
<i>Tillocheles shannonae</i> Woods, 1957	Queensland	Australia	Albian	Early Cretaceous	Woods (1957)

is important to understand the history of this group. After a review of the species included in *Pseudastacus*, we describe a new species from the Early Jurassic of France and discuss the systematics and age of stenochirid lobsters.

MATERIAL AND METHODS

The studied material consists of a single slab with five specimens of lobsters (Fig. 1). All are compressed and laterally and/or dorsoventrally flattened on the surface of the slab. The slab was collected during the digging of a pit in the garden of Mrs Boutang and Mr Gomme, at Chauffour-sur-Vell, Corrèze department, France (Fig. 1A). The sediment corresponds to a marine sublithographic limestone of Sinemurian age, possibly early Sinemurian (see Lefavrais-Raymond *et al.* 1976). After these authors, the limit between the early and late Sinemurian is only cartographic, and the late Sinemurian of the area is well dated by the occurrence of a regional bed yielding the dasycladalean green alga *Palaeodasycladus mediterraneus* (Pia, 1920).

The sample is exposed at the “L’Espace de découverte de la Faille de Meyssac et de la Pierre” (acronym: CN-NMP), the local museum of Noailhac (Corrèze department) located 10 km away from the type locality.

The five specimens were studied using a binocular microscope and a camera lucida. The line drawings were made by the first author. Specimens displaying low contrast were imaged by the second author under cross-polarized light (i.e. light source

equipped with a linear polarizing filter and camera lens also equipped with a polarizing filter whose optical axis is perpendicular to that of light source polarizer) to reduce reflexion and glare. The same specimens were also imaged: 1) under UV-light for those presenting a UV-yellow fluorescence; and 2) covered in 95% ethanol to see cuticular remains with a low angle light to reveal relief. The palaeontological arrangement of this study follows the systematics proposed by Karasawa *et al.* (2013).

SYSTEMATIC PALAEONTOLOGY

Class MALACOSTRACA Latreille, 1802
 Order DECAPODA Latreille, 1802
 Suborder PLEOCYEMATA Burkenroad, 1963
 Infraorder ASTACIDEA Latreille, 1802
 Superfamily STENOCHIROIDEA Beurlen, 1928
 Family STENOCHIRIDAE Beurlen, 1928

Genus *Pseudastacus* Oppel, 1861

Pseudastacus Oppel, 1861b: 360.

TYPE SPECIES. — *Bolina pustulosa* Münster, 1839, by subsequent designation of Glaessner (1929: 350).

INCLUDED SPECIES. — Schweitzer *et al.* (2010) listed five species included in *Pseudastacus*: *P. bakelensis* Fraas, 1878 (Cenomanian, Lebanon), *P. minor* Fraas, 1878 (Cenomanian, Lebanon), *P. mucronatus* (Phillips, 1835) (Hauterivian, United Kingdom), *P. pusillus* Van Straelen, 1925 (Bajocian, France), and *P. pustulosus* (Münster, 1839) (Tithonian, Germany).




FIG. 2. — Holotype CN-NMP-130-1 of *Pseudastacus lemovices* n. sp. from the Sinemurian of Chauffour-sur-Vell, France: **A**, specimen in right lateral view, cross-polarized light; **B**, interpretative line drawing. Abbreviations: **a**, branchiocardiac groove; **b**, antennal groove; **b₁**, hepatic groove; **ca**, carpus; **di**, diaeresis; **en**, uropodal endopod; **ex**, uropodal exopod; **e,e**, cervical groove; **pr**, propodus; **P1-P4**, pereopods 1 to 4; **me**, merus; **r**, rostrum; **sc**, scaphocerite; **se**, setae; **s1-s6**, pleonal somites 1 to 6. Photograph: D. Audo. Scale bars: 5 mm.

TAXONOMIC COMMENTS

Pseudastacus bakelensis was moved to *Notahomarus* by Charbonnier *et al.* (2017). The type material of *Pseudastacus minor* is lost (G. Schweigert pers. comm. 2019) and the illustration proposed by Fraas (1878: pl. 4, fig. 2) is not conclusive concerning the taxonomic determination. Indeed, his illustration is relatively strange for the following reasons: 1) either there is a wrong number of pleonal somites (7 instead of 6 excluding the telson), or the first somite is rather large; 2) the rostrum is exceedingly long in comparison with the rest of the carapace; and 3) as the carapace and pleon seem to be in connection suggesting that the specimen was a corpse, the first chelipeds appear strangely placed and should be positioned more anteriorly; in addition, the general morphology of the chelae does not fit with that observed in *Pseudascatus*.

The type material of *Pseudastacus pusillus* was destroyed during World War II (see Bigot 1945) and the figure proposed by Van Straelen (1925: pl. 9, fig. 7) is not conclusive. Even if the line drawing provided by Van Straelen (1925: fig. 134) shows the carapace groove pattern, it is difficult to confirm the assignation to *Pseudastacus*, which was already proposed with a question mark in the original publication.

The type material of *Pseudastacus mucronatus* corresponds to a fragment of P1 chela figured by Phillips (1829: pl.3, fig. 3) and Phillips (1835: pl. 3, fig. 3), and refigured by Bell (1863: pl. 9, fig. 6). This isolated chela is very massive and the occlusal margins of index and dactylus show very strong tubercles alternating with smallest ones. This general morphology is not consistent with the more elongate and thinner P1 chelae observed in *Pseudascatus*. Following Woods (1931: 89), this P1 chela might be referred to *Hoploparia dentata* (Roemer, 1841).




FIG. 3. — Paratype CN-NMP-130-2 of *Pseudastacus lemovices* n. sp. from the Sinemurian of Chauffour-sur-Vell, France: **A**, specimen in right lateral/ventral view, cross-polarized light; **B**, interpretative line drawing. Abbreviations: a2, antenna; ca, carpus; pr, propodus; P1-P5, pereopods 1 to 5; me, merus; mxp3, third maxilliped; s1-s6, pleonal somites 1 to 6. Photograph: D. Audo. Scale bars: 5 mm.

Pseudastacus lemovices n. sp.
(Figs 2-6)

urn:lsid:zoobank.org:act:D58E5862-AD82-4B39-96D4-56BED134F76D

ETYMOLOGY. — The specific epithet refers to the “Lemovices” (used as a noun in apposition), a Gaulish tribe of Central Europe who established themselves in Limousin and Poitou regions between 700 and 400 BC. The type locality Chauffour-sur-Vell is located in Limousin.

TYPE MATERIAL. — **Holotype**: CN-NMP-130-1.

Paratypes: 4, CN-NMP-130-2 to CN-NMP-130-5. All are preserved in compression at the surface of a single slab of limestone.

TYPE LOCALITY. — Chauffour-sur-Vell, Corrèze department, Limousin, Nouvelle-Aquitaine region, France.

TYPE AGE. — Early Jurassic, Sinemurian (after Lefavrais-Raymond *et al.* 1976).

DESCRIPTION

Carapace

Subcylindrical carapace (length, excluding rostrum: *c.* 11 mm; height: *c.* 6.5 mm); straight dorsal margin; ventral margin poorly preserved; triangular rostrum, elongate, relatively large, with pointed distal extremity, bearing three lateral spines; deep cervical groove, subvertical at level of dorsal margin and ventrally curved before joining concave antennal groove; sinuous branchiocardiac groove joined ventrally to rounded hepatic groove; large and deep orbital incision, ventrally limited by antennal spine.

Pleon

Somite 1, smallest than the others; somites 1 to 5 (s1-s5) with rounded pleura, almost straight anteriorly and concave posteriorly; s2-s5 with longitudinal carina separating tergopleuron from the rest of the tergum; s6 shorter than s5; telson poorly preserved.




FIG. 4. — Paratype CN-NMP-130-3 of *Pseudastacus lemovices* n. sp. from the Sinemurian of Chauffour-sur-Vell, France: **A**, specimen in left lateral/dorsal view, cross-polarized light; **B**, interpretative line drawing. Abbreviations: **di**, diaeresis; **en**, uropodal endopod; **ex**, uropodal exopod; **e,e**, cervical groove; **P1-P3**, pereiopods 1 to 3; **r**, rostrum; **se**, setae; **sk**, endophragmal skeleton; **s1-s4**, pleon somites 1 to 4. Photograph: D. Audo. Scale bars: 5 mm.

Cephalic appendages

Rounded eye (preserved only on paratype CN-NMP-130-4); antennulae (a1) poorly preserved, short, with multi-articulated flagella; antennae (a2) composed of elongate antennal peduncle bearing large, elongate, and pointed scaphocerite, and strong multi-articulated flagella; third maxilliped, elongate and smooth;

Thoracic appendages

Pereiopods 1 to 5 (P1-P5) decreasing in size posteriorly; chelate pereiopods 1 to 3 (P1-P3); P1 larger and longer than P2 and P3; short and stout P1 chela with index and dactylus of equal length, with slightly curved distal extremity and finely serrated occlusal margins; dorsal margin of P1 merus with one longitudinal median row of strong tubercles; P1 carpus

and propodus with smooth dorsal margin; small P2 and P3 chelae with index and dactylus of equal length, slightly curved distal extremity; P4 and P5, smallest and smooth, poorly preserved.

Pleonal appendages

Pleopods not preserved; uropods of equal length, with distal margins fringed by elongate setae; uropodal exopod with longitudinal median carina and straight diaeresis.

Ornamentation

Carapace uniformly covered by small pits; pereiopods smooth; pleon somites smooth. Some specimens show a relatively smooth carapace. Thus, the carapace ornamentation is variable in aspect, according to which of the cuticle layers are



FIG. 5. — Paratype CN-NMP-130-4 of *Pseudastacus lemovices* n. sp. from the Sinemurian of Chaffour-sur-Vell, France: **A**, specimen in left lateral view, cross-polarized light; **B**, interpretative line drawing. Abbreviations: **a**, branchiocardiac groove; **as**, antennal spine; **a1**, antennula; **a2**, antenna; **b**, antennal groove; **b1**, hepatic groove; **e1e**, cervical groove; **o**, eye; **r**, rostrum; **sc**, scaphocerite; **s1-s6**, pleonal somites 1 to 6; **t**, telson. Photograph: D. Audo. Scale bars: 5 mm.

preserved: smooth or finely pitted; both aspects may in fact correspond to diverse degrees of abrasion and/or decortication of carapace (similar cases were observed in erymid lobsters: see Charbonnier *et al.* 2014; Hyžný *et al.* 2015).

DISCUSSION

The new species possesses anatomical characters typical of Astacidea: the first three pereiopods are chelate, the first pereiopods are stronger than the others, there are no mid-lateral hinges between the pleonal somites, the cervical groove is deep and the branchiocardiac groove is oblique. Moreover, following the key proposed by Wahle *et al.* (2012), the new species is assigned to the family Stenochiridae based upon the following characters: rostrum bearing lateral spines; cervical groove extends transversely from dorsomedian; branchiocardiac groove extends obliquely across lateral portion of carapace; uropodal exopod with

diaeresis; pereiopods 2 and 3 bearing very small chelae. More precisely, the new species is assigned to *Pseudastacus* Oppel, 1861b based on the typical groove pattern of the carapace (deep cervical groove crossing dorsal midline; well-developed branchiocardiac groove, joined ventrally to hepatic groove), the elongate triangular rostrum with three lateral spines; the short and stout P1 chela. *Pseudastacus lemovices* n. sp. differs from *P. pustulosus* by the ornamentation of the carapace (uniformly covered by tubercles in *P. pustulosus*; uniformly covered by pits in *P. lemovices* n. sp.) and of the first pereiopod (uniformly covered by tubercles in *P. pustulosus*; smooth in *P. lemovices* n. sp.).

In addition, this association of five specimens may suggest a mass mortality event linked to an environmental stress (e.g., anoxia, temperature). It may also point to a gregarious behaviour as observed in different extant and fossil lobsters (see Klompaker & Fraaije 2012 and references therein).




FIG. 6. — Paratype CN-NMP-130-5 of *Pseudastacus lemovices* n. sp. from the Sinemurian of Chauffour-sur-Vell, France: **A**, specimen in ventral view, cross-polarized light; **B**, interpretative line drawing. Abbreviations: **e₁e**, cervical groove; **md**, dorsal midline; **P2-P4**, pereiopods 2 to 4; **r**, rostrum; **sc**, scaphocerite; **s1-s2**, pleonal somites 1 to 2. Photograph: D. Audo. Scale bars: 5 mm.

CONCLUSIONS

Pseudastacus Oppel, 1861b, was placed for long time in the family Nephropidae Dana, 1852 as reported by Van Straelen (1925), Beurlen (1928), Glaessner (1929, 1969), and Chong & Förster (1976). Albrecht (1983) included *Pseudastacus* in his new family Proastacidae, whereas Tshudy & Babcock (1997) included it in their new family Chilenophoberidae. Garassino & Schweigert (2006) preferred to

follow Albrecht (1983) considering the analysis of Tshudy & Babcock (1997) was not founded by a morphological review of the type material, but only on line drawings in literature. Later, De Grave *et al.* (2009) and Schweitzer *et al.* (2010) maintained *Pseudastacus* in the Chilenophoberidae. Finally, Karasawa *et al.* (2013) proposed to synonymize the Chilenophoberidae with the Stenochiridae Beurlen, 1928, and to place *Pseudastacus* in this latter family. Here we follow this placement.

Until now, stenochirid lobsters had only representatives in the Middle Jurassic of France and Germany, in the Late Jurassic of France, Germany and Chile, in the Early Cretaceous of Australia, and in the Late Cretaceous of Japan (for details see Table 1). In conclusion, the Early Jurassic *Pseudastacus lemovices* n. sp. is the oldest representative of the genus and of the family.

Acknowledgements

We thank the two reviewers, Alessandro Garassino (Milano Museum) and Günter Schweigert (Stuttgart Museum), for their useful comments on our manuscript. We are pleased to acknowledge Mrs Boutang and Mr Gomme who collected the specimen and donated it to the local museum of Noailhac (Corrèze department, France) and Mr Guy Chantepie, who called our attention to the new specimen. We wish to thank Mr Christian Lassalle, mayor of Noailhac, who made the specimen available and carried it to the MNHN, Paris for our study. We are grateful to Philippe Loubry and Lilian Cazes (UMR 7207 CR2P, MNHN, Paris) for the photographic work.

Denis Audo's research was supported by the China Postdoctoral Science Foundation, Yunnan Province Postdoctoral Science Foundation, NSFC grant 41861134032, and Yunnan Provincial Research Grants 2018FA025 and 2018IA073.

REFERENCES

- ALBRECHT H. 1983. — Die Protoastacidae n. fam., fossile Vorfahren der Flußkrebse? *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1983 (1): 5-15.
- BELL T. 1863. — *A Monograph of the Fossil Malacostracous Crustacea of Great Britain, Part II, Crustacea of the Gault and Greensand*. The Palaeontographical Society, London, i-viii, 1-40. <https://doi.org/10.1080/02693445.1863.12027935>
- BEURLEN K. 1928. — Die Decapoden des Schwäbischen Jura mit Ausnahme der aus den oberjurassischen Plattenkalken stammenden. *Palaeontographica* 70: 115-278.
- BIGOT A. 1945. — La destruction des collections et des bibliothèques scientifiques de Caen. *Bulletin de la Société linnéenne de Normandie*, Volume supplémentaire: 1-75.
- CHARBONNIER S. 2009. — *Le Lagerstätte de La Voulte : un environnement bathyal au Jurassique*. Muséum national d'Histoire naturelle, Paris, 272 p. (*Mémoires du Muséum national d'Histoire naturelle* ; 199).
- CHARBONNIER S., AUDO D., GARASSINO A. & HYŽNÝ M. 2017. — *Fossil Crustacea of Lebanon*. Muséum national d'Histoire naturelle, Paris, 252 p. (*Mémoires du Muséum national d'Histoire naturelle* ; 210).
- CHARBONNIER S., GARASSINO A., SCHWEIGERT G., AUDO D., FERNANDEZ S. 2014. — New look at the lobster *Eryma greppini*, OPPEL, 1861 (Crustacea, Decapoda, Erymidae) from the Middle Jurassic of France and Switzerland. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen* 272: 331-339. <https://doi.org/10.1127/0077-7749/2014/0411>
- CHONG G. & FÖRSTER R. 1976. — *Chilenophoberus atacamensis*, a new decapod crustacean from the Middle Oxfordian of the Cordillera de Domeyko, northern Chile. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 1976 (3): 145-156.
- DE GRAVE S., PONTCHEFF N. D., AHYONG S. T., CHAN T.-Y., CRANDALL K. A., DWORSCHAK P. C., FELDER D. L., FELDMANN R. M., FRANSEN C. H. M., GOULDING L. Y. D., LEMAITRE R., LOW M. E. Y., MARTIN J. W., NG P. K. L., SCHWEITZER C. E., TAN S. H., TSHUDY D. & WETZER R. 2009. — A classification of living and fossil genera of decapod crustaceans. *The Raffles Bulletin of Zoology*, Supplement 21: 1-109. <http://hdl.handle.net/10088/8358>
- FRAAS O. 1878. — Geologisches aus dem Libanon. *Jahreshefte des Vereins für Vaterländische Naturkunde in Württemberg* 34: 257-391.
- GARASSINO A. & SCHWEIGERT G. 2006. — The Upper Jurassic Solnhofen decapod crustacean fauna: review of the types from old descriptions. Part I. Infraorders Astacidea, Thalassinidea, and Palinura. *Memorie della Società italiana di Scienze naturali e del Museo civico di Storia naturale di Milano* 34 (1): 1-64.
- GLAESSNER M. F. 1929. — Crustacea Decapoda, in POMPECKJ J. F. (ed.), *Fossilium Catalogus, I: Animalia*, Pars 41: 1-464.
- GLAESSNER M. F. 1932. — Zwei ungenügend bekannte mesozoische Dekapodenkrebse. *Pemphix sueuri* (Desm.) und *Palaeophoberus suevicus* (Quenstedt). *Paläontologische Zeitschrift* 14: 108-121. <https://doi.org/10.1007/BF03041621>
- GLAESSNER M. F. 1969. — Decapoda, in MOORE R. C. (ed.), *Treatise on Invertebrate Paleontology*, Part R, *Arthropoda* 4 (2): 399-533.
- HYŽNÝ M., SCHLÖGL J., CHARBONNIER S., SCHWEIGERT G., RULLEAU L. & GOUTTENNOIRE M. 2015. — Intraspecific variation and taphonomy of a new erymid lobster (Crustacea: Decapoda) from the Middle Jurassic of Belmont (Beaujolais, France). *Geobios* 48: 371-384. <https://doi.org/10.1016/j.geobios.2015.07.006>
- KARASAWA H., SCHWEITZER C. E. & FELDMANN R. M. 2013. — Phylogeny and systematics of extant and extinct lobsters. *Journal of Crustacean Biology* 33 (1): 78-123. <https://doi.org/10.1163/1937240X-00002111>
- KLOMPMAKER A. A. & FRAAIJE R. H. B. 2012. — Animal Behavior Frozen in Time: Gregarious Behavior of Early Jurassic Lobsters within an Ammonoid Body Chamber. *PLoS ONE* 7(3): e31893. <https://doi.org/10.1371/journal.pone.0031893>
- LEFAVRAIS-RAYMOND A., FEYS R., BOISSONNAS J., TALBERT J.-C., GUYONNAUD G., VOGT J. & RAYNAL J.-P. 1976. — Carte géologique de la France à 1:50 000. Brive-la-Gaillarde. Bureau de recherches géologiques et minières, Orléans, (France), carte 785.
- MÜNSTER G. GRAF ZU 1839. — Decapoda Macrura. Abbildung und Beschreibung der fossilen langschwänzigen Krebse in den Kalkschiefern von Bayern. *Beiträge zur Petrefaktenkunde* 2: 1-88.
- OPPEL A. 1861a. — Die Arten der Gattungen *Glyphea* und *Pseudoglyphea*. *Jahreshefte des Vereins für Vaterländische Naturkunde in Württemberg* 17: 108-111.
- OPPEL A. 1861b. — Die Arten der Gattungen *Eryma*, *Pseudastacus*, *Magila* und *Etallonia*. *Jahreshefte des Vereins für Vaterländische Naturkunde in Württemberg* 17: 355-361.
- OPPEL A. 1862. — Ueber jurassische Crustaceen (Decapoda macrura). *Paläontologische Mittheilungen aus dem Museum des königlich Bayerischen Staates* 1: 1-120.
- PHILLIPS J. 1829. — *Illustrations of the Geology of Yorkshire*. Thomas Wilson and sons, York, 192 p. <https://doi.org/10.5962/bhl.title.30592>
- PHILLIPS J. 1835. — *Illustrations of the Geology of Yorkshire. Part 1. The Yorkshire Coast*. 2nd Edition. John Murray, London, 184 p. <https://doi.org/10.5962/bhl.title.127948>
- QUENSTEDT F. A. 1867. — *Handbuch der Petrefaktenkunde*. 2nd edition. Laupp, Tübingen, 982 p. <https://gallica.bnf.fr/ark:/12148/bpt6k97217s>
- ROGER J. & LAPPARENT A. F. DE 1944. — Une nouvelle espèce de crustacé décapode *Palaeophoberus portlandicus*, découverte dans le Portlandien du Pays de Bray. *Bulletin de la Société géologique de France* 14: 365-374.
- SCHWEIGERT G., GARASSINO A. & RIOU B. 2006. — First record of *Stenochirus* Opele, 1861 (Crustacea: Decapoda: Stenochiridae) from the Callovian (Middle Jurassic) of La Voulte-sur-Rhône. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* 2006: 65-77. <https://doi.org/10.1127/njgpm/2006/2006/65>
- SCHWEITZER C. E., FELDMANN R. M., GARASSINO A., KARASAWA H. & SCHWEIGERT G. 2010. — Systematic list of fossil decapod crustacean species. *Crustaceana Monographs* 10: 1-222. <https://doi.org/10.1163/ej.9789004178915.i-222>

- THSUDY D. & BABCOCK L. E. 1997. — Morphology-based phylogenetic analysis of the clawed lobsters (family Nephropidae and the new family Chilenophoberidae). *Journal of Crustacean Biology* 17: 253-263. <https://doi.org/10.1163/193724097X00288>
- VAN STRAELEN V. 1925. — Contribution à l'étude des crustacés décapodes de la période jurassique. *Mémoires de la Classe des Sciences de l'Académie royale de Belgique* 7: 1-462.
- WAHLE R. A., THSUDY D., COBB J. S., FACTOR J. & JAINI M. 2012. — Infraorder Astacidea Latreille, 1802 p.p.: The Marine Clawed Lobsters, in SCHRAM F. R., VAUPEL KLEIN J. C. VON, CHARMANTIER-DAURES M., FOREST J. (eds), *Treatise of Zoology – Anatomy, Taxonomy, Biology – The Crustacea*. Volume 9, Part B. *Eucarida, Astacidea P. P. (Enoplometopoidea, Nephropoidea), Glypheidea, Axiidea, Gebiidea, and Anomura*. Brill, Leiden & Boston: 3-108. https://doi.org/10.1163/9789047430179_003
- WOODS H. 1925-1931. — *A Monograph of the Fossil Macrurous Crustacea of England*. Palaeontographical Society of London, London, 122 p.
- WOODS J. T. 1957. — Macrurous Decapods from the Cretaceous of Queensland. *Memoirs of the Queensland Museum* 13 (3): 155-175. <http://www.biodiversitylibrary.org/page/48733649>
- YOKOI T. & KARASAWA H. 2000. — Additions to Cretaceous decapod crustaceans from Hokkaido, Japan – Part 2. Chilenophoberidae. *Bulletin of the Mizunami Fossil Museum* 27: 191-194.

Submitted on 16 July 2019;
accepted on 23 July 2019;
published on 26 March 2020.