



Food and Agriculture
Organization of the
United Nations

DIETARY ASSESSMENT

A RESOURCE GUIDE TO METHOD SELECTION
AND APPLICATION IN LOW RESOURCE SETTINGS



DIETARY ASSESSMENT

**A RESOURCE GUIDE TO METHOD SELECTION
AND APPLICATION IN LOW RESOURCE SETTINGS**

Food and Agriculture Organization of the United Nations
Rome, 2018

Recommended citation

FAO. 2018. Dietary Assessment: A resource guide to method selection and application in low resource settings. Rome.

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO in preference to others of a similar nature that are not mentioned.

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views or policies of FAO.

ISBN 978-92-5-130635-2

© FAO, 2018

FAO encourages the use, reproduction and dissemination of material in this information product. Except where otherwise indicated, material may be copied, downloaded and printed for private study, research and teaching purposes, or for use in non-commercial products or services, provided that appropriate acknowledgement of FAO as the source and copyright holder is given and that FAO's endorsement of users' views, products or services is not implied in any way. All requests for translation and adaptation rights, and for resale and other commercial use rights should be made via www.fao.org/contact-us/licencerequest or addressed to copyright@fao.org.

FAO information products are available on the FAO website (www.fao.org/publications) and can be purchased through publications-sales@fao.org.

Cover photo credits:

Background photo:
©FAO/Ado Youssouf

Front cover photos:
©FAO/Ruth Charrondiere
©FAO/Alessandra Benedetti
©FAO/Warren Lee
©FAO/Giulio Napolitano

Back cover photos:
©FAO/Oliver Bunic
©FAO/Warren Lee
©FAO/Ami Vitale
©FAO/Ruth Charrondiere

CONTENTS

1. INTRODUCTION	1
2. DIETARY ASSESSMENT METHODS	3
2.1 Indirect methods	3
2.1.1 Food Balance Sheets – national food availability	4
2.1.2 Household Consumption and Expenditure Surveys – household food consumption	7
2.2 Direct methods	10
2.2.1 Retrospective direct methods	10
2.2.1.1 Food frequency questionnaire	10
2.2.1.2 24-hour recall	18
2.2.1.3 Dietary history	24
2.2.2 Prospective direct methods	28
2.2.2.1 Estimated food records	28
2.2.2.2 Weighed food records	33
2.2.2.3 Duplicate meal method	37
2.3 Integration of innovative technologies to improve dietary assessment	39
2.3.1 Personal Digital Assistant (PDA)	40
2.3.2 Image-assisted dietary assessment methods	40
2.3.3 Mobile-based technologies	41
2.3.4 Interactive computer and web-based technologies	43
2.3.5 Scan- and sensor-based technologies	43
2.3.6 Applications and uses of innovative technologies to improve dietary assessment	44
2.3.7 Validation of innovative technologies to improve dietary assessment	45
2.3.8 Strengths and limitations of innovative technologies to improve dietary assessment methods	46
2.4 Qualitative retrospective proxy tools for assessing dietary diversity	51
2.4.1 Minimum Dietary Diversity –Women (MDD-W)	52
2.4.2 Infant and Young Child Dietary Diversity Score (IYCCDS)	53
2.4.3 Applications and uses of individual level dietary diversity score	54
2.4.4 Validity of individual level dietary diversity score	55
2.4.5 Strengths and limitations of individual level dietary diversity scores	56
3. METHODOLOGICAL CONSIDERATIONS	59
3.1 Sources of dietary variation	59
3.2 Individual, community and culturally-specific issues in low resource settings	61
3.3 Intra-household food distribution, shared eating occasions and street food	63
3.4 Estimation of portion size	63
3.5 Availability of food composition data	64

3.6 Measurement errors in dietary assessment	65
3.6.1 Misreporting energy intakes	68
3.7 Reproducibility in dietary assessment	69
3.8 Validity in dietary assessment	70
3.9 Quality control and data analysis on dietary assessment	76
4. SELECTING A DIRECT DIETARY ASSESSMENT METHOD	79
4.1 Study objectives in dietary assessment	79
4.2 Design of the study	81
4.3 Technical and financial aspects to take into account	82
4.4 Supporting information for the selection of a direct dietary assessment method	82
4.4.1 A step-by-step guide for method selection	82
4.4.2 Summary of the major features of the different direct dietary assessment methods	84
4.4.3 Case studies on selection of a dietary assessment method	90
5. KEY MESSAGES AND THE WAY FORWARD IN DIETARY ASSESSMENT	93
5.1 Key messages	93
5.2 The way forward	95
6. FURTHER READING	97
7. REFERENCES	99
8. APPENDICES	115
APPENDIX 1: EXAMPLES OF FOOD FREQUENCY QUESTIONNAIRE (FFQ)	115
8.1 Example 1A: Filled-out qualitative FFQ	115
8.2 Example 1B: Filled-out semi-quantitative FFQ	121
APPENDIX 2: AN EXAMPLE OF BRIEF DIETARY QUESTIONNAIRE	128
8.3 Example 2: Filled-out brief dietary questionnaire	128
APPENDIX 3: AN EXAMPLE OF 24-HOUR RECALL	130
8.4 Example 3: Filled-out 24-hour recall*	130
APPENDIX 4: AN EXAMPLE OF DIETARY HISTORY	133
8.5 Example 4: Filled-out dietary history*	133

APPENDIX 5: AN EXAMPLE OF FOOD RECORD	145
8.6 Example 5: Filled-out three day food record	145
APPENDIX 6: AN EXAMPLE OF MINIMUM DIETARY DIVERSITY – WOMEN (MDD-W) QUESTIONNAIRE	148
8.7 Example 6: Filled-out MDD-W questionnaire	148

TABLES

TABLE 1 - STRENGTHS AND LIMITATIONS OF USING FBS DATA FOR ASSESSING DIETS	6
TABLE 2 - STRENGTHS AND LIMITATIONS OF USING HCES DATA FOR ASSESSING DIETS	9
TABLE 3 - STRENGTHS AND LIMITATIONS OF FFQ	14
TABLE 4 - STRENGTHS AND LIMITATIONS OF 24-HOUR RECALL	21
TABLE 5 - STRENGTHS AND LIMITATIONS OF DIETARY HISTORY METHOD	26
TABLE 6 - STRENGTHS AND LIMITATIONS OF ESTIMATED FOOD RECORDS	30
TABLE 7 - STRENGTHS AND LIMITATIONS OF WEIGHED FOOD RECORDS	35
TABLE 8 - STRENGTHS AND LIMITATIONS OF DUPLICATE MEAL METHOD	38
TABLE 9 - STRENGTHS AND LIMITATIONS OF INNOVATIVE TECHNOLOGIES TO IMPROVE DIETARY ASSESSMENT METHODS	47
TABLE 10 - STRENGTHS AND LIMITATIONS OF INNOVATIVE TECHNOLOGIES INTEGRATED INTO CONVENTIONAL DIETARY ASSESSMENT METHODS	49
TABLE 11 - STRENGTHS AND LIMITATIONS OF INDIVIDUAL LEVEL DIETARY DIVERSITY SCORE	56
TABLE 12 - SOURCES OF ERRORS IN DIRECT DIETARY ASSESSMENT METHODS FOR ASSESSING FOOD AND NUTRIENT INTAKES	66
TABLE 13 - EXAMPLES OF PAIRING TEST DIETARY ASSESSMENT METHODS AND REFERENCE METHODS	71
TABLE 14 - SELECTION OF A METHOD TO MEASURE NUTRIENT INTAKES TO MEET FOUR POSSIBLE LEVELS OF OBJECTIVES	80
TABLE 15 - DIETARY ASSESSMENT METHODS COMMONLY USED IN THE DESIGN OF DIFFERENT STUDIES	81
TABLE 16 - COMPARISON OF THE MAJOR FEATURES OF METHODS FOR ASSESSING DIET	85
TABLE A - DESCRIPTIVE QUALITATIVE FFQ	116
TABLE B - SEMI-QUANTITATIVE FFQ	121
TABLE C - A BRIEF DIETARY QUESTIONNAIRE	128
TABLE D - A 24-HOUR RECALL	130
TABLE E - A DIETARY HISTORY RECORD	133
TABLE F - A THREE DAY FOOD RECORD	145
TABLE G - A 24-HOUR RECALL	149
TABLE H - TABLE OF FOOD GROUPS	150
TABLE I - THE 10 FOOD GROUPS	152

FIGURES

FIGURE 1 - OVERVIEW OF DIETARY ASSESSMENT METHODS TO ESTIMATE FOOD AND NUTRIENT CONSUMPTION AT NATIONAL, HOUSEHOLD AND INDIVIDUAL LEVEL	4
FIGURE 2 - DIAGRAM OF THE TECHNOLOGY ASSISTED DIETARY ASSESSMENT (TADA) SYSTEM THAT STARTS WITH CAPTURING AN IMAGE WITH THE MOBILE FOOD RECORD (MFR)	42
FIGURE 3 - DIFFERENCE IN MEAN IRON INTAKE ESTIMATED BY A 24-HOUR RECALL AND A WEIGHED FOOD RECORD	75

BOXES

Box 1. QUICK GUIDE TO USING AN FFQ	15
Box 2. BRIEF DIETARY ASSESSMENT	17
Box 3. QUICK GUIDE TO USING A 24-HOUR RECALL	21
Box 4. QUICK GUIDE TO USING A DIETARY HISTORY METHOD	27
Box 5. QUICK GUIDE TO USING AN ESTIMATE FOOD RECORD	31
Box 6. QUICK GUIDE TO USING A WEIGHED FOOD RECORD	35
Box 7. QUICK GUIDE TO USING A DUPLICATE MEAL METHOD	39
Box 8. QUICK GUIDE TO USING INNOVATIVE TECHNOLOGIES FOR DIETARY ASSESSMENT	50
Box 9. QUICK GUIDE TO USING INDIVIDUAL LEVEL DIETARY DIVERSITY SCORES	57
Box 10. STEPS AND TIPS ON CHOOSING A DIETARY ASSESSMENT METHOD	83

FOREWORD

Across the world today, there is increasing interest in incorporating robust nutrition information into national information systems. The aim is to inform the implementation and evaluation of nutrition-sensitive agricultural projects, policies and programmes, and to tackle all forms of malnutrition. The need for such robust information was reaffirmed at the Second International Conference on Nutrition (ICN2) in November 2014. It is therefore important that as an organization, FAO works to meet global knowledge demands and gaps in decision-making, by supporting the collection of nutrition information for surveillance, setting targets, measuring impacts, and tracking progress.

Up-to-date and valid assessment of what people eat and drink will help to generate better information and evidence that will contribute to the formulation of effective agricultural and nutrition policies and programmes. It will also benefit consumer education, which in turn will contribute to raising levels of nutrition and help to prevent undernutrition, obesity and non-communicable diseases. This increasingly rigorous approach will lead to a culture of robust dietary data collection, resulting in evidence-based decisions that are crucial to achieving the strategic objectives of the organization.

This resource guide provides an updated overview of the dietary assessment methods that can be used to collect dietary data at national, household and individual levels. The strengths and limitations of various methods are discussed in detail. Its particular focus on low resource settings makes it a valuable tool for users working in environments where resources are limited and rapid nutritional changes might take place. Taking advantage of the proliferation of digital technologies, methodologies involving the use of interactive and web-based technologies for dietary data collection have also been reviewed.

The guide will be a useful resource for programme managers, educators, health care professionals, health promotion specialists, students, extension workers and researchers: in short, anyone involved in food consumption surveys, programme planning, implementation, monitoring or evaluation. It is a one-stop shop for selecting the most appropriate methods for different contexts.

Anna Lartey
Director

Nutrition and Food Systems Division
Food and Agriculture Organization of the United Nations
Rome, Italy



ACKNOWLEDGEMENTS

The resource guide was initially conceptualized and developed by Warren T. K. Lee with contributions from Queenie W. Y. Mak, Nutrition and Food Systems Division, FAO, at an early stage. Warren T. K. Lee, Theodora Mouratidou, Claudia E. Lazarte and Saba Marzara were subsequently responsible for further developing, revising and finalizing the resource guide, and Giorgia Paratore contributed to the development of the Appendices. The advanced draft of the resource guide was peer-reviewed by the following experts on dietary assessment: Jennifer Coates (Tufts University Friedman School of Nutrition Science and Policy, USA), Brooke Colaiezzi (Tufts University Friedman School of Nutrition Science and Policy, USA), Rosalind Gibson (Department of Human Nutrition, University of Otago, New Zealand) and Pattanee Winichagoon (Institute of Nutrition, Mahidol University, Thailand). From within FAO, Pauline Allemand (Nutrition and Food Systems Division), Piero Conforti (Statistics Division), Catherine Leclercq (Nutrition and Food Systems Division), Ana Moltedo (Statistics Division), Salar Tayyib (Statistics Division) and Nathalie Troubat (Statistics Division) acted as internal reviewers. Additionally, we would like to thank Verena Nowak who provided comments on the draft during the initial drafting process, Viviana Panetti, Cristiana Fusconi and Ji Yen Alexandra Tung (Nutrition and Food Systems Division) for their administrative and editorial support on the final version. The publication of the resource guide was made possible with financial support from the European Union and FAO. Final editing by Anthony Jennings, design and layout by Luca Pierotti.

ABBREVIATIONS AND ACRONYMS

ADePT-FSM	ADePT Software
AMPM	Automated Multiple-Pass Method
ANOVA	Analysis Of Variance
ASA24	Automated Self-Administered 24-hour Recall
BMR	Basal Metabolic Rate
CARDIA	Coronary Artery Risk Development in Young Adults
DAFNE	Data Food Networking
DDS	Dietary Diversity Score
DISHES	Dietary Interview Software for Health Examination Studies
DLW	Doubly Labelled Water
DNSIYC	National Survey of Infant and Young Children
EI	Energy Intake
EPIC	European Prospective Investigation into Cancer
ESN	Nutrition and Food Systems Division, FAO
ESNA	Nutrition Assessment and Scientific Advice Group, FAO
EU	European Union
FAO	Food and Agriculture Organization of the United Nations
FAOSTAT	FAO Statistical Databases on Food and Agriculture
FAO/WHO GIFT	FAO/WHO Global Individual Food Consumption Data Tool
FBS	Food Balance Sheet
FCD	Food Composition Database
FCS	Food Consumption Score
FFQ	Food Frequency Questionnaire
FP 24-hR	Food Photography 24-hour Recall
GDD	Global Dietary Database
HAZ	Height-for-Age Z-Scores
HBS	Household Budget Survey
HCES	Household Consumption and Expenditure Survey
HDDS	Household Dietary Diversity Score
HES	Household Expenditure Survey
HIES	Household Income and Expenditure Survey
ICN2	Second International Conference on Nutrition
IHS	Integrated Household Survey

IHSN	International Household Survey Network
INFOODS	International Network of Food Data Systems
IYCDDS	Infant and Young Child Dietary Diversity Score
LCFS	Living Costs and Food Survey
LSMS	Living Standards Measurement Studies
MDD-W	Minimum Dietary Diversity – Women
mFR	Mobile Food Record
MMDA	Mean Micronutrient Density Adequacy
MPA	Mean Probability of Micronutrient Adequacy
NANA	Novel Assessment of Nutrition and Ageing
NDNS	National Diet and Nutrition Survey
PAL	Physical Activity Level
PDA	Personal Digital Assistant
PRA	Participatory Rural Appraisal
TADA	Technology-Assisted Dietary Assessment
TEE	Total Energy Expenditure
USDA	United States Department of Agriculture
WDDS	Women’s Dietary Diversity Score
WHO	World Health Organization

KEY TECHNICAL TERMS AND NOTATIONS¹

A

Analysis of Variance (ANOVA) is a statistical analysis that allows for the comparison of means in more than two groups or in groups defined by more than one qualitative variable.

Anthropometric measurements are measurements of the variation of the physical dimensions (i.e. length, height, weight, weight-for-length, mid-arm circumference head circumference, etc.) and the gross composition (i.e. body fat, fat-free mass) of the human body.

B

Basal Metabolic Rate is the minimal rate of energy expenditure required to sustain life. It is measured in the supine position when the individual is in a state of rest (but not sleeping), mental relaxation, fasted, and in a neutrally temperate environment. It is the largest component of total energy expenditure, typically 60–75 percent when measured over 24 hours.

Biomarkers refer to a chemical, its metabolite, or the product of an interaction between a chemical and some target molecule or cell that is commonly measured in body fluids (blood, serum, urine) and tissue to perform a clinical assessment and/or monitor and predict health and disease states in individuals or across populations and in validation studies.

Bland–Altman plot also known as difference plot in analytical chemistry and biostatistics is a method of data plotting used in analysing the agreement between two different assays.

C

Chi-squared test is a statistical test for categorical variables commonly used to compare observed data with data we would expect to obtain according to a specific hypothesis. The hypothesis states that there is no significant difference between the expected and observed result.

Correlation coefficient is a measure of the interdependence of two random variables that ranges in value from -1 to +1, indicating perfect negative correlation at -1, absence of correlation at zero, and perfect positive correlation at +1. Also called coefficient of correlation.

Covariate is a variable that may be predictive of the outcome under study. A covariate may be of direct interest or it may be a confounder or effect modifier.

Cross-classification is a classification according to more than one attribute at the same time; e.g. the cross-classification of cases was done by age and sex.

D

Doubly labelled water method is a stable isotopic technique for measuring energy expenditure in free-living subjects, it is used to identify underreporting of total energy intake.

¹ The authors used the following sources to compile the list of terms. For further information, users are directed to the original sources. Dietary Assessment Primer, National Institutes of Health, National Cancer Institute: <http://dietassessmentprimer.cancer.gov/>. Medical Research Council, Glossary of terms: <http://dapa-toolkit.mrc.ac.uk/dietary-assessment/da-glossary-of-terms.php>. (Both accessed 23 October 2015.)

E

Energy adjustment is an analytic method by which nutrient or food quantity intake is corrected for the total energy intake.

F

Fisher's exact test is a statistical test used to determine if there are non-random associations between two categorical variables.

It is commonly used when the sample size is small. Fisher's exact test is more accurate than the chi-square test of independence when the expected numbers are small.

Food consumption in the present resource-guide refers to an estimate of the quantity and/or variety of a food or group of foods consumed by an individual, household or a specific population.

Food composition table provides detailed information on the nutrient values of foods – energy, macronutrients (energy, protein, carbohydrates) and micronutrients (vitamins and minerals). Nutrient values are usually expressed in terms of the nutrient content of the edible portion of the food per 100g.

G

Goldberg cut-off is used to identify dietary underreporting and is based on the principle that an individual of a given age, sex and body weight requires a minimum energy intake. The cut-offs can be used at both the individual and group levels.

M

Measurement error is the difference between the true value of a parameter and the value obtained from reporting e.g. dietary intake.

Misreporting in self-report dietary assessment is considered to be unavoidable and can include both under and over reporting.

P

Physical activity level (PAL) is a way to express a person's daily physical activity as a number, and is used to estimate a person's total energy expenditure. In combination with the basal metabolic rate, it can be used to compute the amount of food energy a person needs to consume in order to maintain a particular lifestyle.

R

Random error is a type of measurement error that contributes to variability (reduces precision) but does not influence the sample mean or median. It generates a deviation from the correct results due to chance alone.

Regression is a statistical measure that attempts to determine the strength of the relationship between one dependent variable (usually denoted by Y) and a series of other changing variables known as independent variables (usually denoted by X).

Reliability is a quality of the measurements relating to a technical aspect of measurement noting the ability to accurately measure the real change or achievement in a consistent and comparable manner over time and space.

Reproducibility assesses the degree to which a method provides similar results when used repeatedly (on two or more occasions) in the exact same situation.

S

Sensitivity in dietary assessment (also called the true positive rate) measures the proportion of positives that are correctly identified as such (e.g. the percentage of sick people who are correctly identified as having the condition).

Specificity in dietary assessment (also called the true negative rate) measures the proportion of negatives that are correctly identified as such (e.g. the percentage of healthy people who are correctly identified as not having the condition).

Statistical significance refers to the likelihood that a difference or relationship exists or if it is caused by a mere random chance. Statistical hypothesis tests (e.g. Chi Square, t-test, ANOVA) are traditionally employed to determine if a result is statistically significant or not. Most authors refer to statistically significant as $P < 0.05$ and statistically highly significant as $P < 0.001$ (less than one in a thousand chance of being wrong).

Systematic error (also known as bias) is a type of measurement error in which measurements consistently depart from the true value, in the same direction. Systematic error affects the sample mean and can result in incorrect estimates and conclusions.

T

Total energy expenditure refers to the energy spent, on average, in a 24-hour period by an individual or a group of individuals. Total energy expenditure consists of three components: Basal Metabolic Rate (typically 60–75 percent of total energy expenditure), the thermic effect of food (10 percent), and energy expenditure due to physical activity (15–30 percent).

True intake is the actual intake, which usually cannot be measured among free-living individuals.

T-test is a statistical analysis to test the difference of two populations means that are normally distributed. Commonly applied with small sample sizes, testing the difference between the samples when the variances of two normal distributions are not known.

U

Usual intake is the long-term average daily intake, taking into account both consumption days and non-consumption days.

V

Validity assesses the accuracy of self-report instruments in measuring true intakes.

W

Wilcoxon Signed Rank test is a nonparametric test that compares two paired groups. The test essentially calculates the difference between each set of pairs and analyses these differences. It can be used as an alternative to the t-test when the population data does not follow a normal distribution.

Within-person variation (also known as day-to-day variation) is the difference between assessing a variable or variables collected via a single administration of an instrument, compared with a long-term average based on multiple administrations of the instrument.

EXECUTIVE SUMMARY

The present resource guide provides a comprehensible insight into dietary assessment, and into the challenges and considerations linked to the selection of the most appropriate method. The guide has been developed to provide assistance in the collection of dietary information, to be used to inform a number of programmatic decisions, as well as policy formulation, and to address diet-disease relations. The guide first provides a conceptual background of different dietary assessment methods, highlighting both indirect and direct (prospective and retrospective) methods, and providing a description of their application, validity, strengths and limitations. The guide also provides tips and methodological considerations to take into account during method selection and implementation, along with examples of forms and questionnaires used in previous studies. Lastly, the guide addresses technical and financial considerations, and looks at key factors to be taken into account prior to the selection of a direct dietary assessment method, such as the importance of identifying the study objective and selecting the appropriate study design. This is followed by a step-by-step guide to facilitate the selection of a dietary method along with a summary of the major features of direct methods. Throughout the guide, a special effort is made to include evidence from low resource settings when describing the accuracy, reproducibility, validity and applications of the methods.

The purpose of the resource guide is to facilitate and improve the quality and accuracy of nutrition information collected. The need for this improvement is reflected in international calls for the incorporation of robust nutrition data into national information systems (e.g. ICN2). Selecting the most appropriate dietary assessment method for a given purpose will in turn help generate better evidence for formulating effective nutrition projects, policies and programmes. This resource guide is written for professionals who play a role in the selection of the dietary assessment method for use in regional or national dietary and nutrition surveys, programmes and monitoring frameworks. These professionals may be programme managers, educators, health care professionals (including dietitians, nutritionists and health promotion specialists), students and extension workers. The information presented in the resource guide is intended to be used to direct and help steer the decision on the selection of the most adequate dietary assessment method according to the study objectives, population's characteristics and available resources, and should not be used as a tool to provide all the answers for the selection process. References for further reading have been included to supplement the guide and provide more advanced information for those who would like to go beyond the scope of this publication.



©FAO/Ruth Charrondiere

1

INTRODUCTION

Strengthening and sustaining the capacity of countries to incorporate robust nutrition indicators into their information systems would help to generate better evidence for formulating effective agricultural and nutrition policies. The need for such robust information was recently reaffirmed at the Second International Conference on Nutrition (ICN2), jointly organized by the Food and Agriculture Organization of the United Nations (FAO) and the World Health Organization (WHO) in November 2014. More specifically, as stated in one of the outcome documents from ICN2 (The Rome Declaration):

“Nutrition data and indicators, as well as the capacity of, and support to all countries, especially developing countries, for data collection and analysis, need to be improved in order to contribute to more effective nutrition surveillance, policy-making and accountability².”

The Nutrition Assessment and Scientific Advice Group (ESNA), a branch of the Nutrition and Food Systems Division of FAO, plays an active role in providing technical support to countries in collecting food and dietary information. This support is often provided within a framework

of generating evidence-based policies, implementing ICN2-related follow-up activities, and ensuring government accountability. In order to produce better evidence for formulating effective nutrition projects, policies and programmes, the most appropriate dietary assessment method first needs to be selected. This resource guide has been developed with the purpose of providing a comprehensible review on dietary assessment methods based on the latest research and development and the challenges and considerations that are linked to the selection of the most appropriate dietary assessment method, specifically focusing on low resource setting areas.

Ultimately, the guide can be used as a resource to:

- Strengthen the decision-making process for professionals when used as part of an informed process of selecting the most appropriate dietary assessment method for their particular study, and to provide a resource for those who want a review on the topic.
- Gain a better understanding on the specific challenges and needs that professionals face in

² Paragraph 13g, Rome Declaration on Nutrition, www.fao.org/3/a-ml542e.pdf (Accessed 23 October 2015)

low resource setting areas when they need to assess the diet of individuals and populations, such as the availability of food composition tables, estimation of portion sizes, seasonality, and the characteristics of specific populations and geographical locations.

- Complement other sources of information – such as information generated from needs assessment exercises, scientific expertise and local knowledge and experience – that influence the selection of the final method for measuring food and nutrient intakes for a given purpose.
- Direct users to ask appropriate questions that will lead to the selection of a method that properly considers data needs and the purpose of the study. In so doing, users will need to understand that compromises and more resourceful approaches are needed, especially when working in low resource settings, in order to select methods that take into account, resources required and resources actually available, culturally specific issues, time and human resources available for data collection and analysis, issues pertaining to portion size estimation and the availability of food composition tables.

The guide addresses the fundamental aspects involved in the selection of a dietary assessment method and data collection process. It also highlights the selection of methods for specific programmatic needs by providing a number of examples. It therefore aims to support, not dictate, the selection of a dietary assessment method, by being part of the informed decision-making process that results in a well-thought-out selection. The key objectives of the resource guide are:

- to provide users with practical guidance on available dietary assessment methods and

to enhance users' understanding of their key features, strengths and limitations;

- to describe the main methodological considerations involved in dietary assessment, specifically in low resource settings³;
- to outline and elaborate on the main sources of measurement errors and bias, and to explain why they occur;
- to explain and demonstrate – with a specific focus on low resource settings – the possible consequences of overlooking measurement and methodological considerations during data collection, analysis and interpretation, and their impact on overall data quality.

³ This refers to settings with limited capacity and resources to perform nutritional assessment.



©FAO/Warren Lee

2

DIETARY ASSESSMENT METHODS

Dietary assessment is an evaluation of food and nutrient intake and dietary pattern of an individual or individuals in the household or population group over time. It is one of the four approaches in nutrition assessment to evaluating the nutritional status of individuals comprehensively. The other three are anthropometrics, biochemical parameters and clinical examination (Gibson, 2005). Dietary assessment methods are usually categorized according to the nature of the method used as shown in Figure 1. Indirect methods utilize secondary data for assessing diets, while direct methods collect primary dietary data from individuals. Prospective and retrospective methods refer to the time food consumption is recorded. Prospective methods involve recording the diet when the foods are being consumed; retrospective methods are based on a recall of food intake that have already been consumed. Furthermore, dietary assessment can be qualitative (types of food consumed) or quantitative (types and amounts of food consumed). The latter allows estimation of an individual's food, energy and nutrient intakes. Selection of an appropriate method for dietary assessment depends on the purposes of the study, which may be to measure


food consumption, nutrient intake or eating habits.

2.1 INDIRECT METHODS

Indirect methods use secondary information (e.g. food supply, agricultural statistics, food expenditure) to estimate food available for consumption at the national and household levels. Firstly, we will look at the Food Balance Sheet (FBS)⁴, which provides food consumption information at national level: food consumption is estimated from the point of view of food supply. Secondly, we will discuss about the Household Consumption and Expenditure Survey (HCES), which provide food consumption information *at household level*: food consumption is estimated from the point of view of food demand. Neither of the two methods directly obtain primary dietary data from individuals to evaluate dietary intake or food consumption on an individual basis. Indirect methods are useful for identifying trends in food availability and consumption across different geographical regions and time.

⁴ Others terms that relate to FBSs include: "national food accounts"; "supply/utilization accounts"; "food disappearance data"; and "food consumption level estimates". These terms reflect different methods of calculating food availability.

Figure 1 - Overview of dietary assessment methods to estimate food and nutrient consumption at national, household and individual level



2.1.1 FOOD BALANCE SHEETS – NATIONAL FOOD AVAILABILITY

FBS are compiled by FAO annually and are mainly based on country-level data covering a calendar year. They are used to assess national food availability for consumption. Using these data and the available information on seeds, waste coefficients, stock changes and types of utilization, a supply/utilization account is prepared for each commodity in weight terms.

Besides commodity-by-commodity information, the FAO FBS⁵ also provide total food availability estimates by aggregating the food component of all commodities including fishery products (FAO, 2001; FAO, 2015). The data are conventionally presented per capita, by adjusting for population size, e.g. available amount of food/energy/nutrients per person per day. More specifically, FBS present a comprehensive picture of a country's food supply over time. However, FBS provide little information on the variability of food

⁵ <http://www.fao.org/faostat/en/#home>

intake of individuals within a population. The FBS list food items, i.e. primary food commodities and a number of processed food commodities potentially available for human consumption, including their sources of supply and utilization.

Online FBS data are compiled by FAO annually for approximately 185 countries and for a total of around 100 food commodity groups worldwide that may potentially be available for human consumption in the country. Gross national food supply in a given reference period of a country is calculated from the total quantity of food produced plus the total quantity imported, adjusted for changes at national food stock levels and exports. Net food availability is calculated by subtracting the amounts used for animal feed, seeds, industrial or other purposes and losses in the supply chain. This net value is then divided by the country's population estimate to obtain a final figure describing the availability of food commodities, expressed as kilograms per capita per year. This per capita information can also be linked to food composition data and presented as per capita energy intake (kilocalories per day), protein intake (grams per day) and fat intake (grams per day).

2.1.1.1 Applications and uses of FBS

The FAO FBS are widely used in the food and agriculture sectors to monitor global food patterns and dietary habits. This monitoring also includes trends and changes in overall national food availability, and the adequacy of a country's supply to meet nutritional requirements. Currently, the FAO FBS only provide data on annual per capita consumption of energy, protein and fat, while they do not provide data on micronutrients. FBS have also been used to set public health priorities, formulate policies, undertake intercountry comparisons, and estimate the likelihood of micronutrient deficiencies. For example, FBS data was used to identify the probability of micronutrient deficits in food supply per capita

for 17 countries in the Western Pacific (Gibson *et al.*, 2012). Additionally, FBS were employed to estimate the global prevalence of inadequate zinc intake (Wessells *et al.*, 2012), and to examine the impact of improved nutrient supply on meeting a population's micronutrient needs etc. (Arsenault *et al.*, 2015). Furthermore, (Naska *et al.*, 2009) national FBS data has been used to examine the correlation with mortality statistics. More recently, FBS have been utilized in formative research to examine the nutrition transition (Mattei *et al.*, 2015). Other studies have employed FBS data to analyse variation in adherence to the Mediterranean diet between 1961–1965 and 2000–2003 (da Silva *et al.*, 2009). However, the actual distribution of food consumed among individuals in the population of a country as categorized by socio-economic status, age, or gender cannot be determined by using FBS.

2.1.1.2 Accuracy of FBS

The accuracy of FBS relies on the underlying accuracy and reliability of the statistics that the FBS are based on. These statistics are mostly derived from the official primary commodity production data and the primary and derived commodities trade data. Some adjustments may be required before the data can be used by FBSs. The extent to which the basic data have properly reflected the reality needs to be cross-checked with factors such as food losses and waste and unrecorded trades across national boundaries, etc. Literature on the accuracy of FBS estimates are scarce, and the available literature often focuses on the differences in the trends of food supply and availability over time. Serra-Majem *et al.* (2003) undertook a comparative analysis that evaluated three types of nutrition surveys: the FBS, the Household Budget Survey (HBS) and the individual dietary survey. They analysed inter- and intra-country comparisons of data for different stages of the food chain in Canada, Finland, Poland and Spain. It was concluded that FBS overestimated the energy, alcohol and fat intake of

individuals, and the percentage of energy derived from fat (except in Poland), when compared with individual dietary surveys. FBS estimates were found to have exceeded those from the nationally representative dietary survey data collected in the Global Dietary Database (GDD) for most food groups, namely fruit, vegetables, whole grains, red and processed meat, fish and seafood, and milk, as well as total energy intake, while beans, legumes, nuts and seeds were underestimated. The differences were significant ($P < 0.05$) ranging from 54 percent for total energy intake to 270

percent for whole grain intake (Del Gobbo *et al.*, 2015). In low resource countries, the reliability of FBS data may be further limited by the quality and representativeness of the national primary statistics, and under-reporting of food available through home grown food, hunting and gathering, non-commercial production, etc.

2.1.1.3 Strengths and limitations of FBS

The following table will provide a summary of the strengths and limitations associated with conducting an FBS.

Table 1 - Strengths and limitations of using FBS data for assessing diets

STRENGTHS	LIMITATIONS
<ul style="list-style-type: none"> • Inexpensive source of indirect nutrition data, standardized, accessible by all, relatively simple to analyse • Include nearly all countries worldwide • Monitor global nutrition patterns and dietary habits, including trends and changes in overall national food availability 	<ul style="list-style-type: none"> • Cannot provide disaggregated information across different population characteristics, e.g. socio-economic, demographic etc. • Estimates are derived from primary and/or basic country statistics which themselves may be subject to methodological errors • Do not provide data on seasonal variations in the food supply • Do not provide data on foods not included in national production statistics such as game, wild animals and insects, and non-commercial or subsistence production • Do not provide individual-based dietary estimates • Statistics may be subject to incomplete or unreliable estimates of the total population in a given country • Food waste (domestic and retail), processed foods, home grown food production and food from non-retail sources are not accounted for • Time lag between data collection and availability on FAOSTAT (FAO Statistics Division)

2.1.2 HOUSEHOLD CONSUMPTION AND EXPENDITURE SURVEYS – HOUSEHOLD FOOD CONSUMPTION

Household food consumption has been defined as “the total amount of food available for consumption in the household, generally excluding food eaten away from the home unless taken from home” (Putnam *et al.*, 1994). There is a wide range of multipurpose household surveys, such as the Household Budget Survey (HBS), the Living Costs and Food Survey (LCFS), the Household Income and Expenditure Survey (HIES), the Living Standards Measurement Study (LSMS), the Household Expenditure Survey (HES) and the Integrated Household Survey (IHS) – that measure food consumption or its proxies, are collectively known as HCESs. The central statistical offices in countries are usually responsible for data collection. Household members keep records on all expenses and type of foods consumed during a specific time period, usually one to four weeks, and preferably evenly distributed during different times in the year, which is then provided to enumerators. The collected data are analysed and used to assess food consumption at household level. Surveys of this type are routinely undertaken in many countries to provide information for the calculation of consumer price indices, to study household living conditions and analyse trends in poverty and income distribution (Moltedo *et al.*, 2014). In some low resource settings, information generated from these surveys is the only form of food consumption data that can also be used to calculate estimates of nutrient intake. These estimates are calculated by multiplying the average food consumption data by the corresponding nutrient values for

the edible portion of the food. Data on nutrient values are obtained from food composition tables (Gibson, 2005). However, household surveys do not provide information on the distribution of food consumption between family members, cooking methods or food losses. These surveys are often performed for economic reasons rather than for nutrition or health reasons.

Fiedler *et al.* (2012) provide a comprehensive review on the availability and characteristics of HCES, indicating that currently there are more than 700 published surveys. These surveys differ in terms of the nature and level of detail by country, and cover over one million households in 116 low- and middle-income countries⁶. The key findings of the review are as follows:

- The designs of the surveys were not harmonized or standardized, meaning that they differed in key characteristics such as questions used in the questionnaire, coverage, frequency, sample size and statistical accuracy⁷.
- Routine information was collected including household composition, housing characteristics, income, assets, wealth and livelihood, as well as personal information such as age, sex and education. Information on food consumption was mainly collected via recalls and occasionally via diaries. The latter was reported to be a more accurate but also a more expensive method of data collection.
- The list of food items in HCES questionnaires varied in length and composition.
- All surveys asked whether a food item was purchased, homemade, received as a form of ‘salary’, or received from friends, relatives or a social programme.

⁶ See also <http://blog.usaid.gov/2014/03/the-power-of-household-consumption-and-expenditure-surveys-hces-to-inform-evidence-based-nutrition-interventions-and-policies/> (Accessed 1 December 2016)

⁷ The International Household Survey Network (IHSN) provides a number of resources on HCES. It aims to improve the availability, accessibility, and quality of survey data within developing countries and to encourage the analysis and use of this data by national and international development decision-makers, the research community and other stakeholders. See: <http://www.ihsn.org/home/> (Accessed 23 October 2015)

- Some surveys captured both food quantities and costs, while others captured costs only.
- A majority of the surveys overlapped data on food acquisition and consumption, which can result in overestimating consumption.
- The surveys presented clarifications regarding food stocks, as well as information on the quality of the food composition tables (in some cases food composition information was absent, outdated or available only for a limited number of foods).

2.1.2.1 Applications and uses of HCES

HCES are recognized as an inexpensive and more readily available alternative for tracking food consumption patterns when compared to individual-based methods. HCES have been used by FAO for global monitoring of food security, e.g. for Target 1c of the Millennium Development Goal 1 (“Halve, between 1990 and 2015, the proportion of people who suffer from hunger”) and the similar goal set in 1996 by the World Food Summit. Information on food consumption at the household level allows the derivation of variability parameters such as the coefficient of variation of food consumption which are used to estimate undernourishment (FAO/IFAD/WFP, 2015). Estimation of undernourishment is often conducted in partnership with national and regional institutions, in conjunction with capacity development activities.

To support the use of household surveys, FAO, in collaboration with the World Bank, developed the Food Security module of the ADePT software (ADePT-FSM)⁸. The software allows streamlined and consistent food security statistics to be estimated using HCES data (Molledo *et al.*, 2014). Food security indicators are derived at national

and subnational levels, by population groups, by food commodity groups and by food items, using standardized files as inputs. Examples of such indicators include inequality levels in calorie consumption, as well as consumption levels of macronutrients, micronutrients and amino acids. The wider acceptance of data gathered from HCES received a further boost via the formulation, implementation and evaluation of nutritional policies across Europe with the development of the Data Food Networking⁹ (DAFNE) initiative. The DAFNE databank is based on information collected as part of the HCES which are periodically conducted in various EU countries. A significant achievement of this project is the development of a common classification system for food variables and socio-demographic variables in national HCES, allowing inter-country comparisons. Elsewhere, HCES¹⁰ data have been used to assess household dietary intakes in relation to nutrition transition in Cape Verde (Dop *et al.*, 2012), and to estimate food consumption and micronutrient intakes (vitamin A, iron, zinc) in Bangladesh, thereby identifying population subgroups at risk of inadequate micronutrient intakes (Bermudez *et al.*, 2012).

2.1.2.2 Accuracy of HCES

Smith *et al.* (2007) examined the reliability and relevance of HCES data, including 100 household surveys from low and middle income countries. They concluded that nearly all surveys were appropriate when the aim was to measure the percentage of households purchasing and consuming individual foods. However, for estimating the quantities of individual foods consumed and assessing micronutrient insufficiencies it was shown that this data was useful in less than 10 percent of surveys, mainly

⁸ <http://www.fao.org/economic/ess/ess-fs/fs-methods/adept-fsn/en/> (Accessed 23 October 2015)

⁹ <http://www.nut.uoa.gr/dafneENG.html> (Accessed 23 October 2015)

¹⁰ Smith and Subandoro (2007) have produced a detailed guide for practitioners in assessing food security status in the population using HES.

because of issues related to units of measurement and the estimation of food consumed away from home. The authors appraised the reliability of the surveys in terms of how they addressed different areas of investigation: the recall period for home-bound food data collection; how food acquisition was analysed; completeness of enumeration; comprehensiveness and specificity of the home-bound food list; the quality of data collected on food consumed away from home and how seasonality in food consumption was accounted for. Recommendations to improve reliability focused on three criteria which were not met by approximately half of the surveys, i.e. seasonality, out-of-home eating and specificity of survey food lists. In Uganda, Jariseta *et al.* (2012) compared estimates of nutrient densities in the diet of women and children by HCES and by a 24-hour recall. Nutrient densities were calculated as the nutrient contents per 2 000 kcal of edible portion of food consumed (nutrient content divided by energy intake). The authors found no significant differences between the medians of

energy intake in 7 out of the 14 nutrient densities (i.e. protein, fat, fibre, iron, thiamin, riboflavin, and vitamin B6) estimated by the HCES and 24-hour recall ($P < 0.05$). They concluded that HCES estimates were close proxies for 24-hour recall measures of nutrient density. Whereas HCES may be less precise than individual dietary assessment methods, the relative low costs have made HCES an attractive tool for decision-makers to inform national policies and identify areas where nutritional interventions are needed.

2.1.2.3 Strengths and limitations of HCESs

The following table will provide a summary of the strengths and limitations associated with conducting a HCES.

Table 2 - Strengths and limitations of using HCES data for assessing diets

STRENGTHS	LIMITATIONS
<ul style="list-style-type: none"> ● Inexpensive source of nutrition data since they are also collected for purposes other than nutrition, standardized, accessible to all and relatively simple to analyse, if the design of the questionnaire is appropriate ● Routinely conducted in a number of countries on sample populations that are nationally representative of the national demographic ● Collect information on socio- economic and demographic characteristics of the head of household. This type of survey also enables investigation of subnational variations in consumption patterns, which can be invaluable in designing nutrition programmes ● Statistically representative at the national level, and usually also at the subnational level 	<ul style="list-style-type: none"> ● Limits the ability of the respondent to report completely (via recall or record) all foods consumed by the household. Good training of enumerators and careful questionnaire design could help to alleviate this limitation ● Units used to report food quantity are non-standardized ● Food wasted or food given away is not accounted for: consumption may be overestimated ● Food eaten away from home is not always accounted for: consumption may be underestimated

STRENGTHS	LIMITATIONS
<ul style="list-style-type: none"> Comprehensive, as they contain detailed household food consumption information that allows direct observation of the agriculture and nutrition nexus, through markets, value chains, and other pathways Information collected enables examination of trends and changes in overall national food availability and consumption 	<ul style="list-style-type: none"> Leads to misrecording and/or misreporting of important foods when using a predetermined list of food items The size of the household may be different from the number of people who actually consumed the food over the reference period Does not collect information on individual food consumption or on the distribution of foods among household members In the case of acquisitions surveys, food that is stocked up for an extended period of time remains unaccounted for Nutrient estimates are dependent on the quality of the food composition table.

2.2 DIRECT METHODS

Direct methods using individual-based dietary assessment can be classified into two groups, as described in Figure 1. Retrospective methods measure food intake from the past. These methods include 24-hour recall, food frequency questionnaires (FFQ), and dietary history. Prospective methods assess current food intake. These methods include food records and the duplicate meal method. Estimates obtained from direct methods are used to identify trends in food consumption, food and nutrient intakes, eating patterns, and to evaluate diet–disease associations. Information provided by such methods can also be used to calculate relevant food-based indicators for monitoring and evaluation purposes. This section will describe dietary assessment methods used in individual-based surveys, and analyse their application, focusing mainly on low resource settings, and will also cover the issues of precision and accuracy.

2.2.1 RETROSPECTIVE DIRECT METHODS

Retrospective methods collect information on foods and beverages already consumed. Among the retrospective methods are the FFQ, 24-hour recalls and diet histories. These methods greatly depend on the memory of the respondent and their ability to recall all foods and portion sizes consumed over a reference period of time. Quantities of foods consumed can be obtained by food models, pictures of foods, standard household measuring cups, spoons, etc.

2.2.1.1 Food frequency questionnaire

FFQs assesses the frequency with which foods and/or food groups are eaten over a certain time period. The questionnaire includes a food list (usually close-ended) and a frequency category section, and can be self- or interviewer-administered. Appendix 1 gives an example questionnaire. Depending on the study objectives, data collection might be daily, weekly, monthly or yearly. Furthermore, FFQs can include information

about portion sizes and/or quantity of food intake. These types of questionnaires are known as semi-quantitative, i.e. general portion sizes are described and quantified, or specific portion sizes are recalled and supported by food pictures for each item on the food list. By including portion size as part of frequency, the questionnaire allows for the estimation of food quantities eaten and of nutrient intakes. FFQs can either be developed from basic principles or adapted from existing questionnaires (Cade *et al.*, 2002). In the first case, important decisions and considerations are needed in developing the food list. Several key considerations include:

- Foods selected should encapsulate the objectives of the assessment, e.g. to measure intake of only a few foods and nutrients, or to undertake a comprehensive dietary assessment (Willett *et al.*, 2013).
- Whether to rank individuals' consumption or provide a measure of absolute nutrient intakes.
- It is often preferred to put together a comprehensive list of foods and/or of food groups to allow for energy adjustments. Aggregating foods into food groups can be used as a technique to capture specific nutrient(s) or non-nutrient(s) when these nutrients are confined to a relatively small number of foods (Gibson, 2005). However, aggregation of foods into small groups may lead to underestimation of intakes, whereas larger food groupings can lead to overestimation of intakes. Aggregating food can further lead to over counting due to difficulties in reporting combined frequency for a particular food eaten both alone and/or in mixed dishes (Cade *et al.*, 2002).
- The choice of foods in a list is partly data driven and partly a question of scientific judgment (Patterson *et al.*, 2004). Selected foods can be used to capture the major sources of energy and/or nutrients consumed by the study population, variability in food intake between persons, and of course the study objectives.
- The list of foods in the FFQ cannot be infinite as it could potentially increase the burden on the respondent. Therefore an *a priori* decision should be taken on the foods to be included, their frequency of consumption in the studied population, their cultural importance and their relative value as a specific nutrient source.

Methods for selecting food items to be incorporated into a FFQ food list may range from information obtained from previous dietary assessment reports to focus group discussions and pilot 24-hour recalls.

- A review of the literature can be used to help select the appropriate food items to include in the FFQ. This approach consists of a detailed revision of available dietary data of the target population. Data on the past dietary surveys, cultural beliefs and food choices are reviewed in order to select and confirm foods and food categories to be included in the FFQ. In a cross-sectional study designed to capture diet patterns in women in Micronesia, ethnographic data was used to develop a structured 7-day FFQ (Corsi *et al.*, 2008).
- Focus groups should consist of individuals selected randomly from the target population and be representative of the population. Individuals are asked to identify through discussions the commonly-eaten foods in the area, along with information on ingredients used, food preparation methods, the seasonal-variability of foods and culturally-specific dishes. Later on, the groups can hold an open discussion and agree on the relevant food list to construct for the FFQ questionnaire. For example, 19 focus groups were used to generate data for the FFQ development in Botswana (Jackson *et al.*, 2013). Focus group discussions were organized by home economists together with agricultural demonstrators in each region.
- 24-hour recall (for more information on 24-hour recall, see section 2.2.1.2) can be used as a tool for selecting the appropriate food

items to be included in the FFQ. This approach was used in a study conducted in Colombia, where a random subsample of 100 individuals representative of the target population were asked to record their food intake using a single 24-hour recall. The FFQ was then developed based on the most frequently reported food items, excluding foods that had a low frequency of consumption (Dehghan *et al.*, 2012).

Questionnaires can be modified versions of existing one. However, caution should be employed in assessing the original purpose and validity of the parent FFQ: for example, for whom it was written, when it was developed, whether it had been previously validated etc. (Cade *et al.*, 2002).

2.2.1.1.1 Applications and uses of FFQs

There is a plethora of FFQs available, and they continue to be developed or adapted for different purposes. FFQs are commonly used in large epidemiological studies (Willett *et al.*, 2013) to capture data on dietary intakes and patterns (Corsi *et al.*, 2008; Merchant *et al.*, 2005), to assess diet–disease associations (Liu *et al.*, 2001; McCullough *et al.*, 2002) and to calculate correlations or relative risks (Hutanasu *et al.*, 2009). FFQs can also be used to assess seasonal dietary patterns (Campbell *et al.*, 2014). However, there is an ongoing debate on the use of FFQs in assessing diet, with some authors questioning the validity of results obtained from studies where FFQs were applied (Kristal *et al.*, 2005). Others argue that the value of FFQs in epidemiological applications has been documented objectively by correlations with biochemical indicators (Willett *et al.*, 2007). The FFQ is a time-effective method that is easy to administer and provides a simple data entry procedure. However, FFQs have limited sensitivity to changes in food supply and are not suitable for people with a wide variation in dietary intake patterns (Gibson, 2005). As an example, a semi-quantitative FFQ was used to investigate

intakes of major carotenoids and tocopherols in Costa Rican adolescents. The authors suggested that when assessing the diet of adolescents it is important to ask about specific portion sizes, and to support the questions with food pictures or household measures (Irwig *et al.*, 2002).

FFQs have also been used to measure year-round and seasonal dietary patterns in a large sample of rural Nepalese women (n=15 899) (Campbell *et al.*, 2014). The respondents were asked to report on the frequency of foods consumed over the previous year, using day, week and month as the unit of measure for the frequency of intake. The study revealed that intakes of vegetables, fruits and animal-source foods were infrequent in certain seasons and especially among poorer members of the rural population. The authors noted that the study design could have been improved by including local farmers and resident focus groups for information on seasonal foods. In a study in Micronesia, where an FFQ was used to capture diet patterns in adult women, the authors highlighted the importance of training interviewers on accurate data collection and ensuring a good understanding of local culture and customs (Corsi *et al.*, 2008).

2.2.1.1.2 Reproducibility and validity of FFQs

Methods used to validate FFQs include multiple 24-hour recalls, food records (both weighed and estimated) and biomarkers. Special attention is required to ensure that measurement days captured by the records or recalls reflect the time frame covered by the FFQ, since some FFQ time frames can cover up to one year (Willett *et al.*, 2013). Though a weighed food record is the preferred validation method, it is not error free. A weighed food record is more valid and precise than 24-hour recall to validate FFQ. (see section 3.7 & 3.8: 'reproducibility and validity in dietary assessment'). However, when the study participants are illiterate, the use of multiple 24-hour recalls is more appropriate than a weighed

food record, despite the weaker correlations because both FFQ and 24-hour recall rely on memory and estimation of portion size. Listed below are a number of factors related to FFQ development that could influence validity (Cade *et al.*, 2002):

- The number of food items in the list. This is partly determined by the population characteristics and study objectives.
- The order of the food list, e.g. items of particular interest should be placed at the beginning of the questionnaire.
- The frequency and portion size responses. These should be close-ended rather than open-ended, to minimize errors in coding and transcription.
- The time frame of the recall period.
- The mode of administration.
- Data entry and computation once the survey is complete.

The reproducibility and validity of a 124-item FFQ in assessing habitual nutrient intake of Malay adolescents were validated against a non-consecutive three day 24-hour recall (Nurul-Fadhilah *et al.*, 2012). For reproducibility, the correlation coefficients for energy-adjusted nutrient intakes between the two FFQ administrations ranged from 0.43 (carotene) to 0.86 (fat), indicating a good reproducibility. For validity, correlation coefficients between energy-adjusted nutrient intakes between the methods ranged from 0.22 (zinc) to 0.68 (calcium), indicating a moderate to good agreement between the two assessment methods. In both cases the correlations were weaker for micronutrients than for macronutrients.

In a study conducted in Botswana, Jackson *et al.* (2013) tested the reproducibility and validity of a quantitative 122-item FFQ – repeated after one year – by comparing nutrient and food group intakes against four non-consecutive 24-hour

recalls, covering a time frame of one year. For reproducibility, correlation coefficients for energy-adjusted nutrients ranged from 0.39 (retinol) to 0.66 (vitamin E), indicating good reproducibility. Good validity in estimating most food and nutrient intakes was observed, except for iron, retinol, β -carotene and related food groups (i.e. fruits, dark green leafy vegetables and yellow vegetables). The authors suggested that this could be attributed to the seasonal availability of fruits and vegetables leading to substantial fluctuations in estimated intakes (Jackson *et al.*, 2013).

Kabagambe *et al.* (2001) assessed the reproducibility and validity of a 135-item FFQ in Costa Rican adults using seven 24-hour recalls, plasma samples (for assessment of tocopherol and carotenoid) and adipose tissue samples (for assessment of tocopherol, carotenoid and fatty acid intake). For the reproducibility study – where the administration of the FFQs was 12 months apart – correlation coefficients ranged from 0.33 to 0.77, indicating good to high reproducibility. In the validity analysis, the FFQ gave higher values of energy and nutrient intakes compared with those obtained by the recalls for all nutrients studied, except for vitamin K, iron and caffeine. The authors concluded that FFQ was a valid and reproducible tool to measure the diet of adults. They suggested that biomarkers should be used to complement the FFQ rather than substitute for it, as biomarkers did not perform better than the FFQ in this study. The study furthermore, showed that biomarkers did not give a better result than the FFQ (Kabagambe *et al.*, 2001).

In another study, the reproducibility and validity of a self-administered 130-item FFQ was assessed against biomarkers in urine (nitrogen, potassium, and sodium) and blood (plasma ascorbic acid), and compared to a 7-day food record. For validity, the correlation between urinary potassium and dietary potassium from the FFQ was 0.33 and from the food record was 0.53. The authors concluded that food record (see sections 2.2.2.1 and 2.2.2.2 for

additional information on food records) provides a better estimates of nutrient intakes than the FFQ (McKeown *et al.*, 2001).

2.2.1.1.3 Strengths and limitations of FFQ

The following table will provide a summary of the strengths and limitations associated with conducting an FFQ.

Table 3 - Strengths and limitations of FFQ

STRENGTHS	LIMITATIONS
<ul style="list-style-type: none"> ● Assess the usual intake over a long period of time. ● Can be used to capture a range of foods, specific nutrient(s) (quantitative FFQ) or a specific food group, including rarely consumed food items. ● Can capture portion size estimates (semi and quantitative FFQ), details about cooking and preparation methods. ● An open section added to the end of the questionnaire can allow for addition of foods consumed that are not present in the food list. ● Since it is a retrospective method, it does not affect eating behaviour. ● Low respondent burden. ● Relatively simple to administer and inexpensive when compared with other assessment methods (i.e. 24-hours recall, dietary records). ● Interview-based FFQ questionnaire does not rely on the literacy and numeracy skills of the respondent. ● Can also be self-administered via mail or internet. ● Appropriate for large studies as it can be administered using a machine-scannable format, reducing data-entry errors. 	<ul style="list-style-type: none"> ● The food list cannot cover all the foods consumed by the respondent, which may lead to underreporting. ● Does not give precise information on the estimated portion size consumed. ● Requires respondents who are literate and have numeracy skills (if self-reported). ● Self-administered FFQs can lead to misinterpretation of questions and the omission of food items that are not understood by the respondent. ● Questionnaires need to be adapted and validated to reflect the study population and purpose. This may require a considerable amount of time and resources. ● Not suitable for a population where people have distinctly different dietary patterns. ● Relies heavily on memory; therefore, declined cognitive ability can result in errors when reporting on frequency and portion size estimation. ● Misreporting arises when reporting combined frequencies for a particular food eaten both alone and in mixed dishes.

BOX 1. QUICK GUIDE TO USING AN FFQ

Project objectives and budget determine the study design and sample size

- ✓ Who is the population of interest?
- ✓ Do you want to collect information on specific nutrient(s), foods, food groups or the overall diets?
- ✓ What is the objective of the data collection, e.g. to rank respondents from low to high intakes, or to provide a measure of estimated intake?

Define the target population (elderly, children, adolescents etc.), literacy level, and cognitive ability

Frequency categories in the FFQ: times per day, times per week, times per month, rarely, never etc.

Developing a survey protocol

- ✓ Adapted to local cultural context, e.g. meal patterns, shared dishes, non-standard eating and serving tools.
- ✓ A validation exercise can be applied to test the efficiency of the protocol.

Identifying sources of information

- ✓ Information on foods consumed by a population can be obtained from national or regional survey data, databases, or from undertaking a focus group discussion with the target population.

Development of a food list and assigning food codes

- ✓ Foods selected should represent those commonly consumed by the target population and the food list should be in line with the study objectives.

Adapting an existing food list

- ✓ If a similar FFQ already exists, it can be used in its original form or modified/ adapted by adding or replacing foods with items more commonly consumed in the target population, or by adapting the food list to target a specific nutrient. However, changes to an FFQ will require a validation study.
- ✓ Update the database as required to include all the food components of interest.

Assessing the need for portion size estimation (non-quantitative, semi-quantitative or quantitative)

- ✓ Determining if the FFQ should collect quantitative information on food intake would depend on the objectives of the study, age of respondents, homogeneity of the target population, standard units available, and the type of information to be collected.

Estimation of portion size (semi-quantitative or quantitative FFQ)

- ✓ Using food models, photographs or household measures to help estimate portion size.

Supplementary questions (about cooking methods, brand names, etc.)

Open ended section

- ✓ Respondents may record consumption of other foods that are not included in the close-ended food list.

Mode and time of administration

- ✓ Self-administered using paper or web-based formats, or interviewer administered via face-to-face or telephone interview.
- ✓ To account for seasonality, the survey can be administered at different times of the year (different foods may be available for consumption during different seasons).

Method used for recording (e.g. pen and paper, scannable format)

Length of FFQ

- ✓ To reduce respondent fatigue and reporting error, FFQ length should not be too drawn out and food items should be carefully selected.
- ✓ Increase the number of foods included in a FFQ and at the same time keep the length of the questionnaire short by grouping together items based on food classification or nutrient similarity.

Reference period for the FFQ: e.g. previous weeks, months, etc. Bear in mind that FFQ may not be suitable for recalling diet in the distant past (e.g. the previous year)

Availability of a food composition database

- ✓ Ensure that a food composition database is available which is up-to-date and complete, and includes locally available foods.

BOX 2. BRIEF DIETARY ASSESSMENT

For some assessment situation, a full-length FFQ questionnaire is not practical. Therefore, brief dietary instruments, sometimes referred to as screeners or short dietary instruments, can be employed in these circumstances. Screeners are used to measure the frequency of consumption without including information on portion size or dietary behaviour, via self- or interviewer-administered modes (Perez Rodrigo *et al.*, 2015). They are modified versions of longer FFQs varying in length, frequency categories and number of foods listed. An example is presented in Appendix 2. Consequently, screeners are used in situations when there is no need for comprehensive assessment. They are also used for surveillance, to screen individuals for inclusion in intervention or clinical trials, to identify and separate large numbers of individuals into groups or to distinguish individuals with low or high intakes. Depending on the specific objective of the assessment, brief dietary instruments can be referred to by different names. Instruments that assess specific nutrient and/or food group intakes are often called targeted instruments, and those that assess both dietary and non-dietary components are called brief multifactor instruments (Perez Rodrigo *et al.*, 2015; Thompson *et al.*, 2013).

Brief dietary instruments have further been demonstrated to be useful in low resource settings, specifically, by field workers with limited training in conducting large dietary assessments. Examples of these instruments include the Rapid Rural Appraisal (RRA) and Participatory Rural Appraisal (PRA), developed by Chambers (1994). These instruments are used to rapidly identify key risk factors for malnutrition or inadequate consumption patterns for specific food groups and allow for immediate action to take place (Perez Rodrigo *et al.*, 2015). Screeners have additionally been used to measure diet–disease association, as shown in a study by Jilcott *et al.* (2007). Women from impoverished and resource limited settings in the United States were asked to undertake a brief dietary instrument which included 54 questions on foods commonly eaten in the region. The results were used to provide guidance for dietary counselling.

Additionally, validation of brief dietary instruments have been conducted comparing the method with 24-hour recall. For example, this comparison was examined by Yaroch *et al.* (2012), where the performance of a 16-item screener used in a US adult population to assess fruit and vegetable intake, was compared to multiple 24-hour recalls. A Pearson correlation coefficient for the screener compared to the 24-hour recalls was low ($r=0.37$). The authors concluded that while dietary screeners are a less burdensome and cost-effective method for collecting information on fruit and vegetable intake, it is not recommended to use when trying to measure precise intake levels but rather to be used to obtain overall estimates, and/or to rank individuals with regard to intake levels of a particular food group.

2.2.1.2 24-hour recall

During a 24-hour recall, respondents (i.e. adults, children and their parents or caretakers) are asked, by a nutritionist or dietitian who has been trained in interviewing techniques, to recall and report all foods and beverages consumed over the preceding 24 hours. The 24-hour period starts with the first thing eaten by the respondent in the morning until the last food item consumed before he/she got up the next morning. Thus, the method assesses the actual intake of individuals. However, a single 24-hour recall is not enough to describe an individual's usual intake of food and nutrients. To achieve this objective, multiple non-consecutive 24-hour recalls on the same individual are required in order to capture daily variability (Baranowski, 2013; Gibson, 2005; Patterson *et al.*, 2004). Additionally, multiple day data collection can be used as a comparison method to validate an FFQ. Multiple 24-hour recalls can increase quality control, minimizing errors and maximizing reliability. Further information on the number of recalls required to estimate intakes can be found in section 3.1.

Information on 24-hour recall is collected using an open-ended format. Quantitative information on food intake, as described using portion size, allows for the calculation of energy and nutrient intakes (please refer to Appendix 3 for an example of a 24-hour recall form). Estimation of portion size is facilitated by the use of measurement aids such as standard household measures, photo atlases, food models, etc. To calculate energy and nutrient intakes, the estimated portion size or the amount of food intake is multiplied by the values of nutrient content in foods as found in the food composition tables or databases (for more information on food composition tables, refer to section 3.5).

As 24-hour recall is dependent on memory and cognitive functions, it can be challenging to

study in young children (under the age of eight years), or in elderly and institutionalized groups. In these cases, proxy/surrogate reporters can be used to provide additional assistance during the interview. However, it should be highlighted that the use of surrogate reporters can also introduce error, especially if the respondent is under the care of multiple caregivers. The interview(s) can take place either face-to-face, by telephone, or via a self-administered computer-assisted 24-hour recall version. The latter relies on literacy and an internet connection. It is crucial that interviewers are well trained in order to gather precise, detailed and accurate information to reduce error and bias. They should be familiar with the dietary patterns of the respondents, have a list of foods commonly eaten by the target population and be familiar with composite dishes, their recipes and preparation methods, and be aware of how food is served. Lastly, they should have training on how to use standard probes and prompts properly, how to measure portion size, particularly for mixed dishes, and how to ask questions in a non-judgmental and non-influential manner during the recall.

The importance of using a standardized interview structure such as the 'multiple pass approach' has been found useful (Gibson, 2005). Currently, the United States Department of Agriculture (USDA) computerized Automated Multiple-Pass Method¹¹ (AMPM) uses this approach to collect dietary data in the National Health and Nutrition Examination Survey, either in person or by telephone. This approach employs five steps. **Step 1:** 'quick list' (a list of foods and beverages consumed during the previous 24 hours is collected); **step 2:** 'forgotten foods' (probes for foods possibly forgotten during step 1); **step 3:** 'time & occasion' (time and occasion for each food is collected); **step 4:** 'detail cycle' (detailed description, amount, and additions for each food is collected); and **step 5:**

¹¹ <http://www.ars.usda.gov/News/docs.htm?docid=7710> (Accessed 1 December 2016)

'final probe' (probes for anything else consumed in the previous 24 h) (Blanton *et al.*, 2006).

2.2.1.2.1 *Applications and uses of 24-hour recall*

Depending on the number of days recorded, 24-hour recalls can be used to measure different outcomes. When a study aims to determine the mean intakes for a group or population, a single 24-hour recall per respondent can be carried out, especially when the sample size is sufficiently large. On the other hand, when the objective is to estimate the distribution of intakes, or to examine usual diets and correlations of individual intakes, more than one recall day per respondent is required.

A cross-sectional study examining rural Kenyan adults used two non-consecutive 24-hour recalls to investigate dietary patterns. The quantitative recall attempted to ensure reliable measurements of food and nutrient intakes by using local household measures, local food models and food composition tables (Hansen *et al.*, 2011). Due to the cross-sectional nature of the study, the authors noted that a major limitation was the inability to measure the impact of seasonal variation when assessing dietary patterns.

A non-consecutive two-day 24-hour recall has also been used to monitor household diet adequacy during three different seasons in rural Mozambique (Rose *et al.*, 2003). The decision to use 24-hour recall was related to the simple and inexpensive nature of the method when compared to other assessment methods, such as dietary history and food record. Furthermore, the authors found that conducting multiple 24-hour recalls at different times of the year was useful in accounting for seasonal variation in dietary intakes. Albuquerque *et al.* (2015) used non-consecutive three-day 24-hour recalls to evaluate the association between stature and total energy expenditure of low-income women in Brazil. Portion size was estimated by trained nutritionists with the aid of a photographic atlas.

24-hour recalls have been used in a number of other studies, including: a quasi-experimental complementary feeding intervention in Lombok, Indonesia (Fahmida *et al.*, 2015); a nutrition assessment in three specific regions of Ecuador (Sanchez-Llaguno *et al.*, 2013); a study measuring intakes of young children living in an impoverished South African setting (van Stuijvenberg *et al.*, 2015); and a study examining the impact of water intervention on beverage substitution in a randomized trial of overweight and obese Mexican women (Hernández-Cordero *et al.*, 2015). In summary, 24-hour recall has been used to assess the total dietary intake both of individuals and populations, and to examine the relationship between diet, health and other variables.

2.2.1.2.2 *Reproducibility and validity of 24-hour recall*

There are a number of issues to be considered when undertaking a reproducibility study for a 24-hour recall. These considerations include, but are not limited to, the following factors:

- Using non-consecutive days when conducting multiple 24-hour recalls is recommended since eating habits from consecutive days have been shown to be correlated (Hartman *et al.*, 1990);
- The effect of seasonality on the diet and changes in food availability should be taken into account;
- In repeated 24-hour recalls, action should be taken to avoid the first recall influencing the collection of data from the subsequent recall;
- Rankin *et al.* (2012) determined the reproducibility of two- to five-day repeated 24-hour recalls among urban African adolescents. This study was used to identify the optimum number of 24-hour recalls that need to be conducted in order to give the best reproducibility result. Their findings showed that conducting four or five recalls ensured higher reproducibility when compared to repeating the recall only two or three times.

There are a variety of issues to be considered when addressing the validity of 24-hour recalls. These considerations include, but are not limited to:

- The individual's accuracy in recalling or recording their consumption and estimating portion sizes;
- The comprehensiveness of the food composition tables to capture all foods available for the study;
- Types of foods frequently forgotten or overlooked, e.g. butter and salad dressing;
- Understanding that respondents are prone to over-reporting low intakes and under-reporting high intakes. This pattern is referred to as 'flat slope syndrome' (Baranowski, 2013).

It has been suggested that for the validation of 24-hour dietary recalls, the use of dietary histories or 7-day weighed records is not appropriate, due to differences in the time frame of the recall period and the potential for increasing the burden on the respondent. A single weighed food record may be considered as an alternative validation method (Gibson, 2005). A number of studies have reported good agreement between the methods of 24-hour recall and one day weighed record with trained observers and biomarkers (Thompson *et al.*, 2013).

Estimates from a multiple-pass interactive 24-hour recall in rural Ethiopian women have been compared with estimates from weighed food records (Alemayehu *et al.*, 2011). Their results showed that the median daily intakes of energy and most nutrients obtained were lower when measured by 24-hour recall than by weighed food record ($P < 0.05$). Furthermore negative bias for energy and nutrient intake were confirmed by Bland–Altman plots. The authors concluded that the two methods were not comparable in this setting: a result which could be attributed to the poor portion size estimation. Thus, extra attention must be devoted to the selection of the aids and

tools used to estimate portion size.

In another study, a 24-hour recall to assess the diets of Kenyan children, using mothers as proxies, was validated against a weighed food record (Gewa *et al.*, 2009). Energy intake was underestimated by approximately 6 percent by the mothers and 9 percent by the children. It was stated that further improvements were needed to allow for more accurate recalls to be collected, and to help increase recall of foods that typically go under-reported, namely, sugars, fruits, dairy products and meats. The authors remarked that assessing intakes in large-scale studies might not be feasible until improvements have been made.

Biomarkers have also been used to validate 24-hour recall. Scagliusi *et al.* (2006) conducted a study measuring the level of underreporting of energy intakes in a female Brazilian population, using the doubly-labelled water (DLW) method to validate a 3-day multiple pass 24-hour recall. The authors concluded that the energy intake measured by the multiple-pass 24-hour recalls presented a significant difference compared to energy expenditure measured by doubly-labelled water ($p < 0.0001$). Therefore, 24-hour recall was shown to underestimate energy intake. The authors noted that the outcome of the study could be related to the methodological approach, i.e. number of recall days, portion size estimation, measurement aids and food composition tables.

2.2.1.2.3 Strengths and limitations of 24-hour recall

The following table will provide a summary of the strengths and limitations associated with conducting a 24-hour recall.

Table 4 - Strengths and limitations of 24-hour recall

STRENGTHS	LIMITATIONS
<ul style="list-style-type: none"> Assesses the usual intakes of a large population (provided that the sample population is representative and the days of the week are adequately represented) Captures information on eating patterns, preparation methods, place of consumption, etc. The mode of administration does not affect food choice and eating pattern. Open-ended format used is appropriate for all eating patterns. Recalls intakes over the last 24-hours, therefore there is less burden on the respondents' memory, leading to better accuracy and response rates. Interview-based 24-hour recall does not rely on the literacy and numeracy skills of the respondent. 	<ul style="list-style-type: none"> Needs multi-day recalls to adequately represent the habitual intake of individuals and to take into account seasonality differences. Relies on respondent's memory. Requires well-trained interviewers with knowledge of cultural practices, eating habits, local recipes and preparation methods. Expensive due to the fact that extensive training is needed for the interviewers and the time spent on data entry and food matching with food composition data. Recall bias may be present, as respondents can be selective with the foods they choose to report during the recall. Relies on respondent's literacy and ability to describe the food and to estimate its portion size (self-reported 24-hour recall).

BOX 3. QUICK GUIDETO USING A 24-HOUR RECALL

Project objectives and budget will determine the study design and sample size. It is important to

- ✓ Understand the characteristics of the target population.
- ✓ Define the purpose and research questions of the study.

Requires trained nutritionists and dietitians to conduct this form of assessment

Target population group

- ✓ Characteristics: population groups (e.g. toddlers, pregnant women, elderly, etc.), age, literacy level, numeracy skills and cognitive abilities of the respondents assessed will have an impact on the mode of administration used.

- ✓ Parents can act as a proxy for children less than eight years old and can assist older children.
- ✓ Surrogate reporters can be used as proxies for the elderly if there is evidence of cognitive decline. Note that this increases the chances of error, particularly if the individual is under the care of multiple caregivers during the day.

Food intake and meal patterns. It is important to

- ✓ Understand the food intake and meal patterns of the target population group and also identify specific subgroups of the population (e.g. shift workers, pregnant and lactating women) who may have different meal patterns.

Mixed diets (composite diets). It is important to

- ✓ Understand local recipes, identify and record all ingredients consumed and have a strong appreciation of how to measure portion sizes of mixed dishes. For more information on recording mixed dishes, readers are referred to Gibson *et al.* (2008) pp. 47 to 67 on recording mixed dishes.

Information on dietary supplements (e.g. vitamins, minerals)

Mode of administration

- ✓ Face-to-face interview, computer-assisted recall or telephone administered recall.
- ✓ The researcher should explain to the participants that the goal of this interview is to record everything that the participant ate (meals and snacks) the day before, starting with the first thing eaten by the respondent in the morning until the last food item consumed before he/she got up the next morning.

Method used for recording information (e.g. pen and paper, scannable format)

Number of recorded days

- ✓ Selecting the number of days that will allow for an appropriate estimation of an individual's usual intake.
- ✓ A single 24-hour recall does not represent an individual's usual diet (hence multiple days are needed), nor does it take into account daily, weekly or seasonal variations of an individual's food intake.
- ✓ The time frame for a 24-hour cycle needs to be defined using reference points applicable to the target population (e.g. first food/drink consumed after waking up in the morning to the last food/drink consumed before going to sleep at night).

Days selected for conducting the assessment (weekday vs. weekend)

- ✓ Non-consecutive days are preferable, helping to capture more of the variability in an individual's diet.

- ✓ Including one weekend day in a week is desirable, to capture variability of food intakes during weekends.

Knowledge and skills of the interviewer. Interviewers should

- ✓ Receive training on conducting a 24-hour recall with standardized procedures, including practice interviews prior to the start of the study.
- ✓ Know how to probe the respondent using standardized and non-leading questions that are specific to the food consumption patterns of the target population.
- ✓ Ideally have knowledge of local foods, eating patterns, food preparation methods and the specific cultural practices of the study population.

Conducting a pilot study

- ✓ Select subjects and geographic sites that are representative of the actual target population for the pilot study.
- ✓ Identify any logistical and/or technical problems in the pilot study in order to fine-tune the survey procedures and to identify problems that may occur in the interviews prior to the actual survey, such as discrepancies in interview protocol, recalling and recording of composite dishes and food matching with food composition data.

Estimating portion sizes

- ✓ Using food models, photographs (photographic atlas) or standard household measures to help estimate portion sizes and food intake.
- ✓ For liquids (e.g. soups or beverages), record quantities as volumes, preferably using the respondents' own household utensils after these have been calibrated.
- ✓ Conversion factors or a food composition database are required to convert household measures to weight equivalents (Gibson, 2005).

It is necessary to have access to a food composition database which is up to date, complete and includes locally available foods as much as possible for nutrient estimation

- ✓ Refer to sections 3.4 and 3.5 'Estimation of portion sizes' and 'Availability of food composition data'.

Assessing validity and reproducibility

Procedures to minimize errors

- ✓ Train interviewers prior to the recall to become familiar with the dietary patterns of the study population.
- ✓ Create a standardized interview protocol.
- ✓ Calibrate utensils in the home and use standardized methods for portion size estimates.

- ✓ Use effective probes/prompts to reduce respondent memory lapses.
- ✓ Utilize multiple-pass interviewing techniques.

Reviewing the recall data

- ✓ Check and identify errors in the dietary data with the respondent during the interview. This should be conducted at the same time as the interview in order to ensure the most accurate information is obtained and to limit missing data.

2.2.1.3 Dietary history

Dietary history is a detailed assessment to describe usual food intake and its variation over a long period of time (six months to a year). The original dietary history designed by Burke in 1947 consisted of three parts (Biro *et al.*, 2002; Gibson, 2005):

- an in-depth interview to assess usual food intake and eating patterns;
- a food list;
- a three-day record with portion size estimates used as cross-checks.

Burke used the dietary history method to assess dietary intake. During an interview, the respondent was asked about their usual eating patterns at meal times and in between meals. Food consumed was recorded in common household measures. Questions like “What do you usually eat for breakfast?” were followed by further questioning to include daily variations until a full picture of food variety was obtained for breakfast. The portion size of food was also recorded in its real food size (i.e. a big, medium or small apple) or by using household measures. The interview continued in this way until a comprehensive collection of different kinds of food and their variations was recorded. The dietary history record was subsequently cross checked with a list of food groups, where the respondent was asked the frequency and quantity of a food that was consumed over a specific period of time (Burke, 1947).

The original method by Burke (1947) was however, considered impractical because it relied heavily on trained staff to perform the interviews and to code and enter the data. Modified versions of dietary history have since been developed and used to assess individual food intakes and meal patterns over a predefined period of time. The main difference between these variants and Burke’s original design is that the three-day record is often disregarded. Additionally, computerized versions have been developed to simplify the process, which can be self- or interviewer-administered. However this can still lead to high levels of respondent burden, usually associated with the long duration of conducting dietary histories (Gibson, 2005). When carrying out a three-part dietary history a trained nutritionist is needed to perform the interviews. Please find an example of a dietary history in Appendix 4.

2.2.1.3.1 Applications and uses of dietary history

Dietary histories were originally developed for clinical use, e.g. dietetic counselling, and are not often used in low resource settings, because they require well trained nutritionists with experience in dietary assessment. Computerized dietary history questionnaires, such as the Dietary Interview Software for Health Examination Studies (DISHES 98), have been shown to be useful in assessing dietary intakes and help to reduce the average interview and coding time.

DISHES 98 was used in the German National Health Interview and Examination Survey of 1998. It was designed to assess the usual dietary intake over a 4-week period in a subsample of an adult and elderly German population (Mensink *et al.*, 2001). Dietary history has also been used to study the relationship between early growth and diets and its link to the development of obesity and hyperlipidaemia in Hong Kong children. The dietary assessment was carried out by trained research dietitians to estimate portion size by using standard household measures such as bowls, cups and spoons to indicate the amounts consumed (Leung *et al.*, 2000). The Coronary Artery Risk Development in Young Adults (CARDIA) study employed an interviewer-administered dietary history to collect reliable quantitative data on the intake of young adults from the United States of America (Carnethon *et al.*, 2004). The assessment included a short questionnaire on general dietary practices, an FFQ on the usual intake of foods during the last month, and lastly a follow-up on the portion size and frequency of consumption of each food item selected. The data gathered by the researchers resulted in the collection of detailed information on the habitual eating habits and patterns (McDonald *et al.*, 1991). Dietary histories have also been used to determine the relationship between dietary patterns and tooth decay in a low-income African-American population. The dietary history revealed a high level of sugar and fat intake, with a low level of fruit and vegetable consumption in the study group that contributed to an increased incidence of tooth decay (Lim *et al.*, 2008).

2.2.1.3.2 *Reproducibility and validity of dietary history*

Compared to other assessment methods, there are a limited number of studies in validating dietary history as it is often used as a reference method to validate other methods of dietary assessments, e.g. 24-hour recall and FFQ

(Thompson *et al.*, 2013). A validation study by Mensink *et al.* (2001), compared the results from the computerized version of dietary history, DISHES 98, with results from a 3-day weighed dietary record and 24-hour recall. This study, described above in *Application and uses of diet history*, provides a good example of how the dietary history method has been validated. For most nutrients measured, the mean intakes recorded by DISHES 98 were lower than those measured by the 3-day weighed food records with an average Pearson's correlation coefficient of 0.51. A similar outcome was noted by the authors when the results from DISHES 98 were compared with a 24-hour recall, presenting an average Pearson's correlation coefficient of 0.46. The similarity observed could be due to the tendency of 24 hour-recall and dietary history to be influenced by similar error sources, since they both rely on the memory of the respondent. The authors concluded that the validity of DISHES 98 was deemed comparable to those of other dietary histories and the method was valid for assessing dietary intake (Mensink *et al.*, 2001).

Comparison of results between dietary history and biomarkers have also been demonstrated. Hagfors *et al.* (2005) conducted a validation study to compare the results of a dietary history to estimate energy, protein, sodium and potassium intakes against bio-markers. Protein, sodium and potassium estimates were compared with respective markers from a 24-hour urine sample and good agreement was observed ($r=0.58$). Overall, the authors noted that dietary history methods are adequate in assessing dietary intake, however, due to the small sample size of the study, ($n=32$), results could not be generalized.

The validity of the dietary history questionnaire was also tested against biomarkers to measure urinary excretion of nitrogen and plasma levels of carotenes, vitamin E and vitamin C. The subjects were adult males and females from

the Spanish arm of the European Prospective Investigation into Cancer and Nutrition (EPIC) study. The correlation coefficient between the dietary history questionnaire and nitrogen excretion was 0.58. The validation revealed that the correlation between the questionnaire and the plasma levels of vitamin C, β -carotene and carotenoids, and dietary intake, were 0.46, 0.33 and 0.42 respectively. From these results,

the authors concluded that the dietary history method is able to provide reliable information on usual intake of the nutrients measured in the study (Gonzalez, 1997).

2.2.1.3.3 Strengths and limitations of dietary history

The following table will provide a summary of the strengths and limitations associated with conducting a dietary history.

Table 5 - Strengths and limitations of dietary history method

STRENGTHS	LIMITATIONS
<ul style="list-style-type: none"> Provides details of meal patterns, individual foods consumed and usual food intake after completing a single interview 	<ul style="list-style-type: none"> Relies on respondent's memory, can lead to recall bias
<ul style="list-style-type: none"> Provides quantitative estimates of energy and nutrient intakes 	<ul style="list-style-type: none"> Labour-intensive, time-consuming, may not be suitable for young children and elderly respondents
<ul style="list-style-type: none"> Useful to describe usual food or nutrient intake over a relatively long period of time. It can be used to estimate prevalence of inadequate diets 	<ul style="list-style-type: none"> To obtain detailed information on food intake, longer interview times are needed, resulting in high respondent burden
<ul style="list-style-type: none"> Does not rely on the literacy of the respondent 	<ul style="list-style-type: none"> Portion size estimation of past meals can be difficult, even with the use of aids
<ul style="list-style-type: none"> Provides information on foods that are not regularly consumed 	<ul style="list-style-type: none"> Requires trained personnel with knowledge of local food culture and eating patterns (interview-based dietary history)
<ul style="list-style-type: none"> Does not interfere with normal eating habits 	<ul style="list-style-type: none"> Requires literate respondents with the ability to estimate portion size (self-administered dietary history)
	<ul style="list-style-type: none"> Expensive to administer
	<ul style="list-style-type: none"> Data entry and coding is time consuming and requires trained personnel

BOX 4. QUICK GUIDE TO USING A DIETARY HISTORY METHOD

Project objectives and budget determine the study design and sample size

- ✓ Understanding the specific target population, and the purpose and guidelines for the study.

Population characteristics

- ✓ Age (e.g. toddlers, adolescence, elderly, etc.), pregnant/lactating, literacy level and cognitive abilities of the respondents will have an impact on the mode of administration.

Food and meal patterns

- ✓ Understanding the food and meal patterns of the target population and identifying specific subgroups of the population who may have different meal patterns (e.g. shift workers, pregnant and lactating women).

Mixed diets (composite diets)

- ✓ It is important to have an understanding of local recipes and to identify all ingredients consumed.

Information on dietary supplements (e.g. vitamins, minerals)

Mode of administration

- ✓ Face-to-face with an interviewer or computer-administered (interviewer-based or self-administered).

Length of assessment

- ✓ To reduce respondent fatigue and over-reporting, interview length should be kept short.

Reference period for the interview, e.g. previous weeks or months. Keep in mind that recalling diets from the distant past (e.g. previous year) may result in recall bias

Recording method

- ✓ Use of food models, photographs and/or standard household measures.
- ✓ Brand names, a complete description of the method of preparation and cooking, and the recipes for composite dishes should all be recorded.

Capacity of the dietary assessment coordinator

- ✓ A trained nutritionist is needed to conduct the interview.

- ✓ The nutritionist should have experience in gathering detailed information on the consumption of food and drink, and information related to the respondents' food habits, e.g. food allergies, seasonal variations and dietetic preferences, etc.
- ✓ They should know how to probe the respondent using standardized and non-leading questions.
- ✓ The dietitian or nutritionist needs knowledge of local foods (including brands), preparation methods, recipes and portion size. For more information, readers are referred to Gibson *et al.*, (2008) on recording mixed dishes.

Availability of a food composition database

- ✓ Ensure a complete and up-to-date food composition database is available which also includes locally available foods.

For practical tips on how to conduct an FFQ, see Box 1: 'Quick guide to using an FFQ'. For practical tips on how to conduct a three-day food record for verification, see Boxes 5 and 6: 'Quick guide to using an estimated food record', and 'Quick guide to using a weighed food record'.

2.2.2 PROSPECTIVE DIRECT METHODS

In the prospective methods, diet, including all food and beverages consumed, is recorded at the time of consumption, therefore allowing for current food intake to be recorded. These methods include a weighed food record, an estimated food record and a duplicate meal method. Prospective methods can be more labour intensive depending on the objectives of the study (i.e. the weighed food record and duplicate meal method) than retrospective methods and rely heavily on respondents having good literacy and numeracy skills.

2.2.2.1 Estimated food records

When conducting an estimated food record, respondents are instructed to document all foods and beverages consumed during a predefined period (e.g. 1 to 7 days). The number of days included in the assessment depends on the purpose of the study (Gibson, 2005). If the objective is to collect information on an average

intake of a population group, then one record day will suffice. However, if the purpose of the assessment is to obtain habitual intakes for individual respondents, then more than one day is needed, including one weekend day (Gibson, 2005). During the record period, specific details such as brand names, time of day the food or beverage was consumed, the location and sometimes a description of the occasion should be documented as well. Estimation of the food portion or food weight consumed is normally aided by using standardized household measures, food photographs or models (please refer to Appendix 5 for an example of an estimate food record form). Furthermore, food recording taking place during meal time should take place simultaneously in order to minimize reporting errors due to fading memory. Prior to data collection, respondents are often provided training and practice on recording, in order to become familiar with the steps of the assessment, especially for foods eaten outside the home. Moreover, ways to record food preparation, cooking methods and mixed dishes will also be covered prior to starting the assessment

method. Complete instructions should be given to the participants prior to the start of the project in an effort to minimize errors. In cases where respondents cannot record their consumption, trained interviews, parents, family members and/or caregivers might provide assistance to record intake. To ensure that data collection is accurate, a home visit by trained interviewers on the first day of recording is helpful. In addition to home visits, at the end of the recording period, skilled interviewers should go through the records, clarify the entries, and ask the respondent for any omitted items. After data collection is completed, calculation of nutrient intake is done using a food composition database.

2.2.2.1.1 Applications and uses of estimated food records

Estimated food records are useful in assessing detailed food and nutrient intakes at individual level. A cross-sectional study on obese adolescents in Malaysia examined the role of diet and physical activity on adult morbidity and mortality (Rezali *et al.*, 2012). A self-reported 3-day estimated food record was applied, the respondents were trained on how to use the food record form and how to estimate portion size using standard household measures to assess all food and beverages consumed along with the descriptions of brands and methods of food preparation. The authors commented that a major limitation of the method was the self-reported nature of the study, which can result in under-reporting of dietary intake due to incomplete reporting of the type or frequency of foods and snacks consumed.

Furthermore, estimated food records have been used in large-scale monitoring studies, including the National Diet and Nutrition Survey (NDNS) and the National Survey of Infant and Young Children (DNSIYC) (Stephen *et al.*, 2013). The former is designed to assess the nutritional status of babies aged 18 months or more, living in private households in the UK, while the latter is a single survey of infants and young children

throughout the UK. As part of the survey, a 4-day estimated food record – including two weekend days – was collected to replace a previously used repeated 24-hour recall. The decision for the replacement was based on the fact that energy intake was found to be similar for both methods when compared to DLW. 24-hour recall, however, showed more over-reporting in young children and respondents reported excess burden when conducting 24-hour recalls (Stephen *et al.*, 2013).

2.2.2.1.2 Reproducibility and validity of estimate food record

Methods used to validate estimated food records have included weighed food records and biomarkers. Chinnock (2006) validated a 3- and 4-day estimated food record against those obtained from a 7-day weighed food record in Costa Rican adults from both rural and urban areas. Interviewers visited the respondent's homes during meal preparation and consumption, and observed the weighing of ingredients, and recording of food intakes, as well as left-over food. Training of subjects on data collection was provided before the study. Energy and nearly half of the nutrient intakes were underestimated by the food records obtained. Correlation coefficients ranged from 0.68 for polyunsaturated fats to 0.87 for calcium, indicating good to high agreement. The authors concluded that the results were comparable with other similar studies (Bingham *et al.*, 1994; Bingham *et al.*, 1995; Bonifacj *et al.*, 1997). They supported the use of estimated food records in dietary surveys among Costa Rican adults and also noted that there was a greater difference between weighed and estimated food record estimates for energy in rural residents, where energy intake was underestimated by 312 Kcal. They speculated that subjects in rural areas underestimated the amount of rice and beans consumed. As a result, the authors recommended the use of additional aids in estimating intakes of these two specific foods in subjects living in rural areas (Chinnock, 2006).

Biomarkers have also been used to validate estimated food records. In the EPIC Norfolk cohort (described earlier in the dietary history section), the performance of a 7-day estimated food record was assessed against biomarkers of urine (nitrogen, potassium and sodium) and blood (plasma ascorbic acid). Reproducibility was also measured by comparing one 7-day food record with a second 7-day food record. Results of the reproducibility study did not show a significant difference ($p < 0.001$). For validity, the average correlation coefficients between urinary nitrogen and dietary nitrogen were relatively high ($r = 0.57$ –

0.67); between urinary potassium and dietary potassium they were good, ($r = 0.51$ – 0.55); and between plasma ascorbic acid and dietary vitamin C they were more moderate ($r = 0.40$ – 0.52). The authors concluded that the food record used in the study was a suitable method to estimate dietary intake (McKeown *et al.*, 2001).

2.2.2.1.3 Strengths and limitations of estimated food records

The following table will provide a summary of the strengths and limitations associated with conducting an estimated food record.

Table 6 - Strengths and limitations of estimated food records

STRENGTHS	LIMITATIONS
<ul style="list-style-type: none"> • Provides estimates of actual instead of usual diet. This is dependent on the number of measurement days • Open-ended format appropriate for all eating patterns • Provides a high level of specificity and details regarding foods consumed and occasion • Gives detailed information on eating patterns • Allows for the collection of information from respondents with sporadic eating habits • Does not rely on respondent's memory, since information is recorded at the time of consumption • Allows for real time portion size estimation, reducing errors in the estimation of intake 	<ul style="list-style-type: none"> • Requires literate and motivated respondents with numeracy skills • In low-resource countries, a trained investigator/interviewer is required to collect data • High cost for administration and data analysis • Time consuming and can result in a high level of respondent burden and lower cooperation • Respondents may forget to record specific food items or whole meals • Portion estimation can be difficult to compute if inadequate information is recorded by the respondent • May interfere with normal eating habits • Reliability of records decreases over time, due to increased respondent burden • Missed or less accurately recorded food intake in children when foods are eaten when the child is not in the presence of parents or caregivers

STRENGTHS	LIMITATIONS
	<ul style="list-style-type: none"> • May not capture foods eaten less frequently • Data entry and coding can be time consuming and requires trained staff

BOX 5. QUICK GUIDE TO USING AN ESTIMATE FOOD RECORD

Project objectives and budget determine the design of the study and sample size

- ✓ Understanding the specific target population, purpose and guidelines for the study.

Population characteristics

- ✓ Age (e.g. toddlers, adolescence, elderly, etc.), pregnant/lactating, literacy level and cognitive abilities of the respondents assessed will have an impact on the mode of administration.

Mode of administration

- ✓ Can be completed by the respondent, parent or caretaker.
- ✓ Investigators are advised to conduct home visits during the study period to ensure proper recording procedures.

Number of recorded days

- ✓ Should be selected based on the objectives of the dietary assessment.
- ✓ A single-day food recording is not sufficient to capture a typical diet and does not take into account daily, weekly or seasonal variations in food intake.
- ✓ A standard 3–7 day food record is typically used. Recording periods of more than seven consecutive days can lead to respondent fatigue.
- ✓ Repeated food records conducted at different times of the year or in different seasons will give a better picture of habitual intake.

Days selected for conducting the assessment (weekday vs. weekend)

- ✓ Non-consecutive days are preferable to capture more of the variability in an individual's diet.
- ✓ Including one weekend day in a week is helpful in capturing variability of food intakes during weekends.

Recording methods

- ✓ Use of food models, photographs and/or standard household measurements (cups, tablespoons, etc.), supplemented with measurements using a ruler (e.g. meat and cake) and counts (e.g. bread slices).
- ✓ Food intake can be recorded at the time of consumption or using a portable recording device.
- ✓ Brand names, the method of food preparation and cooking, and recipes for composite dishes should all be recorded.

Capacity of the dietary assessment coordinator

- ✓ Trained dietitians or nutritionists with knowledge of local foods (including brands), preparation methods, recipes, food ingredients and portion size. For more information, readers are referred to Gibson and Ferguson, 1999, pp. 47-67, on recording mixed dishes.
- ✓ To check with the respondent to identify forgotten foods using standardized and non-leading questions.

Capacity of the respondent

- ✓ Understand the instructions for the study (supplied with an instruction booklet to be used during food recording period).
- ✓ Investigators provide training before the study for participants to practice food recording.

Availability of a food composition database

- ✓ Ensure a food composition database is available, up to date and complete, and includes locally available foods.

2.2.2.2 Weighed food records

The weighed food record is often regarded as the most precise method for estimating the food and nutrient intakes of individuals. This approach adopts the same methodological principles as the estimated food record method. However, in this method respondents are asked to weigh using weighing scales (e.g. a digital electronic scale with a tare button to facilitate food weighing). The subjects directly copy the weight of an individual food from the scale while conducting additive weighing without the necessity of manipulating figures, and thus avoiding errors (Marr, 1971). All food and beverages consumed are weighed and recorded, along with a description of portion size, brand names and details on food preparation (please refer to Appendix 5 for an example of a weighed food record form). Similar to the estimated food record, a predefined measurement period (between 1 to 7 days) needs to be set, where the number of days included in the assessment will depend on the purpose of the research question and again, weekend days should be included to account for changes in food intake during weekend days. To obtain even more precise data, leftovers can also be weighed or estimated. For studies requiring a very high level of accuracy it may be necessary to weigh and describe all raw ingredients separately before cooking, and this is especially important for mixed dishes (Gibson, 2005). For foods consumed away from home, the respondents are asked to provide a full description of the foods consumed, the occasion and location. Depending on the design of the study, respondents can be asked to weigh foods consumed outside the house, or the researchers will purchase the particular food from a shop or restaurant to weigh the described portion size at a later time. Prior to the recording period, the respondents should be provided with instructions for recording, with sufficient time for practicing and reminders about maintaining typical dietary habits during the recording period. Furthermore, sufficient supervision during the

study is a prerequisite to obtain accurate and reliable data. Hence home visits are normally arranged during the study period to ascertain that the procedures are carried out properly (Bingham, 1987).

As with estimated food records, in cases where respondents are unable to record their own consumption, interviewers, parents, family members and/or caregivers might be required to weigh and record the food intake. To ensure the accuracy of data collection, home visits on the first day can be arranged to ascertain that procedures are carried out properly (Thompson *et al.*, 2013). It must also be noted that due to the increased respondent burden with the weighed food record, motivated individuals are needed to collect accurate data and avoid changes in usual eating patterns to simplify the measurement procedure. Once all the information from the weighed food record has been completed, food consumption data will need to be converted to nutrient intakes, which requires both a food composition database and an analysis programme.

2.2.2.2.1 Applications and uses of weighed food record

Weighed food records have been shown to be useful in collecting information for different purposes. This method has been used to collect data on group mean intakes, where a single record is sufficient, or to measure the distribution of individual intakes, where multiple record days are needed. The measurement of individual nutrient intakes has been shown in a study investigating pregnant subsistence farmers in rural Malawi using a 4-day weighed food record (Nyambose *et al.*, 2002). Enumerators living in the villages visited the homes of respondents during the study period, from the time they woke up until after the evening meal and weighed all food and beverages consumed. The raw ingredients of all dishes were weighed before cooking, followed by weighing the final cooked dish and the remaining uneaten foods. Furthermore, plates and cups

were supplied to all participants to assist with food weighing. The data were subsequently entered into a nutrient analysis programme updated with additional locally consumed foods. The authors noted a large within-person variation in nutrient intakes, which poses a challenge for dietary assessment and suggests that additional replicate days are required to estimate the mean intake of individuals in subsistence farming.

In addition, weighed food records have also been used to assess micronutrient intakes for example, vitamin A, folate, iron and zinc among young children in Bangladesh. Trained nutritionists recorded all food and beverages consumed, recipe ingredients and preparation methods and any leftover foods over a 12-hour recording period (Arsenault *et al.*, 2013).

2.2.2.2.2 *Reproducibility and validity of weighed food record*

Reproducibility studies conducted on weighed food records have presented good agreement amongst group mean values for energy and the majority of nutrients. These results have been presented in a study by Willett *et al.* (1995), where the reproducibility of a 7-day weighed food record was tested, using Pearson's coefficient and interclass correlation coefficients. The authors showed that the interclass correlation coefficients were between 0.41 and 0.79, with the lower number referring to vitamin A. The study concluded that 7-day weighed food records are sufficient for providing reproducible results (Willett *et al.*, 1995). Weighed food records are considered the gold standard against which other dietary assessment methods are validated; therefore, their relative validity cannot be assessed via other direct assessments (Gibson, 2005). Duplicate meal methods (explained further in section 2.2.2.3) have instead been used as a reference method to validate weighed food records (Gibson *et al.*, 1982; Gibson *et al.*, 2015). A 1-day weighed record was compared with a duplicate meal composite collected on the same day from a

group of rural Malawian women measuring iron intakes ($p < 0.001$). The authors concluded that the intake levels calculated from food composition databases were underestimated since they did not take into account iron contamination in the soil, which the duplicate meal composite was able to capture. Furthermore, recovery biomarkers like the DLW method and urinary nitrogen excretion have also been used to validate energy and nutrient intakes. The validity of a 16-day weighed record to measure habitual food intake was compared against the 24-hour urine nitrogen technique in women aged 50–65 years in the UK. Correlation between dietary and urinary nitrogen estimates was 0.69. In addition, the relationship between weighed records and other potential biomarkers (i.e. potassium and vitamin C) showed relatively high correlation between 24-hour urine potassium excretion and dietary potassium intake ($r = 0.73$), and between plasma vitamin C and dietary vitamin C ($r = 0.86$), among others. The authors concluded that weighed food records provided an accurate estimate of energy and nutrient intakes (Bingham *et al.*, 1995).

2.2.2.2.3 *Strengths and limitations of weighed food records*

The following table will provide a summary of the strengths and limitations associated with conducting a weighed food record.

Table 7 - Strengths and limitations of weighed food records

STRENGTHS	LIMITATIONS
<ul style="list-style-type: none"> • Assess the actual or usual intakes of individuals, depending on the number of measurement days • Is more accurate than other dietary assessment methods. It has been considered the gold-standard method for dietary assessment • Does not rely on memory, since information is recorded at the time of consumption • Provides exact portion sizes, and does not rely on estimation • Provides a high level of specificity and details regarding food consumed and meal patterns • Provides information on foods eaten regularly 	<ul style="list-style-type: none"> • Requires literacy, motivated respondents with numeracy skills (if self-reported) to weigh out foods and record food intakes • In communities with low literacy and numeracy, a trained field investigator is required to collect reliable data • Time-consuming and labour-intensive for both respondent and researcher • High level of respondent burden when compared to the other assessment methods • Respondents may alter eating habits to simplify the procedure due to study fatigue, especially if multiple days are recorded • Costly in both equipment and staff required to provide training and supervision • Errors resulting due to difficulties in weighing foods eaten away from home • Requires a suitable environment for weighing foods • May not capture foods eaten less frequently

BOX 6. QUICK GUIDETO USING A WEIGHED FOOD RECORD

Project objectives and budget determine the study design and sample size

- ✓ Understanding the specific target population, purpose and guidelines for the study.

Population characteristics

- ✓ Age (e.g. toddlers, adolescence, elderly, etc.), pregnant/lactating, literacy level and cognitive abilities of the respondents will have an impact on the mode of administration.
- ✓ Characteristics (e.g. toddlers, pregnant women, elderly, etc.), age, literacy level and cognitive abilities of the respondents assessed will have an impact on the mode of administration.

Mode of administration

- ✓ Can be completed by respondent, parent or caregiver.
- ✓ Investigators are advised to conduct home visits during the study period to ensure proper recording and weighing procedures.

Number of weighing days

- ✓ Should be selected based on the objective of the dietary assessment.
- ✓ A single day food weighing is not sufficient to capture typical diet and does not take into account daily, weekly or seasonal variations of food intake.
- ✓ A 7-day weighing record period is historically most common. A period less than seven days may underestimate the daily variations, while a period greater than seven consecutive days can lead to respondent fatigue.
- ✓ Repeated food records conducted at different times of the year/season will give a better picture of habitual intake.

Weighed recording method

- ✓ A scale (typically a digital electronic scale with a tare button) to weigh all food, drinks and non-foods (e.g. water, dietary supplements, alcohol) at the time of consumption, using a record sheet or a digital audio recorder.
- ✓ Brand names, description of the cooking method and recipes for composite dishes should be recorded. Plate waste is weighed and recorded separately.

Capacity of the dietary assessment coordinator

- ✓ Trained dietitian or nutritionist with knowledge of local foods, preparation methods, recipes and portion size.
- ✓ Ability to check with the respondent to identify forgotten foods using standardized and non-leading questions.

Capacity of the respondent

- ✓ Understanding of the instructions for the study (respondents are supplied with an instruction booklet and visual demonstrations).
- ✓ Competency on how to use weighing scales correctly to measure food and drinks and record composite dishes (weighing of raw ingredients). For more information, readers are referred to Gibson and Ferguson, 1999, pp. 47-67, on recording mixed dishes.
- ✓ Receive instruction and hands-on training prior to the start of the study on how to properly weigh and record foods and drinks.
- ✓ A phone hotline should be made available to provide technical support for the respondents during the study period.

Availability of a food composition database

- ✓ Ensure that a food composition database is available, up-to-date and complete, and includes locally available foods.

2.2.2.3 Duplicate meal method

The duplicate meal method involves setting aside duplicate portions of all foods and beverages consumed throughout a specific time period. These retained duplicate portions are set aside, weighed, either by the respondent or fieldworker, and then sent to a laboratory for chemical analysis to determine nutrient content. During the assessment period, respondents also maintain a weighed food diary to record details of the foods and beverages consumed during the assessment period and provide information on portion size expressed in weights or household portions. The food diary can help to verify the comprehensiveness of collecting duplicate foods and accuracy of the portion sizes of the duplicate meal (Lightowler *et al.*, 1998). In some cases, such as the study by Hurst *et al.*, (2013), enumerators resided in the household during the study period to weigh and create duplicate samples of all beverages and food items consumed, which are then subject to chemical analysis to measure nutrient content. A comparison of nutrient intakes can be made between the values of analysed nutrients and the nutrient intake which is calculated based on the reported portion size and data from a food composition database.

2.2.2.3.1 Applications and uses of duplicate meal method

The duplicate meal method is often used in institutional and metabolic balance studies (where subjects do not control the portions selected) (Pennington, 2004), and is considered the best method for assessing trace element intake (Abdulla *et al.*, 1981). This approach is especially useful to assess the intakes of nutrients/contaminants without nutrient

composition databases, and/or if the nutrients of interest are affected by soil concentrations, e.g. selenium, iodine and zinc. It is suitable for use in developing countries, especially for population groups with low literacy rates or in cases where a substantial portion of food is prepared at home. However, it is costly to purchase and prepare duplicate meals and to analyse nutrient contents by chemical analysis. Furthermore, it imposes a high burden on the respondent and is therefore unsuitable for large-scale studies. Examples of its application in a low resource setting area include a cross-sectional study investigating the risks of iron and zinc deficiency in women aged 15–50 years from six rural villages in Malawi (Siyame *et al.*, 2013). Trained research assistants collected a one-day weighed duplicate diet composite, along with recordings of the type of foods consumed. Diet composites were then transported to the laboratory for processing and analysis of trace elements. To help capture the variability of an individual's diet, weekdays and weekend days were proportionally represented. The authors described how the use of duplicate diet composites improved estimates of trace elements. However, due to the single day nature of measurement, usual intake level could not be obtained. Dietary exposure and assessment of trace elements, specifically the dietary fluoride intake of Ethiopian children was measured using a duplicate portion technique. Mothers were asked to prepare a duplicate portion of food and drink, identical in content and volume to the portion taken by the child, for four consecutive days including one weekend day. The portions were then collected and analysed for fluoride content (Malde *et al.*, 2004).

2.2.2.3.2 Reproducibility and validity of duplicate meal method

Duplicate meals are validated by comparison with biomarkers. However, since this method usually serves as a reference method, few studies have carried out reproducibility and/or validity studies to date. A validation study, conducted on women in Sweden, measured the extent to which the duplicate meal method estimates true dietary intake. Comparisons were made between the results from the duplicate portions and biological markers (urine and faeces collection). The duplicate meal method underreported the

diet by 14 percent, specifically underestimating the intakes of protein, sodium, potassium and calcium. The authors concluded that the method provided a good measure of dietary intake (Johansson *et al.*, 1998).

2.2.2.3.3 Strengths and limitations of duplicate meal method

The following table will provide a summary of the strengths and limitations associated with conducting a duplicate meal method.

Table 8 - Strengths and limitations of duplicate meal method

STRENGTHS	LIMITATIONS
<ul style="list-style-type: none"> • Provides the most precise information on nutrient intake, and it is not affected by the limitation of food composition data • Omission of foods consumed is minimized, when compared to other assessment methods 	<ul style="list-style-type: none"> • It is costly to prepare duplicate meals and conduct the chemical analysis of foods • Time-consuming and burdensome for respondents • Requires literate and motivated respondents with numeracy skills (if self-reported) • In low-resource countries, a trained field investigator is required to collect reliable data • Impractical to use in a large scale study population • Expensive for the participants to purchase and prepare extra food, relies on respondents to provide a complete duplication of foods consumed • Interferes with normal eating habits and may lead to underestimated intakes • Data entry and coding is time consuming and requires trained staff

BOX 7. QUICK GUIDE TO USING A DUPLICATE MEAL METHOD

Project objectives and budget determine the study design and sample size

- ✓ Understanding the specific target population, purpose and guidelines for the study.

Population characteristics and project setting

- ✓ Age (e.g. toddlers, adolescence, elderly, etc.), pregnant/lactating, literacy level and cognitive abilities of the respondents will have an impact on the mode of administration.
- ✓ Study conducted in a controlled environment (e.g. hospital) or unrestricted setting (e.g. home-based assessment).

Mode of administration

- ✓ Weighed record can be completed by the dietary assistant coordinator or by the respondent, parent, caregiver, etc.
- ✓ Investigators can conduct home visits during the study period to ensure proper recording and weighing procedures are followed.

Recording method

- ✓ Food items consumed are weighed, non-edible/non-consumed parts are removed, and duplicate samples are made for laboratory analysis.

Capacity of the dietary assessment coordinator

- ✓ Dietitian or nutritionist with knowledge and understanding of preparation foods for laboratory analysis.

For additional information please refer to the Box 6: 'Quick guide to using a weighed food record'.

2.3 INTEGRATION OF INNOVATIVE TECHNOLOGIES TO IMPROVE DIETARY ASSESSMENT

There is a growing need for more specific and accurate dietary assessment methods. High quality data are essential for research on the association between diet and health, as well as being decisive to understanding dietary

patterns and nutrition-related health problems such as micronutrient deficiencies, diet-related chronic diseases, obesity, cancers, etc. Given its importance, continuous efforts are being made to improve existing dietary assessment methods, as well as to develop more innovative alternatives that are less demanding.

A review published by Poslusna *et al.* (2009) indicated that the main factors influencing misreporting in traditional methods (i.e. dietary recalls, food records) were the reliance on

respondents' memories and their poor ability to estimate portion sizes. New information and communication technologies – i.e. personal digital assistants, mobile phones, interactive computer software – aim to overcome the limitations of a pen and paper method and to obtain more accurate and reliable dietary information. In addition, innovative technologies aim to reduce the cost of collecting and processing dietary information. Compared with traditional dietary assessment methods, new technologies have several advantages: they do not rely on the respondents' memory; they can automatically process data and provide real-time personalized dietary feedback advice. However, they also have a series of limitations: in particular, the feasibility and cultural suitability of integrating the latest innovative technologies in rural areas and in low resource settings, particularly among low-literacy populations, is still limited. This guide provides information on the most recent technologies used to improve dietary assessment in more developed settings, as well as some examples of new technologies used in low resource settings. Based on recent reviews (Forster *et al.*, 2015; Gemming *et al.*, 2015; Illner *et al.*, 2012; Stumbo, 2013), innovative technologies to improve dietary assessment have been classified into four key groups: personal digital assistant (PDA), image-assisted methods (i.e. digital cameras, mobile phones, tablets, etc.), interactive computer and web-based technologies, and scan and sensor-based technologies. In particular, this guide takes a detailed look at image-assisted methods, given their potential use in low resource settings.

2.3.1 PERSONAL DIGITAL ASSISTANT (PDA)

A PDA is a handheld computer that can be used for various purposes. For dietary assessment, a PDA has a specially-designed dietary software program that can be used to register and self-monitor dietary intake. It allows for the evaluation

of short-term dietary intake, facilitating real-time data collection. Prior to data collection, participants must receive training on how to handle the device. They are required to record their food intake right after consumption by selecting appropriate food items from a predefined list of foods. Early PDAs provided users with a selection of about 180 food items; current PDAs offer a much higher number, between 400 and 4 000 items (McClung *et al.*, 2009). However, it has been reported that PDAs increase the respondent burden compared with pen and paper reports, on account of their increasingly extensive list of foods (Welch *et al.*, 2007). PDAs can also provide some type of aid for measuring portion sizes, i.e. electronic prompts, discrete food photographs, picture books or food models, and household measures. These will help participants quantify the amounts of food consumed. For example, one PDA may display colour photographs of each food item together with a default amount (in grams); subsequently the participant can adjust the predefined portion size to the correct consumed amount. Data is then uploaded to be reviewed by dietitians and matched with food composition tables for further analysis (Beasley *et al.*, 2009; Illner *et al.*, 2012; McClung *et al.*, 2009).

2.3.2 IMAGE-ASSISTED DIETARY ASSESSMENT METHODS

Image-assisted dietary assessment methods refer to any method that uses images (i.e. photographs, videos) of food collected during eating episodes to enhance accuracy and reduce respondent burden. The purpose of using images is to support traditional self-report methods (24-hour recalls or records) or as a stand-alone method to provide the primary record of dietary intake (Gemming *et al.*, 2013; Lassen *et al.*, 2010; Lazarte *et al.*, 2012; Martin *et al.*, 2014).

To make use of image-assisted methods, participants should use handheld devices or wearable cameras to shoot images of food

and meals, before and after consumption. It is recommended that participants are provided with a fiducial marker (i.e. a fork or pen), and/or a marked tablecloth, which should be placed close to or beneath the food items before taking the images. The marker is useful to facilitate portion size estimation. Additionally, for increased accuracy the participants can be asked to write down or verbally record a description of food/meals that cannot be captured from images; this is particularly important for obscured images. Alternatively, a 24-hour recall can be conducted to collect further information such as the cooking method, hidden foods and added condiments.

In some cases, if Internet connection is available, respondents are requested to send the pictures immediately after the eating periods (Martin *et al.*, 2012). The images are reviewed by nutritionists and the consumed portion size can be estimated by different methods: from the digital images alone or by comparing them with images of reference portions for known food quantities, the latter being displayed on a computer screen or depicted in a printed food atlas (Lazarte *et al.*, 2012; Martin *et al.*, 2012). Calculation of nutrient intake is based on data from the corresponding food composition tables or databases, and the amount of consumed food (in grams).

Results of image-assisted methods showed that images combined with 24-hour recalls or food records enhanced self-report dietary assessment by revealing unreported foods and identifying misreporting errors not captured by traditional methods alone. This approach helps to reduce memory bias (Gemming *et al.*, 2013; Lassen *et al.*, 2010; Lazarte *et al.*, 2012; Martin *et al.*, 2012).

2.3.3 MOBILE-BASED TECHNOLOGIES

Mobile-based technologies are becoming increasingly accessible and have the potential to address some of the burdens associated with collecting accurate dietary data, allowing users to shoot images or make voice records using a

smartphone or tablet. Moreover, the rapid advance of mobile technology encourages researchers to develop dietary assessment methods based on this technology. This method is especially appealing for children and adolescents, who are often more motivated to use technology and therefore can become potential users (Boushey *et al.*, 2015; Casperson *et al.*, 2015).


Mobile-based technologies allow short-term dietary assessment, i.e. it is possible for these assessments to be conducted in real-time. Participants are asked to take photographs, video or voice records of all the foods and drinks they consume on eating occasions. The identification of foods in the images and portion size estimation can be carried out by the respondents or dietitians. One of the first devices developed with these characteristics was the Wellnavi instrument (Kikunaga *et al.*, 2007), which was initially a personal digital assistant (PDA) with camera (Wang *et al.*, 2006). Then, the Wellnavi was upgraded to a mobile phone with a camera for capturing images of meals. It included a display on which to write the names of the ingredients of the meal in the digital photograph. The images and the written information were then sent to dietitians for portion estimation and data analysis (Kikunaga *et al.*, 2007). Also with the rapid development in mobile technologies, many mobile applications ('apps') have been developed to record food intake on mobile devices. New technologies have been developed as well for food identification and portion size estimation based on automatic analysis of the food depicted in images through a system of digital image segmentation and analysis.

An example of this innovative technology is the mobile food record (mFR) based on the mobile application mFR–Technology Assisted Dietary Assessment (TADA) (Khanna *et al.*, 2010). The mFR–TADA method has the potential to identify foods automatically and estimate portion sizes via the estimation of food volume from the before-

and-after-eating images (Boushey *et al.*, 2015; Zhu *et al.*, 2010). A fiducial marker is included in the images as a measurement reference to help with the reconstruction of a three-dimensional image that allows for estimation of the volume of foods and drinks consumed (Chae *et al.*, 2011; Khanna *et al.*, 2010). A diagram of the mFR–TADA system indicating each step involved in capturing an image is shown in Figure 2. The image analysis is linked directly to food composition databases for calculating the energy and nutrient content of the foods and drinks consumed. A similar approach is the food record mobile application (FRapp) (Casperson *et al.*, 2015). To date, these

technologies have only been tested in terms of the usability of the mobile application among adolescents, and their willingness to do so (Boushey *et al.*, 2015; Caspersen *et al.*, 2015). The use of mobile technologies as a stand-alone method of primary recording of dietary intake appears to be promising. However, methods that rely on automated image analysis can be prone to underestimation if users do not capture images of adequate quality before and after intake of food and drink, and on account of missed meal images which cannot be recorded retrospectively (Casperson *et al.*, 2015).

Figure 2 - Diagram of the Technology Assisted Dietary Assessment (TADA) system that starts with capturing an image with the mobile food record (mFR)



(1) A user captures an image of an eating occasion; the image is sent to a server. (2) The image is analysed to identify the foods and drinks. (3) The labelled image is returned to the user for the 'review process', as shown by the dotted green line. (4) The user confirms the automatic labels or corrects the labels. (5) The image is returned to the server for final identification and volume estimation. (6) Identified foods and amounts are matched for nutrient analysis to the Food and Nutrient Database for Dietary Studies. (7) Images and data are stored in a server for use by researchers or clinicians. Source: Boushey *et al.* (2015), reproduced.

2.3.4 INTERACTIVE COMPUTER AND WEB-BASED TECHNOLOGIES

Interactive computer and web-based technologies involve the use of interactive dietary assessment programs installed on a desktop or portable computer; the 'web-based' moniker refers to these devices being connected to the Internet. The aim of these technologies is to collect dietary data during a specific period in the recent or distant past (short- or long-term dietary assessment). The methods associated with these technologies are based on pen and paper traditional methods which are introduced into a computer program together with a series of multimedia attributes, i.e. colours, food photographs, audio narration, animated guides, graphics and/or touch screens, pop-up functionalities and webcams. Several authors have developed interactive computer-based dietary assessment methods from pen and paper traditional methods such as FFQ (Wong *et al.*, 2008), 24-hour recall (Kirkpatrick *et al.*, 2014; Zoellner *et al.*, 2005), food records (Timon *et al.*, 2015) or dietary history (Beasley *et al.*, 2009). For data collection, participants are asked to report their food intake during a specific period, and add this information to the previously-developed computer software. The software can also include a comprehensive system for probing forgotten items. Once the food items are introduced and coded, the system calculates the intakes by means of multimedia features. Web-based technologies also include several software components, e.g. adjustable images of portion sizes. An advantage of web-based technologies is that they allow data collection to take place at any time – i.e. real-time data collection and analysis – and at a location and in a language that are more convenient for the participants (Holm *et al.*, 2015; Illner *et al.*, 2012). Nevertheless, to handle an interactive computer and/or web-based technology, the users may need to possess a high level of literacy and computer skills (Illner *et al.*, 2012).

As an example, an interactive computer-based method called Novel Assessment of Nutrition and Ageing (NANA) was developed to assess the dietary intake of the elderly. NANA consists of a touch-screen computer-based food record, plus a fixed webcam for capturing any foods and drinks participants consume, in real time. The touch-screen button selections allow participants to navigate and select food items from 12 high-level food groups. The subsequent determination of portion size is carried out by nutritionists rather than participants (Timon *et al.*, 2015). Another example is the web-based Automated Self-Administered 24-hour Recall (ASA24), which consists of a respondents' website used to collect data either in English or Spanish, and a researchers' website used to manage logistics and undertake data analysis. The ASA24 system presents a series of audio-visual aids to help respondents complete the 24-hour recall, and has shown results comparable with an interviewer-administered 24-hour recall (Kirkpatrick *et al.*, 2014). However, web-based technologies such as ASA24 require high-speed Internet access for optimum performance, as well as a high level of literacy in the respondents. These requirements may limit their viability in low resource settings.

2.3.5 SCAN- AND SENSOR-BASED TECHNOLOGIES

Scan-based technologies allow participants to scan the barcodes of purchased food items; therefore, the applications of this technology is restricted to commercial or institutional settings (Illner *et al.*, 2012). A more innovative approach is sensor-based technologies, which are designed to be memory-independent and almost entirely passive for the participants. This approach also minimizes intrusion to avoid alteration of participants' eating habits. A sensor-based technology named eButton consists of a small electronic device to record food intake automatically. It contains a miniature camera, a

microphone, a memory card and several other sensors. Participants are requested to wear the device around their neck during eating occasions. The device collects visual data immediately in front of the participants, storing the data on its memory card. The data are constantly transferred to the dietitians' computers for data analysis (Sun *et al.*, 2010).

Estimation of portion size takes one of two approaches. First, servings of food with different portion sizes consumed in the participant's home can be estimated via automated image analysis using fiducial markers, such as a marked tablecloth and/or reference pictures and known dimensions of plates; these measurements are made by the participant prior to the study. Second, where food intake occurs outside the home, the device emits lights – small beams produced by laser diodes – to project a dimensional referent into the visual field that allows for calculation of portion sizes (Sun *et al.*, 2010). Once the food and portion sizes are determined, the system is linked to a food database for calculation of nutrient content.

Counting of chews and swallows has also recently been incorporated into sensor-based devices. These are a precise measure of biological movements related to eating activities, and as such are a new method for detecting and quantifying food intake (Fontana *et al.*, 2015; Sun *et al.*, 2015). These innovative technologies suggest a more objective dietary assessment. They may offer a promising alternative to overcome some of the limitations of traditional self-report methods. However, these are still emerging technologies that need to be adapted and validated before they can be deemed both feasible and suitable in different settings and populations.

2.3.6 APPLICATIONS AND USES OF INNOVATIVE TECHNOLOGIES TO IMPROVE DIETARY ASSESSMENT

Innovative technologies have been shown to be useful in assessing current dietary intakes, as most of them are simple and non-invasive. The majority of reported innovative dietary assessment methods or technology-assisted dietary assessment methods have been used to quantify participants' food intake behaviour in a variety of controlled settings such as schools, colleges and university cafeterias (Wang *et al.*, 2006; Williamson *et al.*, 2003), or in hospitals and community centres (Kikunaga *et al.*, 2007; Ptomey *et al.*, 2015; Rollo *et al.*, 2011). Fewer studies have been carried out in free-living conditions (Gemming *et al.*, 2013; Lassen *et al.*, 2010; Martin *et al.*, 2009; Martin *et al.*, 2012) and in low income populations (Fowles *et al.*, 2008; Lazarte *et al.*, 2012).

PDAs have been widely used in industrialized countries and controlled settings (Forster *et al.*, 2015). However, their application is still limited to low resource settings. PDA food records require high literacy in the respondents; as such, children and older or less literate populations might have difficulties in using a PDA for recording food intake (Fowles *et al.*, 2008; Ortega *et al.*, 2015).

Image-assisted methods were used to assess dietary intake in children (11 months to 8 years) from remote communities in Australia (Liberato *et al.*, 2015), in adolescents (12–17 years) with intellectual disabilities (Ptomey *et al.*, 2015), and in overweight and obese adults (Martin *et al.*, 2012). Mobile-based technologies were used to address difficulties reported by adolescents (11–15 years) on conducting dietary assessment (Boushey *et al.*, 2015; Casperson *et al.*, 2015) and for recording dietary intake in adults with type 2 diabetes (Rollo *et al.*, 2011). The results of these studies have shown that image-facilitated

dietary assessments can be suitable in different populations and settings. One study (Lazarte *et al.*, 2012) showed the feasibility of using images in the rural areas of a low resource country, where the characteristics of the population (i.e. low level of literacy) and the resources they possessed (i.e. lack of Internet connection) were taken into consideration for the development of the method.

The interactive computer-based food record NANA has been used to assess the dietary intake of older adults (Timon *et al.*, 2015). Meanwhile a web-based FFQ has been used to evaluate eating practices and diet quality in a large sample size of 7 531 people from Norway, Denmark, Sweden and Finland (Holm *et al.*, 2015). In all cases, the use of PDAs, mobile applications, interactive computer and web-based technologies has been shown to require an adequate level of literacy and technical skills on the part of the participants. These requirements might limit their usage in some low literacy populations. Adequate training can facilitate the use of some technologies, especially in selected population groups such as adolescents and adults in some low-resource settings, who may already have access to mobile phones and computers. Sensor-based technologies may be easier to administer, but these are still limited in both industrialized and low resource settings. As mentioned earlier, while some of these technologies are promising, they are also emergent and at a developmental phase.

2.3.7 VALIDATION OF INNOVATIVE TECHNOLOGIES TO IMPROVE DIETARY ASSESSMENT

Validation studies have been carried out comparing results from innovative dietary assessment methods with results from traditional dietary assessment methods. Comparisons undertaken have included weighed food records (Kikunaga *et al.*, 2007; Lassen *et al.*, 2010; Lazarte *et al.*, 2012; Schap *et al.*, 2014; Wang *et al.*, 2006), estimated food records (McClung *et al.*, 2009;

Yon *et al.*, 2006), 24-hour recalls (Gemming *et al.*, 2013) and a few comparisons with more objective methods such as DLW (Gemming *et al.*, 2015; Martin *et al.*, 2012).

Gemming and colleagues evaluated whether 24-hour recall using the wearable camera SenseCam could reduce under-reporting of energy intake by providing visual aids (Gemming *et al.*, 2013). Their findings showed that images increased self-reported energy intake by approximately 12.5 percent ($P=0.02$) compared with the 24-hour recall alone. The improvement was mainly explained by the reporting of forgotten food items and the adjustment of some portion sizes. The validity of SenseCam-assisted 24-hour recall was also assessed against the DLW technique (Gemming *et al.*, 2015). Energy intake was assessed in free-living conditions by three multiple-pass 24-hour recalls alone, and by SenseCam-assisted 24-hour recall. The results compared total energy expenditure (TEE, from DLW) with energy intake calculated from the multiple-pass 24-hour recalls alone and from the SenseCam-assisted 24-hour recall. They found that underestimation of energy intake was significantly reduced ($P<0.02$), in the range of 6 to 8 percent, when the wearable camera assisted the 24-hour recall. The authors concluded that wearable cameras enhanced the accuracy of self-report methods such as 24-hour recalls.

Image-assisted methods have also been validated against the weighed food record – the so-called ‘gold-standard’ – in free-living conditions in industrialized countries (Kikunaga *et al.*, 2007; Lassen *et al.*, 2010) as well as by one study in free-living conditions in low resource settings (Lazarte *et al.*, 2012). When the results were compared with the reference method, small differences were shown (Kikunaga *et al.*, 2007; Lassen *et al.*, 2010; Lazarte *et al.*, 2012). Lazarte and colleagues (2012) developed a food photography 24-hour recall (FP 24-hR) to evaluate food consumption among women in rural areas in Bolivia. When FP 24-hR was validated against the weighed food

record, the authors found small differences in nutrient intakes: from 0.90 percent for vitamin C to 5.98 percent for fat ($P < 0.05$). Furthermore, Lassen and colleagues (2010) have developed an image-based food record. Their method was validated against weighed food records of evening meals, with a negative difference of 11.3 percent for energy intake ($P < 0.001$) (Lassen *et al.*, 2010). In these studies, high correlations between the image-assisted methods and weighed food records were reported (Lassen *et al.*, 2010; Lazarte *et al.*, 2012). However, high correlations did not necessarily imply good agreement between the methods; a better approach was to examine the Bland–Altman plots. In this analysis, acceptable limits of agreement between the methods were found for both energy and nutrient intakes. This indicated that differences were random without systematic bias (Lassen *et al.*, 2010; Lazarte *et al.*, 2012). It was suggested that by improving the quality of the photos it might be possible to reduce the differences. Image-assisted dietary assessment methods are promising to enhance the accuracy of some traditional self-reported dietary assessment methods, although there is still room for improvement in accuracy.

A PDA technology named MiHealthLog (McClung *et al.*, 2009) was compared with traditional written food records and validated against DLW. The authors found that energy intake estimated by food records (difference of 3 percent) and PDA (negative difference of 8 percent) were similar to energy expenditure calculated by DLW ($P > 0.05$), concluding that estimation of energy intake is similarly biased for both PDA and food records (McClung *et al.*, 2009). Another PDA named DietMatePro (Beasley *et al.*, 2009) was compared with food records and validated against DLW. Beasley and colleagues reported comparable correlation between DietMatePro ($r = 0.41$ to 0.71) and written food records ($r = 0.63$ to 0.83). It was concluded that DietMatePro may improve diet adherence compared with participants adhering to a written food record ($P = 0.039$). However, it does

not appear to produce more valid data than the pen and paper approach (Beasley *et al.*, 2009).

The computer-based method NANA (Timon *et al.*, 2015) was validated against written food records and against biomarkers of nutrient intake. Good correlations were found between these dietary assessment methods for energy ($r = 0.88$) and macronutrient intakes ($r = 0.80$). The authors also found positive correlations between urinary urea and dietary protein using either written food records ($r = 0.56$) or NANA ($r = 0.47$). They concluded that NANA compared well with written food records and offered a potential alternative for estimating dietary intake in community-living older adults (Timon *et al.*, 2015). Nevertheless, the different technologies and alternatives available in the literature need to be adjusted according to the characteristics of each study area and validated before their actual application.

2.3.8 STRENGTHS AND LIMITATIONS OF INNOVATIVE TECHNOLOGIES TO IMPROVE DIETARY ASSESSMENT METHODS

The strengths and limitations of innovative technologies to improve dietary assessment methods are presented in Table 9, based on information extracted from the latest reviews published on this topic (Gemming *et al.*, 2015; Illner *et al.*, 2012; Martin *et al.*, 2014; Ngo *et al.*, 2009; Stumbo, 2013).

Table 9 - Strengths and limitations of innovative technologies to improve dietary assessment methods

METHOD	STRENGTHS	LIMITATIONS
All innovative technologies used in dietary assessment	<p>Costs for data collection can be lower (less need for person-to-person interaction)</p> <p>Convenient for users, good acceptability, may improve compliance</p> <p>Do not rely on respondent's memory</p> <p>Record of qualitative information (e.g. date and time of recording)</p> <p>Significantly cut down data processing time</p>	<p>Larger up-front investments (i.e. purchase of mobile phones, cameras, computers, software development, etc.)</p> <p>Risk of losing devices</p> <p>Risk of technical problems (i.e. low battery, loss of Internet connection) could impede data collection</p> <p>A backup method is required to collect information, if technical problems occur</p>
Personal Digital Assistant (PDA)	<p>Is portable and can be easily carried by study participants</p> <p>Facilitates real-time data collection, entry and coding</p> <p>It is possible to set an alarm within the PDA to alert participants to record their food intake</p> <p>Can be programmed to allow participants strict access to the dietary intake software only</p>	<p>Face-to-face training of the participants is required</p> <p>Low level of dietary data details because of pre-coded food listings</p> <p>Increases the respondent burden compared with pen and paper records, due to the extensive list of foods (depending on each PDA)</p> <p>Reports of difficulty using the search function and inability to find certain foods</p>
Image-assisted methods, i.e. digital photographs	<p>Easy to use</p> <p>Suitable for low literacy populations (if the technologies are easy-to-use, e.g. digital cameras)</p> <p>Quality of digital cameras keeps improving and pictures with higher resolutions can help improve the accuracy of analysis</p> <p>Suitable for subjects with memory impairments and for children</p>	<p>Participants may forget to take some images</p> <p>Not all information can be captured with a single photograph/image</p> <p>Difficulties in estimating portion size of food consumed from common mixed dishes</p> <p>Lack of details about cooking methods</p>

METHOD	STRENGTHS	LIMITATIONS
	<p>Lower under-reporting compared with some traditional methods</p>	<p>Probably more limited accuracy for countries with a wide range of mixed dishes (e.g. Asia)</p> <p>Still needs a written record of foods obscured in photos, and when details of ingredients are required</p>
<p>Image-assisted methods, i.e. mobile-based technologies</p>	<p>Possible higher quality control of data because of shorter delays and real-time responses</p> <p>Possibility of sending reminders</p> <p>Internet access would allow respondents to send instant photos, thereby minimizing systematic mistakes</p>	<p>It is costly and time-consuming to develop the application as an interface, and the software for automated portion size estimation</p> <p>Certain types of foods, such as mixed dishes (soups, stews, casseroles, etc.) can be difficult to analyse with automated image analysis</p> <p>Requires certain level of literacy</p> <p>Requires network/Internet access (for real-time data collection)</p>
<p>Interactive computer and web-based technologies</p>	<p>Efficiency in terms of recording information and data processing (i.e. reduced costs and interviewer workload)</p> <p>Increased levels of quality control</p> <p>Include interactive visual and audible aids</p> <p>Suitable for large, geographically dispersed samples; can include different countries/languages (web-based)</p> <p>Data processing can be finalized at any time and location (web-based)</p> <p>Can provide personalized dietary feedback (web-based)</p> <p>Possibility of sending interactive reminders (web-based)</p>	<p>Some imaging algorithms are not sufficiently advanced to identify foods correctly and to accurately estimate the quantity of food in the computerized images</p> <p>Need adaptation of the software to local settings</p> <p>Require high levels of literacy and computer skills from the participants</p> <p>Possibility of collecting less food details (i.e. ingredients, methods of preparation, etc.)</p> <p>Require Internet access (web-based)</p>

METHOD	STRENGTHS	LIMITATIONS
Scan- and sensor-based technologies	<p>Incorporation of barcode scanning and PLU (Price Look Up) codes allows automatic identification of foods</p> <p>Objective dietary assessment (sensor-based)</p> <p>Passive for the participants, reduces burden (sensor-based)</p>	<p>Scanning barcodes is applicable to packed food only</p> <p>Have only been used in controlled settings to date</p> <p>Their application to date has been restricted to small test studies</p>

The applications of innovative technologies in dietary assessment are subject to constraints such as cost, location, available resources, literacy level of the population and number of interviews. Therefore, some of these technologies remain relatively costly for implementation in large-scale epidemiological studies. As mentioned earlier, innovative technologies may be used as stand-alone dietary assessment methods, or can be added as an aid to improve data collection in conventional dietary assessment (i.e. 24-hour recall, dietary record and FFQ). Using innovative

technologies along with a conventional dietary assessment may reduce the up-front cost of the stand-alone innovative methods and avoid some technical drawbacks, i.e. problems with data transfer and storage, battery life, Internet access, etc. For the dietary assessment methods used in epidemiological studies, Shim and colleagues reviewed the strengths and limitations of innovative technologies incorporated into conventional dietary assessment methods, found in Table 10 below (Shim *et al.*, 2014).

Table 10 - Strengths and limitations of innovative technologies integrated into conventional dietary assessment methods

	24-HOUR RECALL	DIETARY RECORD	FOOD FREQUENCY QUESTIONNAIRE
Required technology	Software, Internet, mobile phone, digital camera, etc.	Software, Internet, PDA, mobile phone, application, etc.	Skip algorithms, questions that ask for multiple details, pictures of foods, etc.
Strengths	Standardized data collection is possible (reducing interviewer bias); probable reduced time and cost; improves feasibility	Standardized, real-time data collection is possible; probable reduced time and cost; improves feasibility	Able to collect complex information and highly accurate data
Limitations	Inherent bias related to self-reporting	Inherent bias related to self-reporting; requires participant training on how to use the technology	Measurement errors related to methodology remain

Adapted from Shim, Oh and Kim (2014) with permission.

The limitations discussed in Table 10 may restrict the usage of some innovative technologies (i.e. computer web-based methods, scan- or sensor-based methods) for dietary assessment in rural areas, in low resources settings and among low literacy populations. These limitations might be overcome with the addition of new and evolving

technologies and better ways to control bias. Currently, image-assisted methods (i.e. digital cameras, mobile phones) may be the most suitable for low-resource settings, and could have a positive impact on dietary assessment in these settings.

BOX 8. QUICK GUIDE TO USING INNOVATIVE TECHNOLOGIES FOR DIETARY ASSESSMENT

To integrate innovative technologies in supporting conventional methods (dietary records and 24-hour recalls)

Define the project objectives, the design of the study, the sample size and the budget

Take into consideration the specific characteristics of the population and settings: e.g. literacy level, computer skills, eating from a common pot, highly mixed dishes, availability of electricity and Internet connectivity

Choose the technologies that will be used for dietary assessment, e.g. digital cameras, mobile phones, tablets, computers, mobile applications, computer software, etc

Select the method for estimating portion sizes

- ✓ Only digital images;
- ✓ Photo atlas with standardized food portion sizes (printed or digital) for comparison with the digital images;
- ✓ Automatic analysis of food identification and portion size estimation, e.g. FRapp (Food Record application), ACTi Pal (www.actipal.com), mFR app, TADA system (www.tadaproject.org).

Define the number of days to be recorded: three days or more is recommended when the aim is to evaluate the individual's usual diet

Define the time frame, and days selected for conducting the method (e.g. weekdays vs. weekend days, consecutive vs. non-consecutive). It is recommended to include one weekend day

Compile information on composite dishes (e.g. soups, stews) and/or dietary supplements (e.g. vitamins, minerals). This information is difficult to extract from images, so the following alternatives are suggested

- ✓ *In case of image-assisted record methods*, the respondent should keep a record with the description of the composite dishes, including lists of ingredients and quantities, given that these may be difficult to identify in the images.

- ✓ *In case of image-assisted recall methods*, the interviewer should ask for a description of the composite dishes, including lists of ingredients and quantities.

Establish the procedure to follow in case of obscured or missing images of food events

- ✓ Train the respondent to review the images after taking them and to take a second image if necessary.
- ✓ Ask the respondent to write down a description of the consumed foods if the images are not clear or if he/she forgot to record the images.
- ✓ Innovative technologies assisting 24-hour recall allow the interviewer to ask for information necessary to clarify obscured images and to obtain detailed information on food ingredients.

Capacity development for the respondents

- ✓ Training in the appropriate use of digital cameras, smartphones, mobile applications and/or other devices used to collect dietary information.
- ✓ Computer skills (required for computer-based and web-based methods).

For innovative technologies used as stand-alone methods, keep in mind the following aspects

Be aware that images, if used as the only source of information, could not describe food ingredients in composite dishes. Thus, the method may increase the likelihood of under-reporting.

- ✓ Additional voice or writing records would be needed in case of obscured images.

Stand-alone innovative methods may not be suitable to estimate portion size for certain populations, for example

- ✓ When the family eats from a common pot;
- ✓ When the majority of the meals are mixed dishes such as soups and stews, or have sauces served on top of the other ingredients.

2.4 QUALITATIVE RETROSPECTIVE PROXY TOOLS FOR ASSESSING DIETARY DIVERSITY

The Dietary Diversity Score (DDS) is a proxy tool based on the concept that 'dietary diversity is a key element of diet quality and a varied diet helps ensure adequate intakes of essential nutrients that promote health' (Ruel *et al.*, 2013). DDS is easy to use, inexpensive, practical, and can be

utilized to quickly obtain results. Hence, it can be used for rapid assessment of dietary diversity in selected population groups, in contexts where resource and capacity constraints might restrict the use of more detailed dietary assessment methods. DDS provides a simple score which represents a number of different foods and/or food groups consumed over a given reference period (Hoddinott *et al.*, 2002). There are various DDSs published and used for different purposes. They differ in the number and definition of food

groups or food items, reference periods and units of analysis (i.e. household or individual level), and are as follows:

- *At household level:*
 - Household Dietary Diversity Score (HDDS) (FAO, 2010; Swindale *et al.*, 2006) is based on the count of 12-food groups. The information is extracted from one qualitative 24-hour recall, excluding foods consumed outside the home. It accesses the quality of food access at household level. Additionally, it is useful for identifying emergencies related to food security and malnutrition (Ruel, 2003a). Studies have shown that an increase in dietary diversity at household level was positively associated with household food security (Hoddinott *et al.*, 2002).
 - Food Consumption Score (FCS) (WFP, 2008) is based on the count of 9-food groups. The score is calculated taking into account the frequency of consumption of different food groups over a period of 7 days, before the survey, and the relative nutritional importance of the food groups. The results of the analysis categorize each household as having either poor, borderline or acceptable food consumption. It aims to reflect the quality and quantity of food access at household level.
- *At individual level:*
 - To assess dietary diversity and micronutrient adequacy of the diet at individual level, many indicators have been developed varying in the number of food groups, target population and cut-off point (FAO/FANTA/IRD, 2014). On account of their importance, the indicators Minimum Dietary Diversity – Women (MDD-W) and Infant and Young Child Dietary Diversity Score (IYCD) (WHO/UNICEF, 2010) are presented in further detail below.

2.4.1 MINIMUM DIETARY DIVERSITY –WOMEN (MDD-W)

The MDD-W indicator (Martin-Prével *et al.*, 2015, 2017) aims to be used as a global indicator of dietary diversity, and was recently developed from the Women's Dietary Diversity Score (WDDS) (FAO, 2010). WDDS (FAO, 2010) was constructed with the simple count of a 9-food group indicator, but it did not have an established cut-off value. There is an increasing need for a dichotomous indicator that is useful for target-setting purposes and advocacy (FAO, 2015b). As a result, WDDS was updated to the MDD-W indicator. MDD-W has been defined as a dichotomous indicator of whether or not women who are 15–49 years of age have consumed at least five out of ten defined food groups during the previous day and night. The proportion of women of 15–49 years of age who reach this minimum dietary diversity (five or more food groups) in a population can be used as a proxy indicator for micronutrient adequacy, which is one important dimension of diet quality (FAO/FHI360, 2016; Martin-Prével *et al.*, 2015). MDD-W was developed through secondary analysis of dietary intake (from multiple 24-hour recalls) of nine data sets from rural and urban areas in six countries: Bangladesh, Burkina Faso, Mali, the Philippines, Mozambique and Uganda (Martin-Prével *et al.*, 2015). The nine datasets were used to calculate the mean probability of micronutrient adequacy (MPA) for 11 micronutrients – vitamin A, thiamine, riboflavin, niacin, vitamin B6, folate, vitamin B12, vitamin C, calcium, iron and zinc – using the probability of adequacy approach. This approach is based on information or assumptions about both the distribution of nutrient requirements in the population and the day-to-day variations (intra-individual) in nutrient intakes (Barr *et al.*, 2002). The 10-food group indicator and the cut-off of the MDD-W were chosen by examining the sensitivity, specificity and total misclassification of the MDD-W with several cut-offs and thresholds of MPA (i.e. 0.50, 0.60, and 0.70). Each one of the

10-food group indicator holds the same weight. The food groups are as follows:

1. all starchy staples
2. beans and peas
3. nuts and seeds
4. all dairy
5. flesh foods
6. eggs
7. vitamin A-rich dark green leafy vegetables
8. other vitamin A-rich vegetables and fruits
9. other vegetables
10. other fruits

A cut-off point of five was established as the ‘best cut off’ since it performed accurately in a sufficient number of data sets (six out of nine data sets) and at different MPA levels. The cut-off of five was, furthermore, recommended as a reasonable predictor of an MPA >0.60 (for more details on the MDD-W cut off selection process, refer to Martin-Prével *et al.*, 2015). The performance of food group indicators was evaluated through a sensitivity–specificity analysis, examining error rates of classification across a range of cut-offs and for various MPA cut-offs. Low sensitivity was found in cases of very varied diets containing small quantities of nutrient-rich foods, e.g. diets in Asia. In addition to the low sensitivity, the authors reported that results from the secondary analysis were consistent enough that MDD-W could be recommended for global use in population-level assessment and advocacy (Martin-Prével *et al.*, 2015, 2017).

It is generally agreed that dietary diversity is a key element of diet quality. Thus, dietary diversity should be measured as reliably as possible. MDD-W data is relatively simple to collect and the score is easy to calculate. Food consumption of individuals during the preceding 24 hours is recalled using a qualitative 24-hour recall questionnaire, including foods that have been eaten outside the home. Alternatively,

the information can be collected with a list-based questionnaire (a form of FFQ), where the interviewer asks the respondent if she has eaten foods from different food groups, using a culturally-adapted food list (i.e. common food names, local foods). The information on food consumption, collected either by 24-hour recall or list-based questionnaire, that matches the items in the 16-food group list are underlined. To create the MDD-W score, the preliminary list of the 16-food group is aggregated into the 10-food group indicator (Appendix 6), where one point is allocated whenever a food item contained in the 10-food groups has been underlined in the 16-food group list. Even if each food group has one or more underlined food item only one point is allocated, thus contributing equally to the final score. The value of zero is given if none of the food items in the food group were consumed. The sum of the consumed food groups represents the total DDS.

Only food items that were consumed in a quantity greater than 15 grams (roughly a tablespoon) should be considered and included in the 10-food group indicators. The minimum quantity rule of 15 grams was defined by secondary analysis where a comparison was made between two indicators: the first, imposing a 1 gram rule and the second, imposing a 15 grams rule (Arimond *et al.*, 2010). A minimum consumption of 15 grams of a food group per day is required in order for a food group to be accounted for in the score and, therefore, helping to provide a better indication of MPA.

2.4.2 INFANT AND YOUNG CHILD DIETARY DIVERSITY SCORE (IYCDDS)

IYCDDS is defined as the “Proportion of children 6–23 months of age who receive foods from four or more food groups” (WHO/UNICEF, 2010). This means that children who received four or more food groups out of seven during the previous day were more likely to have an adequate diet than

children who consumed food items from less than four food groups. The indicator is based on the count of 7-food groups. It was developed for children of 6–23 months old through secondary analysis of ten data sets from a multi-country study that included countries in Africa, Asia and Latin America. The 7-food groups were selected by examining the sensitivity and specificity of the indicators performance with a different number of food groups. The indicator was further investigated using a 1 gram and 10 gram minimum consumption cut-off, and concluded that a 1 gram cut-off per day provided a better result. Therefore, since the cut-off is so low the consumption of any amount of food from each food group is sufficient enough to be accounted for in the construction of the score, except if an item is only used as a condiment (WHO/UNICEF, 2010). The food group classification includes:

1. Grains, roots and tubers
2. Legumes and nuts
3. Dairy products (milk, yogurt, cheese)
4. Flesh foods (meat, fish, poultry and liver/organ meats)
5. Eggs
6. Vitamin-A rich fruits and vegetables
7. Other fruits and vegetables

The cut-off of at least four of the above 7-food groups indicates a minimum dietary diversity in infants and young children. The cut-off was selected because it was shown to be associated with better quality diets for both breastfed and non-breastfed children (WHO *et al.*, 2008). Briefly (in a similar approach as that explained in section 2.4.1), the selection of the cut-off was based on data analysis from the multi-country study, where the association between IYCDs and mean micronutrient density adequacy (MMDA) of the diet for nine and ten nutrients was tested (WHO *et al.*, 2008; WHO/UNICEF, 2010). The indicators were positively and significantly associated with MMDA at all ages in all countries and for both

breastfed and non-breastfed children. Although sensitivity and specificity analyses failed to identify a cut-off point that performed best across all contexts, the cut-off point of four was selected based on extensive stakeholder consultations and discussions (Leroy *et al.*, 2015). The consultation agreed that the consumption of foods from at least four food groups on the previous day would mean that in most populations the child had a high likelihood of consuming at least one animal-source food and at least one fruit or vegetable that day, in addition to a staple food (grain, root or tuber) (WHO/UNICEF, 2010).

To construct the IYCDs score, data on child food consumption are collected through an interview with the parents or caregivers. The interviewer asks about the different types of foods the child ate the day before the interview, using a 24-hour recall approach. The food items recalled by the respondent are then underlined and given a value of 1 in one of the 7-food groups for the construction of the IYCDs score (WHO/UNICEF, 2010). Results may be reported separately for breastfed and non-breastfed children. However, diversity scores for breastfed and non-breastfed children should not be directly compared, because breast milk is not counted for in any of the above food groups, since the indicator is meant to reflect the quality of the complementary diet. It is recommended that the indicator be further disaggregated and reported for the following age groups: 6–11 months, 12–17 months and 18–23 months (WHO, 2008). Detailed steps for IYCDs data collection are presented in the guide published by WHO (WHO, 2010).

2.4.3 APPLICATIONS AND USES OF INDIVIDUAL LEVEL DIETARY DIVERSITY SCORE

DDS at the individual level has been used to evaluate the probability of having a varied diet, and has also been associated with the probability of micronutrient adequacy in the diet (Arimond *et*

al., 2010; Ruel, 2003a). In this regard, Kennedy and colleagues found that DDS was associated with the adequacy of 11 micronutrients – vitamin A, vitamin C, vitamin B6, vitamin B12, thiamine, riboflavin, niacin, folate, calcium, zinc and iron – in the diets of women of reproductive age in an urban area of Mali (Kennedy *et al.*, 2010). DDS was also shown to be a useful indicator of micronutrient adequacy in children (24–71 months old) in the Philippines (Kennedy *et al.*, 2007).

Since the tool is simple and inexpensive to use, studies can be repeated several times in 1 year, helping to capture the seasonal effect. Savy and colleagues have used DDS to study the seasonal variations of women’s diets in a rural area of Burkina Faso. The authors assessed the DDS of ten standard food groups at the beginning and at the end of the seasonal cereal shortage. They concluded that DDS could help to identify vulnerable individuals from a socio-economic and nutritional point of view when DDS was measured at the end of the shortage season (Savy *et al.*, 2006). Additionally, DDS has been used to evaluate variations of dietary intakes between populations in different settlements or regions in South Africa (Acham *et al.*, 2012). It has also been used to investigate the associations between environmental, physiological and sociodemographic variables and dietary diversity in Bolivia, Botswana and Burkina Faso (Benefice *et al.*, 2006; Clausen *et al.*, 2005; Savy *et al.*, 2006). Furthermore, the dichotomous MDD-W indicator has been used for the first time in Tajikistan to collect baseline nutrition information across four regions of the country. The study was useful to identify the percentage of women that consumed five or more food groups, and were classified as being more likely to have an adequate diet and meet their nutritional requirements. Regions in Tajikistan with less varied diet diversity, along with the food groups that were consumed less by Tajik women, were identified in the study (Lazarte *et al.*, 2015). As a result of such studies, DDS has been shown to be a potentially low-cost,

field-friendly tool for rapid assessment of dietary diversity, an important feature of dietary quality and micronutrient adequacy.

2.4.4 VALIDITY OF INDIVIDUAL LEVEL DIETARY DIVERSITY SCORE

Several authors have constructed and validated DDS by secondary data analysis (Arimond *et al.*, 2004; Arimond *et al.*, 2010; Kennedy *et al.*, 2010; Martin-Prével *et al.*, 2015). This involves the comparison of results from DDS with existing results from individual dietary assessments methods (i.e. quantitative 24-hour recall), anthropometric measurements and/or biochemical markers. One major limitation of DDS studies validated through secondary data analysis is that the MPA and the food group indicators were generated from the same data sets; as such, measurement errors may be correlated and this could bias the results of associations (Arimond *et al.*, 2010). With secondary data analysis, DDS has been shown to associate with the MPA for micronutrients calculated from previous data of quantitative 24-hour recalls (Arimond *et al.*, 2010; Kennedy *et al.*, 2010; FAO, 2015b). Additionally, DDS has been compared with the results of energy intake, nutrient adequacy and anthropometric measurements calculated from previously collected dietary data in adults (Arimond *et al.*, 2010; Hoddinott *et al.*, 2002) and children (Arimond *et al.*, 2004; Steyn *et al.*, 2006). In one study, HDDS (a 12-food group indicator) was presented as a proxy indicator of household food access and was associated with energy availability calculated from data sets in ten countries: India, the Philippines, Mozambique, Mexico, Bangladesh, Egypt, Mali, Malawi, Ghana, and Kenya. The authors found that a 1 percent increase in dietary diversity was associated with a 0.7 percent increase in total per capita caloric availability (Hoddinott *et al.*, 2002). Meanwhile WDDS, a 9-food groups indicator (Arimond *et al.*, 2010) was presented as a proxy indicator for

micronutrient adequacy in women of reproductive age. WDDS was associated with the MPA for 11 micronutrients – vitamin A, vitamin C, vitamin B6, vitamin B12, thiamine, riboflavin, niacin, folate, calcium, zinc and iron. Where MPA was calculated from data sets of quantitative 24-hour recalls from five low resource settings (Burkina Faso, Mali, Mozambique, Bangladesh and the Philippines), the authors reported that MPA for micronutrients in all sites were correlated with all 9-food groups of DDS. Associations were stronger for WDDS when a 15 gram minimum consumption cut-off was required for a food item to be included in the food groups for the final score (Arimond *et al.*, 2010). A similar approach was used to assess the associations between MDD-W and MPA (Martin-Prével *et al.*, 2015). Additionally, Arimond *et al.* (2004) reported the association between DDS (7-food group indicator) and height-for-age Z-scores (HAZ) in children 6 to 23 months old. They extracted and analysed data from the demographic and health surveys in 11 countries: Benin, Cambodia, Nepal, Ethiopia, Haiti, Colombia, Peru, Malawi, Rwanda, Mali and Zimbabwe (Arimond *et al.*, 2004). The data were adjusted for a number of potentially confounding factors, i.e. age, sex, breastfeeding status, mother’s education, household wealth/welfare residence area, etc. The authors reported that in 10 out of the 11 countries, significant associations were found between DDS (7-food group indicator) and HAZ, either as a main effect or as an interaction with other factors, e.g. age, breastfeeding, urban or rural living area (Arimond *et al.*, 2004).

Torheim and colleagues validated a DDS (10-food group indicator) by comparing the scores with nutrient adequacy assessed via a 2-day weighed food record collected by two studies in western Mali. They found acceptable correlations ($r=0.25$, $P<0.05$) between DDS and nutrient adequacy estimated by the weighed food records in one of the studies (Torheim *et al.*, 2003). Furthermore, if DDS aims at assessing the risk of inadequate micronutrient intake, a validation study comparing the results with biochemical indicators is recommended if resources are available. Only a few studies have assessed DDS against biomarkers. One study by Fujita *et al.* (2012), evaluated how well a 10-food group DDS predicts vitamin A status, measured as serum retinol concentration, in adult women from northern Kenya. Their findings showed that DDS was positively related to retinol concentration ($P=0.045$) and that dietary diversity had a significantly positive effect on serum retinol concentration. Results suggested that women having more diversified diets had improved intake of vitamin A and serum retinol as compared with women who had less diversified diets with equivalent vitamin A content (Fujita *et al.*, 2012).

2.4.5 STRENGTHS AND LIMITATIONS OF INDIVIDUAL LEVEL DIETARY DIVERSITY SCORES

Based on several published reviews and reports (FAO/FANTA/IRD, 2014; Herforth, 2015; Rose *et al.*, 2008; Ruel, 2003b), Table 11 shows the strengths and limitations of DDS.

Table 11 - Strengths and limitations of individual level dietary diversity score

STRENGTHS	LIMITATIONS
<ul style="list-style-type: none"> • A useful indicator for assessing dietary diversity (a key element of dietary quality) • Can potentially be used as a global indicator to classify individuals at risk of micronutrient inadequacy 	<ul style="list-style-type: none"> • Collects only qualitative dietary data • Cannot represent habitual intake of an individual person

STRENGTHS	LIMITATIONS
<ul style="list-style-type: none"> • Has a cut-off point to identify adequate diets for IYCDSS and MDD-W • Quick, simple and inexpensive data collection and analysis • Non-intrusive, low participant burden • Showed high response rate among target populations • Suitable for populations eating from common bowls (does not require detailed quantitative information of the consumed food) 	<ul style="list-style-type: none"> • Does not assess the full picture of dietary quality of subjects • Not designed to collect information on within-person variation • Relies on respondent's memory when data collection is based on 24-hour recall • Cannot assess dose-response effects via follow up intervention studies • Nutrition supplements are not taken into account • Dietary diversity may significantly vary by season. Data interpretation needs to be cautious when comparing dietary diversity data across seasons and regions

The DDS indicator is potentially suitable and feasible in low resource settings with limited resources and capacity to conduct more robust dietary assessment such as diet records or 24-hour recalls. DDS can provide valuable information on dietary diversity, which is an important factor in diet quality and micronutrient adequacy. The duration of a DDS interview is relatively short.

The tool is easy to understand and execute, and the data analysis and interpretation are straightforward. However, caution must be taken when adapting the tool to specific populations, taking into consideration context-specific characteristics (i.e. dietary habits, seasonal variations, geographical differences, etc.).

BOX 9. QUICK GUIDE TO USING INDIVIDUAL LEVEL DIETARY DIVERSITY SCORES

Define the project objectives, sample size and budget

Define the unit of analysis/data collection

✓ At individual level (e.g. MDD-W, IYCDSS)

Select the method used to collect information

✓ 24-hour recall (open free recall) or list-based method

Select the mode of data collection (e.g. pen and paper, tablets, etc.)

Define the number of days recalled

- ✓ Normally one day. DDS does not represent an individual's usual diet even if the information is collected for more than one day.

Conduct adaptation of the DDS questionnaire to population characteristics (REF, training course)

- ✓ Local adaptation of the DDS questionnaire can be conducted by meeting with local key informants and by focus group discussions.
- ✓ Understanding the food and meal patterns of the target population.
- ✓ Foods classified in the DDS questionnaire should represent those commonly consumed by the target population.
- ✓ The list of foods should be revised and modified (add or delete food items).
- ✓ The questionnaires should be translated into the local language.

Define how to proceed for mixed or composite dishes

- ✓ All the ingredients included in composite dishes should be recalled. It is recommended that the enumerators should have an understanding of local recipes to identify and record all ingredients consumed.

Define a minimum quantity rule

- ✓ A minimum quantity rule should be set and followed for foods to be counted in the construction of the score (e.g. 15 grams).
- ✓ It is recommended to validate the minimum quantity within the target population.

Consider seasonal variation: food availability can be greatly affected by seasonal variations

Define the days for conducting the method

- ✓ Any day of a habitual food intake is preferred; avoid days when the eating patterns may be significantly different from habitual (e.g. fasting days, feast days)

Consider conducting capacity development for enumerators

- ✓ Training on dietary data collection;
- ✓ How to probe the respondent using standardized and non-leading questions;
- ✓ Knowledge about the foods and food preparation methods of the study population is an asset.



©FAO/Sebastian Liste

3

METHODOLOGICAL CONSIDERATIONS

Before conducting a dietary assessment, there are numerous factors to be considered by the researchers, from culturally-specific issues to obtaining a comprehensive food composition database. These methodological challenges can be encountered throughout different stages of the research process starting from the initial planning phase to the data collection and data analysis phases. In order to reduce barriers, issues pertaining to the study purpose will need to be addressed before selecting the most suitable dietary assessment method to fit the particular study objective and setting. The aim of this chapter is to address some of the fundamental methodological considerations that have a bearing upon the assessment of diet. The information in this section is presented with an emphasis on low-resource settings. Firstly, the chapter will address sources of dietary variation, followed by individual, community and cultural specific issues, food distribution, eating habits, consumption of street foods, issues pertaining to portion size estimation and food composition tables. Secondly, measurement errors in dietary assessment are presented along with comprehensible information on validation, reproducibility and quality control.

3.1 SOURCES OF DIETARY VARIATION

Characterizing habitual diets of individuals is one of the purposes for dietary assessment studies, as many nutritional studies aim to implement interventions that are intended to affect long-term dietary behaviours and health benefits. Additionally, when estimating diet–disease associations, or examining correlates of individual intakes, the assessment of usual intake is required. Dietary intakes in many individuals can vary widely from day to day through changes in meal patterns, food availability, food habits, changing environments and cultural food factors. Therefore, it is important to address common sources of dietary variation prior to choosing the most appropriate dietary assessment method to capture the most accurate and reliable data. The following describes the main sources of dietary variations:

Day-to-day variability in individual food intakes (within- or intra-subject variability) is common when measuring dietary intake due to changes in food availability, meal patterns, household food habits, etc. This variation creates difficulties when an individual tries to determine their usual dietary

habits accurately. In order to account for individual variability, multiple day recalls and/or records are the appropriate methods to employ, which should include both weekdays and weekends. How many records or recall days are needed in order to account for the intake variability of the nutrient of interest, (e.g. some nutrients demonstrate large day-to-day variability such as vitamin A and C) as well as the degree of between- and within-individual variation. In order to obtain the most accurate data of usual intake, the number of measurement days would need to be increased (Thompson *et al.*, 2013). However, by increasing the number of measurement days this can affect recruitment level, compliance and the cost of data collection.

In a study conducted in Finland to investigate the relationship between diet and disease in middle-aged men, the authors attempted to determine the number of food record days needed in order to characterize habitual intake. The results suggested that 7–14 days are needed to accurately account for usual intake, which may be impractical for most large-scale studies (Hartman *et al.*, 1990). Thus, if resources and time are limited, more sophisticated statistical modelling (i.e. multivariate regression models) can be used to compensate for this limitation through analytical estimation and the removal of within-subject variation in the diet. Please refer to Dodd *et al.* (2006) for further details.

Between- or inter-individual variability is the measure of how individuals differ from each other in regard to their usual daily intake and is dependent on the target population and the nutrient being measured. In order to differentiate between individuals and their usual food intake, between-subject variability needs to be higher than within-subject variation. To increase this variability, the sample size should be as large as possible. Furthermore, it should be noted that for most nutrients, the level of within-subject variability is higher than between-subject

variability, making it difficult to measure the usual mean intake of an individual but easier to assess the group mean intake (Beaton *et al.*, 1997). Fukumoto *et al.* (2013) conducted a study in Japanese adults to calculate between- and inter-subject variability by age and sex in the intakes of energy, along with 31 nutrients. Subsequently, the sample size needed in order to differentiate between individual usual intakes was examined. The authors concluded that the number of days required to assess individual usual intake was higher in men, in particular for those aged 30–49 years. Therefore, depending on the objective of the study, the number of measurement days or the size of the study sample needs to be taken into account in order to influence between- or inter-individual variability.

Seasonality effects can also impact dietary variability, especially in low resource settings. In communities where people often rely heavily on subsistence farming, food availability may be moderated by the season. It is therefore important to consider, in the design stages, the possible effects of seasonal variance. Multiple recalls spread across different seasons can be used to calculate typical food intakes and avoid misinterpretations of the overall diet (Kigutha, 1997). In a study by Rao *et al.* (2009), rural Indian mothers measured the effect of seasonality in maternal intake on neonatal size. The study area was defined by having three seasons, summer, monsoon and winter. A 24-hour recall and FFQ were conducted during both the summer (lean period) and winter (harvest period). The authors concluded that maternal intakes during the harvest season had a positive association ($P=0.04$) with birth size, while the lean period presented a negative association ($P=0.002$) with birth size.

Age and sex within a population group can also cause variations in nutrient intakes, contributing to the between-subject variability. Pereira *et al.* (2010), for example, observed that the variability

of food intakes in a Brazilian population, was higher for female adolescents and male adults and estimated that 14 replications for boys and men, 15 for girls and 23 for women, were required to estimate energy intake with a precision of 90 percent. Meanwhile four replications for boys and seven for men, girls and women were required to rank individuals' energy intake (Pereira *et al.*, 2010). Furthermore, some studies have also emphasized the need to adjust within- and between-subject variability for demographic differences between subjects, because variations in intakes can result from differences in age, sex, smoking status or physical activity, as well as the season and the size of family. For example, results obtained by Stote *et al.* (2011) showed that in the United States of America, to estimate energy and macronutrient intakes during a six-month period, overweight and obese adults needed 5–10 and 12–15 days of 24-hour recalls, for men and women respectively. However, a balance between scientific rigour and practicality is often what actually occurs in practice, commonly using 4–5 days to assess individual intakes of energy and macronutrients.

Number of days of recalls or records of food intake is also a source of dietary variability. Results of a landmark study conducted in the United States of America by Basiotis *et al.* (1987) observed a large variation in the number of days of food intake records needed to predict typical nutrient intakes among individuals for the same nutrient, and within individuals for different nutrients. They reported that to assess energy intake, the fewest number of days required was 31. Meanwhile the longest period required was for vitamin A: an average of 433 days.

The day-to-day variability in low- and middle-income countries is less frequently examined than in high income countries. The popular belief that:

'diets of poor people in non-industrialized parts of the world are homogenous; therefore, day-to-day variability does not need to be taken into account

is often debatable. Variability in low- and middle-income countries is something that deserves serious consideration'

because even small economic differences might impact intakes, and subsequently increase between-person variation (Willett, 2013b). Similarly, Harrison (2004) suggests that this popular belief is not supported by evidence and intra-individual variability should be taken into account. Any assumption that the assessment of homogenous diets might require a smaller number of assessment days may not be correct. It was reported that a 6-day 24-hour recall was sufficient enough to estimate intakes of energy, carbohydrates, iron and vitamins A and C in pregnant Indonesian women, with a precision of ± 20 percent of the true intake (Persson *et al.*, 2001). Besides, other authors reported large within-person variation in the nutrient intakes of pregnant women in rural Malawi, despite limited dietary diversity. In this case, 8–23 days were required to measure energy, protein, carbohydrates and fibre intakes, and 95–213 days for micronutrient intakes, within an error range of ± 20 percent from the true intake (Nyambose *et al.*, 2002).

3.2 INDIVIDUAL, COMMUNITY AND CULTURALLY-SPECIFIC ISSUES IN LOW RESOURCE SETTINGS

Accounting for cultural aspects when assessing diet helps to ensure data accuracy and project success (Cassidy, 1994; Harrison, 2004; Kigutha, 1997). Awareness of such aspects and their influences is of utmost importance, both at individual and community level. At individual level, food preferences, taboos, specific socio-economic characteristics (such as high priced foods, preservation and transportation facilities, low socio-economic status and inability to adapt to temporary shortfalls in income) and also demographic ones (literacy and household

composition) affect food habits. It must be noted that dietary assessment in low resource rural areas poses additional methodological challenges and differs from low resource urban areas in terms of food systems, food habits, culinary practices, accessibility and food supply. Studies comparing food consumption of individuals from these two areas reported significant differences in dietary patterns and nutritional intakes, often determined by differences in dietary habits and demographic composition (Andrissi *et al.*, 2013; Omigbodun *et al.*, 2010; Oninla *et al.*, 2007). In a study conducted among Peruvian children, the authors measured the relationship between dietary pattern and growth in both urban and rural settings. The authors noted a difference in dietary pattern in the two study locations, specifically in the level of fat consumption (Andrissi *et al.*, 2013). This variation in dietary pattern highlighted the importance of considering differences associated with distinct study settings prior to conducting a dietary assessment.

Furthermore, a number of additional factors should be considered prior to conducting dietary surveys in low resource areas, including low literacy rates, accessibility to target population groups, and costs (Kigutha, 1997). Assessing the diets of people with low literacy levels also requires careful choice of an appropriate dietary assessment method. Collection of 24-hour recalls over multiple days, and food records collected by trained field workers under direct observation, are both appropriate methods. To improve compliance and quality of data collection, field workers should be aware of the cultural aspects related to the community's attitude towards face-to-face interviews, especially regarding local customs related to food behaviour and meal patterns. Food in many societies is related to social status, therefore over-reporting to impress the interviewer is not unusual (Kigutha, 1997). Similarly, when survey objectives have not been communicated clearly, it can lead to misunderstanding of the study purpose and to a belief about unjustified

benefits (i.e. food or monetary aids) for reporting low consumption (Kigutha, 1997). Furthermore, under-reporting of energy intakes has been shown to increase with an increased BMI (Livingstone *et al.*, 2003). Furthermore, studies have revealed that underreporting is associated with lower levels of education and socioeconomic status, as well as high socioeconomic status and high levels of knowledge on the subject of health and diet (Rasmussen *et al.*, 2007). Moreover, if innovative technologies such as digital cameras, mobile phones and computers are to be used for dietary data collection in low resource areas, issues such as literacy level of the respondents, availability of electricity and Internet connectivity come into play (Gemming *et al.*, 2015; Illner *et al.*, 2012).

Recruiting locally trained staff to undertake dietary assessment is strongly encouraged: such individuals would be able to build strong relationships with community leaders to increase participation rates and to make sure accurate information is communicated regarding the purpose and procedures of the assessment. He or she would be familiar with the community's food practices and cultural values, provide important cultural information in order to adapt the study protocols and tools, and to translate the dietary assessment tool and relevant materials into the local language. The time frame for data collection should be carefully considered, given that meal patterns – including the frequency and times of eating occasions – can vary by area and season. Factors to be considered include the distance of travel to collect water and wood fuel for cooking, and the fact that during the harvest season, people spend long hours in the field, and meals for the whole household may be served in one pot, with leftovers being used again on the following day, etc.

3.3 INTRA-HOUSEHOLD FOOD DISTRIBUTION, SHARED EATING OCCASIONS AND STREET FOOD

Sharing food, including eating from shared pots and plates, is common among populations in low resource settings, leading to difficulties in assessing individual food intakes. Extra efforts should be made to capture accurate records of intakes, taking into account the local cultural context and examining its influence on recorded intakes. Kigutha (1997) describes one technique to overcome the difficulties faced when measuring individual intakes from shared plates. The author suggested asking the family to serve food to an individual participating in the study on a separate plate from the rest of the family. However, this may not be an acceptable cultural practice for the family, or may cause the respondent to change their usual intake, leading to bias. In another study, investigating 45 Senegalese children, a region where eating from a common pot is the norm, the field workers calibrated handfuls of specific foods taken by a mother from the common pot (Dop *et al.*, 1994).

Eating patterns and meal frequency are not fixed in many rural settings and can vary depending on the season, geographical area, or proximity to water/fuel. Seasonal variations can arise due to the demands for agricultural work. During the lean season, when a large amount of time is spent in the field, it is more common for food to be prepared in one pot for the whole day. A method for gathering information on the local cultural context, including local and cultural eating habits and patterns, recipes, typical food portions and serving sizes, and food sharing practices would be to employ key informants in the community and establish focus group discussions. In other cultures, including regions of western Nepal, food

sharing is not frequently observed as shown in a study by Gittelsohn (1989). This is due to the cultural belief that the pollution of food occurs by touching another's served portion.

A further methodological consideration to highlight is the consumption of street food (food and beverages). Street food has become more prevalent in many urban areas due to rapid urbanization. In many locations, it can account for one or even two meals per day. Therefore, in low resource settings, the implications of street food consumed outside the home must be considered when developing a dietary assessment protocol and when assessing food intake. Another factor to consider is that dietary quality is affected by the consumption of processed foods and food products. Depending on the degree of processing, classification and data analysis will change, and have an effect on the information that should be collected by food consumption surveys. For this reason, FAO has produced a publication, *Guidelines on the collection of information on food processing through food consumption surveys*¹². It provides guidance on how to collect accurate and reliable information on processed foods and food products when conducting a dietary survey.

3.4 ESTIMATION OF PORTION SIZE

Estimation of portion size can be challenging, especially in low resource settings given the absence of standard portion size models for local foods, the absence of non-standard serving sizes and measurement tools. Those involved in the preparation of the survey can use different ways to avoid potential errors introduced to the survey during portion size estimation. These include visual aids such as a food atlas, food models or empty food packages that include local foods. It is important to make efforts to devise serving sizes in a way that is conventional, simple and

¹² <http://www.fao.org/3/a-i4690e.pdf>

easy for respondents to follow. This can be done using tools that the respondents are familiar with – such as household measures, i.e. cups, glasses, bowls, plates and spoons – calibrated prior to the study (Cassidy, 1994). Providing respondents with training prior to the study can also help reduce measurement error. Additionally, protocols to estimate individual portion sizes need to be developed to account for families that eat from shared pots or plates (Kigutha, 1997). In this regard, innovative technologies such as image-assisted measurement methods to overcome some of the pitfalls associated with portion size estimation can be employed (Gemming *et al.*, 2015; Illner *et al.*, 2012). However, many of these technologies are still at an early stage of development hence the effectiveness of using such methods in low resource settings would need to be evaluated. As such, validation studies are recommended before adopting new technologies in different settings. It is important to take into account the eating habits of the studied populations in order to make portion size estimation easier, for example stews being served on top of a dish can cover other foods underneath.

3.5 AVAILABILITY OF FOOD COMPOSITION DATA

A food composition database (FCD) is needed to convert dietary data into energy and nutrient intakes. The number of food items included in FCDs may vary from database to database. It is essential that FCDs are continuously updated to provide new foods and food composition values. Furthermore, the best values included in FCDs should be analytical data derived from a laboratory that include local foods as much as possible. Nonetheless due to financial constraints this may not be feasible, leading to the estimation of values based on borrowed data from non-local foods

(Schakel *et al.*, 1997). It is important that users of this guide are aware of the value of identifying a full set of updated national/international FCDs for nutrient analysis, as accurate determination of nutrient intakes from dietary studies is dependent on the availability of up-to-date FCDs specific to the countries, regions or localities.

Due to a lack of resources and capacity in low resource settings food composition information is often borrowed from publicly-available resources such as the USDA¹³, or FCDs from neighbouring countries. This approach may not always be directly applicable to the country in question, as these FCDs may be lacking information on local foods consumed or are not as reliable and up to date as other sources. This may result in the introduction of a variety of errors. Kigutha (1997) remarked that a lack of an appropriate database that takes into account regional specific consumption patterns and local foods was one of the main barriers when interpreting food consumption data in the study. Harmonizing FCDs from different countries can also occur, creating comparability problems. However, programmes have been developed to merge different FCDs, such as *The European Food Information Resource*, which helps to harmonize databases among European countries¹⁴.

Outdated FCDs may not include updates on new foods or new food compositions (e.g. a manufacturer's formulation may change) or up-to-date information on fortified or processed items. This can lead to inaccurate measures of nutrient intake. Converting food consumption data into measures of nutrient intakes via matching to specific FCDs is challenging and critical for obtaining high quality estimates of nutrient intakes. Food matching aims for the highest quality match possible by identifying the most appropriate food in the most applicable FCD. If

¹³ USDA, National Nutrient Database <https://ndb.nal.usda.gov/ndb/search> (Accessed 1 December 2016)

¹⁴ The European Food Information Resource <http://www.eurofir.eu/home/> (Accessed 1 December 2016)

obtaining this is not possible, it is recommended to use a stepwise approach, i.e. to search for the highest quality food match. To properly assess nutrient intake, there can be no missing food composition values, and therefore, a food match must be made. For further information on how to perform food matching, readers can refer to the FAO INFOODS *Guidelines for Food Matching*¹⁵. INFOODS¹⁶ is also assisting countries in compiling and publishing food composition databases and tables according to international standards. The INFOODS website provides additional information on work that has been undertaken regarding food composition databases, and a number of relevant technical documents on food matching, internal checks on food composition data *prior* to publication and guidelines for converting units, denominators and expressions¹⁷.

3.6 MEASUREMENT ERRORS IN DIETARY ASSESSMENT

Measurement error¹⁸ can be in the form of random error and/or systematic error. The former reduces precision and the latter results in incorrect estimates. Hence, it is important to quantify and correct for these effects. In general, the errors that affect the validity of a dietary assessment method are systematic and those associated with reproducibility are random (Gibson, 2005; Willett, 2013a).

- Random errors may occur across all respondents and all days, causing associations to be underestimated and even failure to detect associations in the first place. This type of error

can be minimized by increasing the number of measurements.

- Systematic errors may be respondent-, food- or interviewer-specific and can result in underestimated or overestimated associations; these type of errors cannot be minimized by increasing the number of measurements.

Major measurement errors are due to: nonresponse bias, respondent biases, interviewer biases, respondent memory lapses, incorrect estimation of portion size, supplement usage, coding errors, mistakes in the handling of mixed dishes, etc. (Gibson, 2005). Table 12 describes possible sources of error that should be considered in different dietary assessment methods. Depending on the population group being studied, it is important to employ appropriate strategies to optimize the information being retrieved and reported to the investigator, and to minimize errors. Considerable efforts have been made in developing statistical techniques to deal with these errors and to enhance the performance of various methods. Linear regression calibration, energy adjustment and analysis of variance can be used to correct for random and systematic errors during the data analyses stage (Slimani *et al.*, 2015).

¹⁵ FAO INFOODS <http://www.fao.org/docrep/017/ap805e/ap805e.pdf> (Accessed 1 December 2016)

¹⁶ FAO INFOODS <http://www.fao.org/infoods/infoods/en/> (Accessed 1 December 2016)

¹⁷ FAO INFOODS <http://www.fao.org/infoods/infoods/standards-guidelines/en/> (Accessed 1 December 2016)

¹⁸ For an interactive way to learn more about the implications of measurement error, readers are also encouraged to visit the [Measurement Error Webinar Series](#), organized by collaborators from the National Cancer Institute, the Office of Dietary Supplements, the United States Department of Agriculture, the Gertner Institute, Texas A&M University, and Wake Forest University.

Table 12 - Sources of errors in direct dietary assessment methods for assessing food and nutrient intakes

SOURCE OF ERROR	ESTIMATED FOOD RECORDS*	WEIGHED FOOD RECORDS*	24-HOUR RECALL* [∞]	DIETARY HISTORY	FFQ	BRIEF DIETARY INSTRUMENTS	DUPLICATE MEAL METHOD
Food composition table	+	+	+	+	+	-	-
Food coding	+	+	+	+	+	+	+
Incorrect weighing of food	-	+	-	-	-	-	+
Reporting error	+	+	+	+	+	+	-
Diet variations with time and season	+	+	+	+	+	+	+
Wrong frequency	-	-	-	+	+	+	-
Modified eating pattern	±	±	+	+	+	+	+
Respondent memory lapses	-	-	+	+	+	+	-
Portion size estimation	+	-	+	+	++	-	-

SOURCE OF ERROR	ESTIMATED FOOD RECORDS*	WEIGHED FOOD RECORDS*	24-HOUR RECALL * [∞]	DIETARY HISTORY	FFQ	BRIEF DIETARY INSTRUMENTS	DUPLICATE MEAL METHOD
Respondent bias	±	± ***	+	+	+	+	-
Interview bias	±	±	±	+	±	-	-

Adapted from FAO (2002).

* Image-assisted dietary assessment methods inherit the sources of errors of conventional dietary assessment methods

[∞] Dietary diversity score inherits the sources of errors of the 24-hour recall, when data was collected by a 24-hR recall questionnaire

± Means that there is a possibility for the method to be affected by the source of error

** Only in cases of quantitative FFQ

*** Occurs in cases where a field worker or nutritionist is weighing the food (happens in low resource countries)

3.6.1 MISREPORTING ENERGY INTAKES

Misreporting is characterized by the reporting of implausibly low (underreporting) or high (overreporting) energy intakes¹⁹ at person or group level. Energy intakes are often used as proxies of dietary intakes, therefore, if energy intakes are underestimated then intakes of other nutrients are also underestimated. Underreporting is a widely acknowledged limitation of dietary assessment methods characterized by unlikely low reports of habitual energy intake. Underreporting can occur because of the presence of systematic bias and differential misreporting, and can lead to an overestimation of undernourishment prevalence.

The magnitude of underreporting varies with different dietary assessment methods (Burrows *et al.*, 2010; Poslusna *et al.*, 2009). A better understanding of other factors and mechanisms that are inter-related when dietary data are gathered will help enhance the quality and accuracy of dietary assessment surveys. Such factors include body weight and BMI, gender, socio-economic position, memory disturbances, motivation and social expectations (e.g. dieting), and the nature of the study environment. Specific social mechanisms involved in producing errors may include lack of motivation and deliberate or subconscious errors in recordings. These may be inevitable in some groups or linked to the nature of the activity itself. Memory disturbances can be especially prevalent in the elderly, while other social groups may demonstrate disturbed perceptions of body image, a preoccupation with weight, and therefore an unhealthy attitude towards food leading to the unintentional or intentional misreporting of dietary intakes (Hill *et al.*, 2001).

Validating dietary surveys to identify the degree of misreporting can be carried out by comparing

reported energy intake with Total Energy Expenditure (TEE), often measured by using biomarkers for energy intake (DLW technique) (Subar *et al.*, 2003). However, this method is expensive because it requires laboratory-based investigation. A review conducted in low resource countries found that only a few studies concerning underreporting have been conducted, and none of them have used DLW. Most of these studies have used an alternative approach to identify the degree of underreporting, using the standard equation for estimating basal metabolic rate (BMR) for different ages, gender and body weight, as proposed by (Schofield, 1984). On the other hand, the Goldberg cut-off method is also used, despite concerns about its accuracy (Tooze *et al.*, 2012). It is based on the assumption that energy intake (EI) is equal to total energy expenditure (TEE) assuming weight stability and using a 95 percent confidence interval for statistical comparison with reported EI/BMR and physical activity level (PAL), given by TEE/BMR (Black, 2000; Coward, 1998). Information on PAL can be obtained by observing physical activity, applying lifestyle questionnaires to the target population, or generating information from very similar samples. An EI/BMR ratio below the calculated cut off indicates implausible reported energy intake, i.e. energy intake is too low to reflect the true habitual energy requirements. Increasing the number of days of dietary assessment may help to resolve this issue, because it increases precision and reduces within-person variability. However, it should also be noted that long recording periods reduce reporting accuracy because they increase fatigue and boredom. Rankin *et al.* (2011) assessed energy intakes by 24-hour recalls investigating a group of peri-urban African adolescents and compared the results with estimated energy expenditure. The latter was estimated using BMR equations and estimated physical activity factors

¹⁹ In a state of energy balance, energy intake is assumed to be equal to TEE considering that the individual's weight is stable.

derived from the previous day's physical activity recall. After calculation of energy expenditure, the relative validity of reported energy intake derived from multiple 24-hour recalls was tested using correlation analysis. It was reported that the 24-hour recalls collected at five different measurements over two years offered poor validity between energy intakes reported (Elrep) and energy expenditure estimated (Elest). Goldberg's formula identified cut off points for under and over reporting of energy intake. The ratio Elrep: Elest was calculated and compared with the Goldberg cut-offs, indicating that 87 percent of the boys and 95 percent of the girls underreported their dietary intakes, whereas only 2 percent of boys and girls over reported their energy intakes (Rankin *et al.*, 2011).

Managing misreporting prior to data analysis has been carried out in different ways: for example, by excluding individuals who have an EI/BMR ratio below the calculated cut off (note that this may leave out individuals who do in fact have a true low intake level), or by performing energy adjustments. Misreporting cannot be completely avoided, because of the self-report nature of the dietary assessment methods. Nevertheless it is important to attempt to deal with this, especially in regard to factors affecting a subject's memory lapses – one of the main causes of under-reporting (Poslusna *et al.*, 2009). Solutions can include a more careful design of the study so that it uses appropriate methodologies and standardized procedures, along with the use of visual memory aids and effective interpersonal communication between respondents and field workers to ensure the quality of the reported data.

3.7 REPRODUCIBILITY IN DIETARY ASSESSMENT

Reproducibility, also referred to as precision in dietary assessment, assesses the degree to which a method provides similar results when used repeatedly (on two or more occasions). Reproducibility is commonly determined by a "test-retest" design, where the same dietary method is repeated on the same subject over the same time period, after a defined time interval between repetitions. True reproducibility cannot be assessed, because it is not possible to replicate observations in dietary intakes; only estimates can be obtained (Gibson, 2005). Reproducibility of a dietary assessment method is affected by a series of factors that need to be taken into consideration (Gibson, 2005; Patterson *et al.*, 2004; Willett *et al.*, 2013):

- Time interval between administrations; the selection of the time interval depends on the time frame of the dietary method used. It is preferable to choose non-consecutive days. An interval of four to eight weeks was recommended (Block *et al.*, 1989) to ensure that the second measurement does not influence the previous one.
- The mode of administration will depend on the selected dietary method. The same method of administration should apply to other respondents of the same study.
- The method used to estimate portion size, including approaches and tools, must be consistent throughout.
- Seasonality can cause changes to food habits, and therefore must be taken into consideration in any dietary assessment. This is especially important in low-resource countries where seasonality may greatly alter food access and food availability for people.

- Sample size should be calculated by taking into consideration the degree of precision of a specific dietary method, provided that the within-subject variation is known.
- The intake of nutrients may have changed between the intervals of the two assessment periods as a result of daily variation of food intake.

Increasing the number of recalls or records improves reproducibility by decreasing the effect of random measurement error on the mean. The number of recall and record days would also depend on the variability of the nutrient of interest and the dietary method used. The reproducibility of any dietary method is a function of the measurement errors (see section 3.6 on measurement errors) and uncertainty resulting from true variability in daily nutrient intake and modulating factors such as sex, age, seasonality, dietary habits, etc. Steps can be taken to minimize measurement errors and confounders; however, actions should not be taken to minimize true variation in daily intake since it characterizes the true usual intake of a group of subjects. The sources of true variability in nutrient intakes include: between- and within-subject variability, age, sex, daily variation, seasonality and sequence effect (Gibson, 2005). These are described in further detail in section 3.1, sources of dietary variation.

Between- and within-subject variability can be estimated statistically using analysis of variance, as long as data of two or more days of food intake is available, in at least a subsample of the population. This implies that between- and within-subject variability cannot be calculated for FFQ or dietary history, which collect retrospective data that do not vary from one day to another. Statistical methods to assess reproducibility, at group level, based on the test-retest design include: a paired t-test for comparing the mean intakes (when data is normally distributed) and a Wilcoxon matched-pairs or signed-rank test for comparing the median intakes (for non-normally distributed

data). If the aim is to assess reproducibility at an individual level, the simplest statistical method is to calculate the percentage of misclassification, by comparing the number of pairs with exact agreement. Alternatively, the limits of agreement and 95 percent confidence intervals between two replicates can also be calculated (Bland *et al.*, 1986). The source of true variability may certainly affect reproducibility. Nonetheless, although there may be disagreement between the data collected from dietary assessments on two separate occasions, the method may not have a poor reproducibility but the food intake of the individuals may indeed have changed (Gibson, 2005). A dietary assessment method may have good reproducibility and yet have poor validity, but a method that has a good validity cannot have poor reproducibility.

3.8 VALIDITY IN DIETARY ASSESSMENT

Validity assesses the degree to which a test method actually measures what it is intending to measure. Dietary assessment methods developed to evaluate usual intakes in subjects are the most difficult to validate because the true intake level can never be known with absolute certainty. Thus, the absolute validity of dietary assessment methods is only determined in dietary studies that involve a limited number of subjects or cover a short time frame (Gibson, 2005).

Alternatively, relative validity studies assess the extent of agreement between the “test” and the “reference” method (please note that throughout this resource guide, the words “validity” and “relative validity” will be used interchangeably). Relative validity is a process where dietary estimates from a test method are compared with those from a reference method, which has a greater degree of demonstrated validity (Brink, 1991). Table 13 presents examples of the pairing of test dietary assessment methods and the

Table 13 - Examples of pairing test dietary assessment methods and reference methods

TEST METHOD	REFERENCE METHOD
Single 24-hour recall	Single day weighed record
Multiple 24-hour recalls	Multiple day weighed records
FFQ over 1 year	Four 7-day weighed records at 3-month intervals over 1 year and spaced to account for seasonal variation
Dietary history over 1 month	Single day weighed record spaced evenly over 1 month, the number depending on the nutrients

Reproduced from Gibson, 2005.

reference methods. The purpose of conducting validation is to better understand how the test method works in particular research settings and to use that information for more accurate interpretation of the results from the overall study.

When designing validity studies, please refer to:

- The selection of subjects, who should be representative of those in the main study. Validation studies are relatively expensive to conduct, and are done in small samples compared with the size of the main study. The validation sample, however, should be large enough to estimate the correlation between the test and the reference method with reasonable precision (Thompson *et al.*, 2013). Increasing the number of respondents and decreasing the number of repeat measures per respondent can often help to increase precision without extra cost. It is recommended that the subsample is randomly selected with the sample size of the main study.
- The subject's physiological characteristics, including sex and age, as validation studies have shown that the response of women to dietary assessment differs from that of men (Johnson *et al.*, 1994). Thus, validity studies should be tested separately on both men and women.

- Socioeconomic status, ethnicity and health status of the subjects. This may affect the outcome of validity studies through their link with dietary diversity, as was shown in the validation of a FFQ (Kristal *et al.*, 1997).
- The study objectives. The reference method must have the same study objective and must measure similar parameters over the same time frame (i.e. the current, past or usual intake) as the test method (see the possible levels of objectives in Table 14) (Gibson, 2005).
- The sequence and spacing of test and reference methods also needs to be considered carefully. In general, the test method should be administered prior to the reference method in validation studies, in order to simulate the situation chosen for the proposed study. Spacing between the administration of the methods is equally important, so that the completion of the test method does not influence responses to the reference method (Gibson, 2005).

The reference method must also differ from the test method in terms of specific primary errors. This means that errors in the reference method should be independent from those in the test method and also of the true intake (Nelson, 1997). Therefore, the reference method must differ from

the test method in aspects such as reliance on memory or the method used for estimating portion size (Gibson, 2005). More about measurement errors can be found in section 3.6. The subsequent data analysis, in validation studies, quantifies the relationship between the test method and the reference method, and the resulting statistics can be used for a variety of purposes. Readers are advised to seek statistical advice before conducting this task. The statistical methods to be used to evaluate validity will depend on the objectives of the study. For example, for a level one objective (See Table 14) only the extent of the agreement on a group basis is required, but for levels two to four objectives, an assessment of the validity of individual dietary intake is required. Any assessment of validity should consider each nutrient of interest separately. In addition, confounders such as age, sex and food habits must be considered in the interpretation of the results (Gibson, 2005). There are different statistical approaches that can be used in validation studies, for detailed information readers are recommended to consult chapter 8 of *Design Concepts in Nutritional Epidemiology* (Nelson, 1997). In general, it is suggested that more than one statistical method should be used for more accurate interpretation of the results. The first step prior to conducting any statistical tests is to check if the data generated are normally distributed (Shapiro *et al.*, 1965; Vickers, 2005). If this is the case, a parametric statistic test²⁰ should be used. If the distribution of the nutrient intakes is skewed (non-normally distributed), attempts should be made to normalize the data (i.e. log-transformations) or follow non-parametric statistical tests (Nelson, 1997; Vickers, 2005). Please find below a brief description of some of the statistical methods used to assess validity (Bountziouka *et al.*, 2010; Gibson, 2005; Lombard *et al.*, 2015; Nelson, 1997):

Comparison of means and medians

- Tests to measure the comparison of means or medians (unpaired comparisons) for a level one objective to assess validity at group level. The validation should assess the ability of the test method to reflect the group mean intake. Comparison between test and reference measures of mean or median nutrient intakes are best examined using a t-test (parametric) or a Wilcoxon signed rank (non-parametric) test respectively (Gibson, 2005; Nelson, 1997). These tests will inform if the two means or medians are statistically different (e.g. $P < 0.05$, $P < 0.01$) at some predetermined probability level (e.g. probability 95 percent, 99 percent). Careful interpretation of the results should follow the statistical analysis. For example, if differences between the means/medians for the test and reference methods (i.e. $P > 0.05$ for a 95 percent probability) are not significant for multiple nutrients and if the differences are in different directions (i.e. positive and negative) there is no bias present in the test. This means that the test method can be used to replace the reference method. Landais *et al.* (2014) assessed the validity of a FFQ in Morocco by comparing it with a 24-hour recall. The authors used a Wilcoxon signed rank test to compare the mean food intakes. It was found that the test method FFQ slightly underestimated mean fruit and vegetable intakes by 10.9 percent, and this underestimation was significant at $P = 0.006$. The authors also used Spearman correlations and Bland–Altman plots to assess the validity of the test method. They concluded that the FFQ in this study was valid to be used at population level but not at the individual level (Landais *et al.*, 2014).

²⁰ Parametric statistics is a branch of statistics which assumes that sample data comes from a population that follows a probability distribution based on a fixed set of parameters. A non-parametric model differs precisely in that the parameter is not fixed and can increase, or even decrease if new relevant information is collected.

Correlation analysis

- Correlation coefficients, i.e. Pearson (parametric) and Spearman (non-parametric), measure the strength of the relationship at individual level, between the intakes when conducting the test and the reference methods. Correlations have a maximum value of +1, indicating positive correlation and a minimum value of -1, indicating negative correlation. Values around 0 reflect no linear relationship between the two measurements. For example Coulibaly *et al.* (2009) used correlations (in addition to comparisons between groups t-tests) for assessing the validity of an FFQ compared with 48-hour dietary recall in Mali. In this study the correlation between protein intakes estimated by the FFQ and the 48-hour recalls was 0.63 ($P < 0.0001$) (Coulibaly *et al.*, 2009). The P value indicates that the correlation between the two methods was highly significant. In addition, in most cases, the reference method is itself imperfect and subject to within-person variation or day-to-day variability (see measurement error section 3.6). In these cases, measures such as correlation analysis may underestimate the level of agreement with the actual usual intake. This type of underestimation is known as “attenuation bias”; it can be addressed through the use of measurement error models i.e. de-attenuating the correlation coefficients, using the ratio of intra- to inter-subject variation (the variance ratio) as calculated from the replicate observation in the reference method. This correction will result in estimates that nearly reflect the correlation between the diet measure and true diet (Carroll *et al.*, 1998). Likewise, various researchers have recommended energy-adjusting nutrient intake

prior to conducting the correlation analysis (Bingham *et al.*, 1997). Energy adjustment is determined by means of the nutrient density, dividing each nutrient value by the energy intake of that particular subject. Nutrient density is then used in statistical analysis instead of the original value of nutrient intake (Bingham *et al.*, 1997). The correlation coefficient describes only one aspect of agreement between the two dietary methods. Several authors have noted that poor agreement between the test and reference method can exist even when correlation coefficients are high (Bingham *et al.*, 1994; Bland *et al.*, 1986). A detailed review on the limitations of correlation analysis for validation of dietary assessment methods was addressed by Bland and Altman (Bland *et al.*, 1986). Thus, a preferable alternative is to use linear regression analysis, and/or additional statistical measures of agreement to characterize the relationship between the test and reference methods.

Regression analysis²¹

- Regression analysis can be considered an extension of correlation analysis. In a linear regression, total energy intake is designated as an independent variable (x) and intake of the nutrient of interest as a dependent variable (y). This analysis aims at finding the best mathematical model ($y = a + mx$) for predicting the dependent variable (y) from the independent variable (x) (Beaton, 1994). To measure how well the data fit, a regression coefficient can be obtained by calculating r-squared (r^2) (the closer to 1 the better the fit). When the model is adjusted a t-test can be used to assess whether the slope (m) of the regression line is statistically significant and how far away it is from zero. If so, the model indicates that

²¹ Linear regression attempts to model the relationship between two variables by fitting a linear equation ($y = a + mx$) to observed data. One variable is considered to be an explanatory variable (x), and the other is considered to be a dependent variable (y). For example, one can use this to relate the weights of individuals to their heights, or in this case to relate the results from two dietary assessment methods. <http://www.stat.yale.edu/Courses/1997-98/101/linreg.htm> (Accessed 1 December 2016)

the test method is valid to be used instead of the reference method. (Hernández-Avila *et al.*, 1998) in Mexico validated a semi-quantitative FFQ against 24-hour recalls, where validity was evaluated using regression analysis (and correlation coefficients). The authors found significant regression coefficients between 24-hR recall and FFQ, ranged from 0.147 to 0.55 for Vitamin B6 and carotenoids respectively. The regression correlations were significant ($P < 0.05$, $P < 0.0001$) for most of the nutrients ($n=24$), except for polyunsaturated fat, folic acid, vitamin E and zinc. It was concluded that the test-FFQ provided a useful estimate by which to categorize individuals by level of past nutrient intake.

Additionally, more complex multiple regression models can be applied. Such models can allow for control over the effect of confounders (i.e. smoking, total energy intake, BMI, etc.) (Nelson, 1997). Readers can find more examples of these multiple regression models in the following references: Martin-Moreno *et al.*, 1994; Sanz-Gallardo *et al.*, 1999.

Cross-classification

- Cross-classification consists of grouping subjects into different categories of consumption (i.e. tertiles (thirds), quartiles (fourths) or quintiles (fifths)) by the test and reference methods. The percentage of subjects being correctly categorized into the same category and the percentage misclassified in the opposite category is determined (Gibson, 2005). Cross-classification evaluates the ability of both methods to classify individuals similarly in categories of nutrient intake. This method is commonly used to validate FFQ. The cross-classification will reflect how well the FFQ separates the subjects into classes of intakes and thus provides an estimate of the validity of the test methods. If the test and reference methods were in perfect agreement, every subject would be classified

in the same category of distribution according to both methods. This means that perfect agreement would be reached with 100 percent of subjects falling in the same category by both methods, and zero percent in adjacent or opposite thirds (Nelson, 1997). (Jackson *et al.*, 2001) evaluated the validity of a quantitative FFQ questionnaire among Jamaicans of African origin, against 24-hour recalls. Results from both methods were grouped in quartiles with cut-off points for quartiles determined separately for each method. Agreement in cross-classification by the two methods was assessed as the proportion of participants similarly classified in the highest or lowest quartiles, and misclassification was assessed as the proportion classified into the opposite extreme quartile, for nutrients. The percentage of participants similarly classified by both instruments ranged from 31.6 percent for retinol to 100 percent for alcohol in the lowest quartile, and from 24 percent for retinol to 60 percent for alcohol in the highest quartiles. Misclassification was low (one or two persons) for most nutrients but was higher for retinol (16 percent) in the lowest quartile and polyunsaturated fat (29 percent) in the highest quartile. It was concluded that this test-FFQ showed reasonable validity and was suitable for estimating the habitual intakes of energy and macronutrients, but was poor for some micronutrients like retinol (Jackson *et al.*, 2001). More examples of cross-classification can be found in FFQ validation studies (Klipstein-Grobusch *et al.*, 1998; Martin-Moreno *et al.*, 1993).


Bland-Altman plots

- Bland-Altman plots are a more reliable method than the correlation coefficient when assessing validity (Bland *et al.*, 1986). The plots are constructed using the mean and the standard deviation of the difference between the test and the reference method for energy and each nutrient intake, evaluated on the

same subject (Bland *et al.*, 1986). The Bland Altman plot is drawn for each nutrient and depicts the mean of the test and reference intake (for each subject) plotted against the difference between each pair of observations (See Figure 3). If there is no bias in the test method, the differences will cluster along the horizontal line at zero ($y=0$), and the mean difference will be close to zero. The plot can also indicate if the differences between the two methods become progressively smaller or larger while increasing the intake (Bland *et al.*, 1986; Gibson, 2005). In addition, it is recommended to calculate the 95 percent confidence limits for the difference between both methods. Therefore, the plots and the confidence limits will indicate if the agreement between the test and the reference methods is acceptable. For example, Figure 3 shows the plot for iron intake from a study validating an

image-assisted 24-hour recall compared with a weighed food record conducted in Bolivia by Lazarte *et al.* (2012). The authors found that the variation around the mean difference line was in the range of 0.9 to 6.0 percent for nutrients and energy intake. For iron, a difference of 5.1 percent with 95 percent confidence limits at 16.7 and 6.3 percent was found. Moreover, the plot indicated that the differences were random and did not exhibit any systematic bias or trend; therefore, this is consistent over different levels of mean intake (Lazarte *et al.*, 2012). Thus the test method, in this study, would be likely to provide comparable results to those obtained from the reference method for the assessment of individual intake.

Figure 3 - Difference in mean iron intake estimated by a 24-hour recall and a weighed food record



Source: Lazarte *et al.*, 2012, reproduced

The validity of self-reported intakes has been questioned on a number of occasions, on account of the influence of misreporting and in particular of under-reporting. A more objective approach to validate dietary assessment methods is the use of biomarkers. Biomarkers are external independent markers of nutrient intake. They respond to the intake of nutrients, preferably in a dose-dependent manner and are available for assessing energy and a limited number of nutrients. They are increasingly being used to overcome the limitations of reference dietary assessment methods (Gibson, 2005; Hedrick *et al.*, 2012; Thompson *et al.*, 2013; Trabulsi *et al.*, 2001; Willett *et al.*, 2013). However they are also subject to error. The high cost of these techniques also make them impractical for validating every dietary assessment study. Most biomarkers are measured in body tissue or fluids. Biomarkers such as DLW²², used to measure energy expenditure (Black *et al.*, 1997; Hill *et al.*, 2001), 24-hour urinary nitrogen excretion to validate protein intake (Black *et al.*, 1997) and urinary potassium as a biomarker of potassium intake (Freedman *et al.*, 2014; Freedman *et al.*, 2015) are widely used and often considered as reference standards to validate nutrient intakes.

In addition, several factors affect the reproducibility and validity of dietary assessment methods:

- respondents' abilities to respond accurately, due to cognitive abilities, literacy and numeracy skills;
- respondent characteristics (i.e. children, adolescents, elderly, low literacy, etc.);
- measurement errors of the instrument;
- the error-proneness of the response format affected by the questionnaire design;
- quality control of coding;
- dietary changes in the time between administrations of the questionnaire;
- quality and adequacy of the reference data (i.e. food composition tables).

Most of these factors are under the control of the investigator, and thus can be controlled and improved. However, as in any survey methodology, dietary assessment methods are subject to measurement error, which is explained in detail in section 3.6. Due to time and financial constraints, such reproducibility and validity studies are often undertaken in a subsample of the main population, and used as test-reference approaches. Results of the evaluation study are taken into consideration when interpreting the dietary evidence obtained from the main study method and sometimes result in a partial correction of these results (Willett, 2013a).

3.9 QUALITY CONTROL AND DATA ANALYSIS ON DIETARY ASSESSMENT

Prior to beginning the analytical process, careful consideration needs to be given to assessing missing data and outliers to ensure data quality. One method used to avoid errors when conducting a dietary assessment is to review the forms during the study, to ensure all information has been entered correctly, either by the investigator or the respondent. However, this method may be useful when assessments are carried out over the phone or by mail (Dennis *et al.*, 2003). Furthermore, criteria need to be set for the exclusion of over-reporting and under-reporting of energy intakes. However, any proposed criteria must be carefully selected to avoid excluding valid subjects who should stay in the final data set. Criteria for inclusion/exclusion have been developed by generating prediction equations

²² DLW technique involves enriching the body water of a subject with an isotope of hydrogen (²H) and an isotope of oxygen (¹⁸O), and then determining the washout kinetics of both isotopes. It's a method for measuring energy expenditure in free-living people, providing estimates of habitual expenditure over a time period of 10–20 days. https://inis.iaea.org/search/search.aspx?orig_q=RN:21093729 (Accessed 20 April 2016)

based on indirect calorimetry or DLW (Ventura *et al.*, 2006). Ranges based on arbitrary allowances (500–3 500 kcal/day for women, 800–4 000 kcal/day for men) have also been utilized to exclude outliers from the final data analysis. In addition, the following actions may help to ensure data quality:

- the organization of an international nutrition coordinating team to supervise nutritionists at country and field levels, and to certify dietary interviewers;
- country-specific training and certification of all staff by competent international and national authorities, based on standard procedures adapted for each country (forms, manuals, materials, and quality control procedures);
- complete and valid food consumption nutrient databases to determine nutrient intakes.

Prior to the start of a study, a statistician needs to be consulted on the correct sample size calculation, study design to be used and what data analysis method should be employed. The statistical analysis method used is dependent on the objective(s) of the study. The investigators must ask themselves if the aim of the study is to characterize mean intakes of a group (requiring the measurement of a single day food intake) or to define the usual intake distribution of a group from the dietary information collected at individual level (requiring the collection of more than one day of dietary data). Assessing the distribution and central tendency of dietary data is an important first step for the data analysis process. If data are skewed, transformations can be applied to normalize the distribution before applying statistical procedures. However, if the distribution cannot be transformed, non-parametric statistical tests can be applied. In many dietary intake studies, a representative sub-sample of the population is randomly selected to participate in the study. The sample is assumed to be representative of the whole population; however, there is always a degree of uncertainty

when extrapolating the results. This is expressed as the confidence interval measuring the precision of the sample estimates.

The mean intake of a population group, based on the measurement of a single day dietary intake that is normally distributed, is the best estimate of the central tendency. If the objective of the study is to investigate the differences between the mean values of two separate population groups, and to determine if the differences present between them truly exist or are by chance, a t-test can be applied. Moreover, a Fisher's or Wilcoxon's test can be used to assess whether the variances are homogenous. Alternatively, if mean intakes of two or more target population groups are under investigation, an analysis of variance test (ANOVA) can be conducted. Analyses of the percentage of individuals at risk of inadequate intake levels can be generated by looking at usual dietary intake data (where collection of dietary intake takes place over a minimum of two days). The risk of inadequate intakes can be examined against other variables (e.g. education, sex, socio-economic status) and the magnitude of the association between the variables can be measured using a number of different tests, including a chi-squared test, relative risk or odds ratio. For further details on statistical tests and procedures, refer to Gibson *et al.*, 2008; Willett *et al.*, 2013.



©FAO/Warren Lee

4

SELECTING A DIRECT DIETARY ASSESSMENT METHOD

Throughout this guide, the need to address a number of issues prior to selecting a dietary assessment method is emphasized. Depending on the study objectives, available resources, target population characteristics and cultural considerations, a suitable method that imposes minimum respondent burden and provides accurate and reliable dietary information would be the best option. This section will address key factors to be considered prior to the selection of a direct dietary assessment method, such as the importance of identifying the study objective, selecting the appropriate study design and technical and financial considerations. This will be followed by a step-by-step guide (Box 10) to facilitate the selection of a suitable dietary method from the *Oxford Handbook of Nutrition and Dietetics* (Webster-Gandy *et al.*, 2011) along with a summary of the major features of direct methods, including their strengths and limitations, to provide readers with a summary of information described in section 2. Lastly, case studies of some past projects are also illustrated, underlining the key steps required in the process of choosing a suitable method.

4.1 STUDY OBJECTIVES IN DIETARY ASSESSMENT

The study objective, a specific result that a research project aims to achieve within a specified time frame and given resources, will influence the dietary assessment method chosen for measuring dietary intakes. According to Gibson (2005), the possible study objectives could be:

- to determine the overall nutritional status of a population or subpopulation;
- to identify areas and populations/subpopulations at risk of chronic malnutrition;
- to characterize the extent and nature of malnutrition within the population/subpopulation;
- to identify possible causes of malnutrition within the population/subpopulation;
- to design appropriate interventions for high-risk populations/subpopulations;
- to monitor the progress of changing nutritional, health or socioeconomic influences, including intervention programmes;
- to track progress towards the attainment of long-term goals.

Table 14 presents the most suitable assessment information in selecting the appropriate dietary method for a particular study objective. Four method for assessing dietary intakes according levels of accuracy and precision have been to your study objective. shown below, which will provide guiding

Table 14 - Selection of a method to measure nutrient intakes to meet four possible levels of objectives

LEVEL OF ACCURACY AND PRECISION	DESIRED INFORMATION	PREFERRED APPROACH
One	Mean nutrient intake of a group	A single 24-hour recall, or single weighed or estimated food record, with a large number of subjects and adequate representation of all days of the week
Two*	Proportion of population "at risk" of inadequate intake, i.e. prevalence of inadequate intake	At least two observations on each individual or a subsample using 24-hour recall or weighed or estimated one-day food records
Three*	Usual intakes of nutrients in individuals for ranking within a group	Two or more observations of 24-hour recall or food records or a semi-quantitative FFQ
Four*	Usual intakes of foods or nutrients in individuals for counselling or for correlation or regression analyses	Even larger number of recalls or records for each individual. Alternatively, a semi-quantitative FFQ or a dietary history can be used

*Levels 2–4 depend on within-person variation of the nutrient of interest, which in turn depends on the chosen dietary assessment method, population, and seasonal variation of intake

Source: Reproduced from Gibson (2005) with permission.

4.2 DESIGN OF THE STUDY

Similarly, the design of a study should also be considered and preferably decided upon prior to final decisions on method selection. Readers are directed to Thompson *et al.* (2013) for more information, who provide an excellent review of this topic. A number of questions should be

considered when selecting an appropriate design for a study. These include the type and scope of information needed, time frame, costs and the level of accuracy and precision required. Table 15 provides a description of the methods commonly used in four main study designs, including cross-sectional, case-control, cohort and intervention studies.

Table 15 - *Dietary assessment methods commonly used in the design of different studies*

STUDY DESIGNS	DIETARY ASSESSMENT METHOD
Cross-sectional	24-hour recall FFQ Brief assessment instruments
Case-control (retrospective)	FFQ Dietary history
Cohort (prospective)	FFQ Dietary history 24-hour recall Food records
Intervention	FFQ 24-hour recall Brief dietary instruments

Source: Reproduced from Thompson (2013) with permission.

4.3 TECHNICAL AND FINANCIAL ASPECTS TO TAKE INTO ACCOUNT

Technical and financial considerations related to expertise and capacity development, time frames, availability of personnel (including interviewers and statisticians), equipment, software, etc. all influence method selection. Drewnowski (2001) has suggested that the minimum cost for conducting the analyses of a four-day food record is US\$120, whereas the cost of analyses for a FFQ is US\$0.50 per respondent or US\$3.50 for a FFQ that is optically scanned and machine analysed. Costs in low resource settings may be higher as a consequence of tool adaptation required for local use, and taking into account access to remote or inaccessible geographical areas and communities.

It must be noted that the cost will depend on factors specific to each country or situation, e.g. human resources, sample size, data processing and analysis, availability of a food composition table, etc. Fiedler *et al.* (2013) reported that the implementation of a 24-hour recall similar to a HCES on a representative sample of around 8 500 households would cost approximately 75 times more than a typical HCES analysis to generate nutrient intake indicators. The authors also provided estimated costs for various parameters, including number of provinces, number of health areas per province, number of supervisors and enumerators, and number of days' training, including field testing. Survey designers should always consider that financial aspects directly affect sampling methodology decisions in surveys (Harrison, 2004). Some basic guidance on choosing and carrying out sample design, and ways to avoid sampling bias, is provided by Gibson *et al.* (2008).

4.4 SUPPORTING INFORMATION FOR THE SELECTION OF A DIRECT DIETARY ASSESSMENT METHOD

The following step-by-step guide to selecting an appropriate diary assessment method should complement the readings in earlier sections of the manual, helping to guide the method selection process. Please use the steps below to supplement your decision making process, providing guidance in choosing the most appropriate assessment method to complement your specific study.

4.4.1 A STEP-BY-STEP GUIDE FOR METHOD SELECTION

The following step-by-step guide is presented to support readers on the selection of a dietary assessment method. The issues presented under each step should be carefully considered before selecting a method.

BOX 10. STEPS AND TIPS ON CHOOSING A DIETARY ASSESSMENT METHOD**STEP-BY-STEP GUIDE TO FACILITATE THE SELECTION OF A DIETARY ASSESSMENT METHOD****1. Defining the objective**

- ✓ The method is determined by the reason for the assessment, e.g. is it part of a research study or to clarify an individual's deficiency? (Refer to Table 14 for more details)

2. Food and/or nutrients?

- ✓ Are data needed at the nutrient level or will a description of food patterns suffice?

3. Decide the conceptual time frame for collecting dietary intake data

- ✓ Past, present, usual?
- ✓ Will a retrospective or prospective method be most applicable?

4. Decide the actual time frame for the respondent to record or recall dietary intake

- ✓ Day, week, or year?
- ✓ Number of interviews (e.g. multiple 24-hour recalls)?
- ✓ Consecutive days?
- ✓ Weekdays and/or weekends?
- ✓ Account for seasonality?
- ✓ Record or recall period is over the last three days or last week?

5. Who will be interviewed and by whom?

- ✓ Children (interview parent, caregivers or the child directly?)
- ✓ Who will conduct the dietary assessment? Do they need training?
- ✓ Who will record the information? (Interviewer or interviewee?)
- ✓ Where will the assessment take place?

6. What type of food and drink need to be assessed?

- ✓ If all, then a prescribed method like a FFQ may be inappropriate.
- ✓ Is information needed on whether foods are raw or cooked and the cooking method used?

7. Estimating how much is eaten and/or how often

- ✓ Is measuring the frequency of food consumption enough to meet the objectives?
- ✓ If the aim is to estimate nutrient intakes then portion sizes will be needed. What method is best suited to the context?
 - Direct quantification: weight, measure volume
 - Indirect quantification: household measures, estimated with utensils and food packages
 - Standard food portions
 - Actual dimensions of the food
 - Food models
 - Photos of food with a range of portion sizes
 - A count of handfuls
 - Proportion of the prepared meal that was consumed, e.g. household intake

8. Comparing intake with recommendations

- ✓ If data analysis includes nutrient analysis: are all foods available in food composition tables?
- ✓ Which nutrient analysis program should be used?
- ✓ Selecting the most appropriate dietary reference values
- ✓ Dietary patterns could be explored or diet scores created to compare with food-based dietary guidelines

9. Assessments resources

- ✓ Available to support the measurement of dietary intake (funds, study duration, food composition data, human resources, study design, software purchases and training of interviewers and interviewees)

10. Effective management, logistical plan and mobilizing intangible resources

(Oxford Handbook of Nutrition and Dietetics 2008 – reproduced with permission)

4.4.2 SUMMARY OF THE MAJOR FEATURES OF THE DIFFERENT DIRECT DIETARY ASSESSMENT METHODS

information provided in the table should provide readers with a summary of information described in section 2.

The following table summarizes the major features (including strengths and limitations) of existing dietary assessment methodologies. The

Table 16 - Comparison of the major features of methods for assessing diet

	ESTIMATED FOOD RECORD	WEIGHED FOOD RECORD	24-HOUR RECALL	FFO	DIETARY HISTORY	BRIEF ASSESSMENT INSTRUMENTS	DUPLICATE MEAL METHOD	IMAGE- ASSISTED DAM	DIETARY DIVERSITY SCORES
Type of information attainable:									
Detailed information about foods consumed	✓	✓	✓	✓ ¹	✓		✓	✓	
General information about food groups consumed				✓		✓			✓
Mea-specific details	✓	✓	✓		✓		✓	✓	
Scope of information sought:									
Total diet	✓	✓	✓	✓ ²	✓		✓	✓	
Specific food groups				✓		✓			✓

	ESTIMATED FOOD RECORD	WEIGHED FOOD RECORD	24-HOUR RECALL	FFQ	DIETARY HISTORY	BRIEF ASSESSMENT INSTRUMENTS	DUPLICATE MEAL METHOD	IMAGE-ASSISTED DAM	DIETARY DIVERSITY SCORES
Reference period:									
Short term (e.g. yesterday, today)	✓	✓	✓		✓		✓	✓	✓
Long term (e.g. last month, last year)				✓	✓	✓			
Adaptable for diet in distant past?									
Yes				✓	✓	✓			
No	✓	✓	✓				✓	✓	✓

	ESTIMATED FOOD RECORD	WEIGHED FOOD RECORD	24-HOUR RECALL	FFQ	DIETARY HISTORY	BRIEF ASSESSMENT INSTRUMENTS	DUPLICATE MEAL METHOD	IMAGE-ASSISTED DAM	DIETARY DIVERSITY SCORES
Cognitive requirements:									
Measurement or estimated	✓	✓					✓	✓	
Memory of recent consumption			✓	✓	✓				✓
Ability to make judgments of long-term diet				✓	✓	✓			
Interference with normal eating habits: (Willett W., 2013a)									
High	✓	✓					✓	✓	
Low			✓	✓	✓	✓			✓

	ESTIMATED FOOD RECORD	WEIGHED FOOD RECORD	24-HOUR RECALL	FFQ	DIETARY HISTORY	BRIEF ASSESSMENT INSTRUMENTS	DUPLICATE MEAL METHOD	IMAGE-ASSISTED DAM	DIETARY DIVERSITY SCORES
Time required to complete data collection:									
Less than 15 minutes						✓		✓	✓
Greater than 20 minutes	✓	✓	✓	✓	✓		✓		
Measurement Errors (Slimani <i>et al.</i>, 2015):									
Systematic misreporting (under-reporting or over-reporting of intake?)	✓	✓	✓	✓	✓	✓	✓	✓	✓
Respondent burden (Slimani <i>et al.</i>, 2015):									
High	✓	✓		✓	✓		✓	✓ ⁴	
Low			✓	✓		✓		✓ ⁵	✓

	ESTIMATED FOOD RECORD	WEIGHED FOOD RECORD	24-HOUR RECALL	FFQ	DIETARY HISTORY	BRIEF ASSESSMENT INSTRUMENTS	DUPLICATE MEAL METHOD	IMAGE-ASSISTED DAM	DIETARY DIVERSITY SCORES
Literacy level is required (Slimani <i>et al.</i> , 2015):									
Yes	✓	✓		✓	✓	✓	✓ ⁵	✓ ⁵	
No			✓					✓ ⁵	✓

Adapted from Thompson (2013) with permission.

¹ This applies when using a quantitative or semi-quantitative FFQ

² Will be dependent on the object of the study (i.e. measuring all food groups or a specific food group)

³ This will vary depending on the specific population group under study (i.e. anorexic individuals will tend to over-report intakes and obese individuals will tend to under-report intakes)

⁴ Depending on the innovative technology and/or the conventional dietary assessment method used

⁵ Except from when data collection is done by trained field workers

4.4.3 CASE STUDIES ON SELECTION OF A DIETARY ASSESSMENT METHOD

The following case studies present details from past projects, specifically highlighting some of the steps taken (shown in box 10) in the process of choosing a particular method. These case studies are just some of the examples available and should be interpreted with care.

- *National UK dietary assessment study in low-income households*

A comparison of four dietary assessment methods, which included both prospective and retrospective techniques (24-hour recall, food checklist, semi-weighed method and weighed food record), has been compared to identify the most appropriate tool for a sample of low-income households in the UK. The best method was selected to be used in the UK national study of diet and nutrition among low-income households for measuring nutrient intakes. The target households commonly had low literacy and numeracy skills and lacked the ability to speak the English language. The study was commissioned by the Ministry of Agriculture, Fisheries and Food. The authors were asked to collect the most valid data and recommend an appropriate dietary method for the target population. Each method was assessed for four non-consecutive days, which included a weekend day. In every household, one child (2–17 years) and one adult (over 18 years) were chosen to take part in the study. Children below the age of 12 were assisted by parents/caregivers for the duration of the study. A different approach was used to estimate portion sizes for each of the four dietary assessment methods. For 24-hour recall, a photographic atlas, household measures and food packaging in different sizes were used to estimate portion size. For the food checklist, conducted either retrospectively or prospectively, household measures were used to

estimate food consumed. For foods consumed that were not available in the food composition table, such as uncommon recipes or items where dietary data was missing, the closest match was used. Following the study, the authors concluded that respondents provided the most detailed and consistent responses across all age groups when using the 24-hour recall method. This result can be attributed to the low response burden of the method and low reliance on numeracy, literacy and language skills. Therefore, recall was chosen as the most appropriate direct dietary assessment method for the national UK study (Holmes *et al.*, 2009; Holmes *et al.*, 2008).

- *Dietary assessment methods used in rural areas of Kenya*

A study was conducted in a group of small landowners in rural areas of Kenya to investigate the influence of seasonality on food consumption in the elderly and preschool respondents. The aim of the study was to evaluate dietary intake using two methods, i.e. 24-hour recall and weighed food records, for identifying the most suitable assessment method for the target population, and find the technique that would best reduce time, cost and respondent burden, as well as improve accuracy and reliability. A retrospective method, 24-hour recall, was selected for its yield of reliable information in rural areas and its low cost and low respondent burden. Additionally, 24-hour recall was simple to conduct even with respondents low in literacy and numeracy skills. Three non-consecutive day 24-hour recalls were conducted at three different periods of the agricultural cycle to determine usual dietary intake of the target population and account for seasonality. Household measures were used to estimate food intake. Nutrient values were determined by a food composition table developed by the Technical Centre for Agriculture and Rural Cooperation – East, Central and Southern Africa (CTA–ECSA). However, not

all foods were included in the table and further sources of information had to be used (Kigutha, 1997).

- *The EFSA study: Harmonizing food consumption data across Europe*

The main objective of the European Food Safety Authority (EFSA) is to produce available, detailed, harmonized and high quality food consumption data for use in dietary exposure assessments. EFSA coordinated the pan-European food consumption survey methodology (“EU Menu”) in close collaboration with the EU Member States. The project aimed to harmonize data collection in food consumption across Europe and was endorsed by the EFSA Network on Food Consumption Data to collect data from population groups whose ages ranged from 3–79 months and 10–74 years. The use of a food record was considered the most appropriate method for the “EU Menu”, because it enabled parents and/or caregivers to record detailed information about all foods and drinks consumed by the infant or child, including information on brands and specific products. Furthermore, 24-hour recall was selected for the adult population because of its low response burden. Portion size and food intake estimates were made using a picture book, which included country specific dishes (EFSA, 2014).



©FAO/Paballo Thekiso

5

KEY MESSAGES AND THE WAY FORWARD IN DIETARY ASSESSMENT

5.1 KEY MESSAGES

- The aim of this resource guide is to support readers in selecting a dietary assessment method for a context specific purpose and to provide information on the principles and practical considerations that are important in the selection process, with a focus on low resource settings.
- This guide was written for professional readers, who have knowledge in nutrition and/or dietetics, and more specifically, dietary assessment. The purpose of creating the manual is to support an informed decision-making process for professionals to select a dietary assessment method and to provide a resource for those who wish to refresh or update their knowledge in the subject. In this way readers could gain a better understanding of the rationale for choosing an appropriate dietary assessment method, prompting them to consider the most relevant questions to guide the selection process, while others are provided with a resource for updating their knowledge and information on the topic of dietary assessment methods, specifically in low resource settings.
- Dietary assessment estimates can be obtained at individual or household levels, and their results can be presented as energy and nutrient adequacy, dietary quality, food patterns, intakes of individual foods and food groups, intakes of macronutrients and micronutrients, and diet composition. These estimates are obtained via direct and indirect retrospective or prospective assessment methods. The information gathered is then used to inform various purposes, including target setting, risk assessment, monitoring and surveillance of food consumption patterns and analysis of diet–disease relationships. The information is also useful for the development of a range of food-based indicators for policy process across governmental departments such as agriculture, health, education, finance, planning, trade, etc.
- The study objectives will guide and determine what needs to be assessed, in what detail, of whom and by whom and what are the determining factors that drive the method selection. Furthermore, the level of accuracy and precision needed for the study can also play a role in guiding the selection process for the most appropriate dietary assessment method to choose.

- It is important to be aware that uncertainties, errors, and the unsuitability of different dietary methods for specific programmatic decisions can lead to inaccurate interpretations of findings, resulting in wasted resources. When selecting a method, consideration should be given to the study objectives, resources available and characteristics of the study population, i.e. its culture, religion and particularly the cultural food habits.
- It is important to address a number of methodological considerations prior to selecting the most appropriate dietary assessment method which can help to guide the decision making process. These methodological considerations include: the number of days required to estimate intakes; training needs of the participants; seasonality effects; geographical location; individual, community and culturally-specific issues; intra-household food distribution; shared eating occasions, street food and eating out habits; estimation of portion sizes and food composition databases.
- It should be noted that all dietary assessment methods need to be tested for their validity and reproducibility in assessing food and nutrient intakes. Furthermore, one should also keep in mind that all the dietary assessment methods are subject to measurement errors, which can be systematic or random. Understanding these and defining strategies to address sources of error could result in better estimates and measurements.
- While the integration of advanced innovative technologies – i.e. smartphones, mobile applications, software, sensor-based recording – to improve dietary assessment has shown progress in developed countries, their usage in low resource settings and in low literacy populations is still limited. Nevertheless, technologies such as digital photographs have been successfully used in low resource settings; researchers in developing countries should be encouraged to investigate these technologies further.
- Proxy tools such as dietary diversity scores are useful in low resource settings due to their simplicity and low cost. Nevertheless, they need to be adapted to the local context and the specific characteristics of the population, prior to their application. It should also be noted that they are not suitable methods for assessing individual-based and/or quantitative dietary intakes.
- One of the few available reviews on the application of dietary assessment methods in programmatic decisions concluded that none of the methods reviewed could be considered a “gold standard”; some are better suited to particular contexts than others; the validity and usefulness of a method for a given purpose should guide its selection (taking into consideration sources of potential bias and error); and that simple informed modifications to data collection and processes can improve their usability in programmes.
- When using this guide, users should remember that dietary assessment is one of the four nutrition assessment components, the others being anthropometry, biomarkers and clinical assessment. These other complementary methods should not be overlooked when planning an assessment but are outside the scope of this manual to expand upon further. Equally crucial to nutrition-related programmes is the assessment of food intake practices, knowledge, and beliefs. (Note that these are likely to be of greater importance within the context of developing countries than developed countries.) No description of the assessment of these practices is provided here, but readers interested in this topic are directed to the *FAO Guidelines for assessing*

*nutrition-related Knowledge, Attitudes and Practices*²³.

5.2 THE WAY FORWARD

- The need for such a resource guide is reflected in global economic developments and changing patterns across the food chain, which necessarily entail a corollary impact on individuals' diets and population health. The information presented complements ongoing work in FAO's Nutrition and Food Systems Division that supports global efforts to strengthen national nutrition information systems *vis-à-vis* monitoring and implementation processes. Examples of such work include the FAO/WHO Global Individual Food Consumption Data Tool (FAO/WHO GIFT), which aims "to collect, harmonize, and disseminate data available at national and sub-national level all over the world through an FAO hosted web-platform"²⁴. This platform, intended for use by both experts and a broader audience, aims to facilitate access to microdata and to compute food-based indicators – such as average leafy vegetable consumption, main food sources of vitamin A or high levels of fish consumption – so that data is comparable between different population groups and geographical areas.
- Moving on from ICN2 to the post-2015 sustainable development agenda requires coordinated and synchronized efforts in dietary assessment. Countries can learn valuable lessons from each other's experiences; quite often the international community raises such issues, including calls for stronger national mechanisms for monitoring and evaluation of a population's nutrition. When resources are available and infrastructure is in place,

monitoring and evaluation of a population's diet is relatively straightforward. Within such contexts, opportunities exist or arise to improve individual-based methods, understand their errors better and their influences on data interpretation. However, when resources, capacity and infrastructure are not sufficient or are non-existent, professionals may want to make use of the existing nutrition information structures already in place. An example is using the HCES as a secondary source of data to inform policies with recent calls for improving their relevance and reliability in nutrition policy-making.

²³ <http://www.fao.org/docrep/019/i3545e/i3545e00.htm>

²⁴ <http://www.fao.org/nutrition/assessment/food-consumption-database/en/>



©FAO/Vasily Maximov

6

FURTHER READING

The following list of links supplements the content of this guide. The list is by no means exhaustive, but aims to provide further material of interest to readers for their information. Readers are also encouraged to perform their own searches of the literature for more specific information related to their needs.

Dietary assessment methodology

- **World Health Organization.** *Preparation and use of food-based dietary guidelines.* 1996. Report of a joint FAO/WHO consultation, Nicosia, Cyprus. Geneva, WHO.
- Epidemiology and Genomics Research Program, United States National Cancer Institute.
- **Gibson, R.S. and Ferguson, E.L.** 2008. *An interactive 24-hour recall for assessing the adequacy of iron and zinc intakes in developing countries.* Washington, DC, International Food Policy Research Institute (IFPRI).
- **Cade, J., Thompson, R., Burley, V. and Warm, D.** 2002. Development, validation and utilisation of food-frequency questionnaires – a review. *Public Health Nutrition*, 5(4): 567–87.
- **Coates, J., Colaiezzi, B., Fiedler, J., Wirth, J., Lividini, K., and Rogers, B.** 2012. *Applying Dietary Assessment Methods for Food*

Fortification and Other Nutrition Programs. Gain Working Paper Series No. 4. Geneva, Global Alliance for Improved Nutrition (GAIN).

- **Winichagoon, P.** 2008. Limitations and resolutions for dietary assessment of micronutrient intakes. *Asia Pacific Journal of Clinical Nutrition*, 17(S1): 296–298.
- **Gibson, R.S.** 2005. *Principles of Nutritional Assessment.* 2nd edition. New York, Oxford University Press.

Food consumption manuals for field studies

- **FAO.** 2004. *Uses of food consumption and anthropometric surveys in the Caribbean: how to transform data into decision-making tools.* Rome. Food and Agriculture Organization of the United Nations.
- **den Hartog, A.P., van Staveren, W.A. and Brouwer, I.D.** 2006. *Food habits and consumption in developing countries: Manual for field studies* Wageningen (Netherlands), Wageningen Academic Publishers.
- **European Food Safety Authority.** 2009. General principles for the collection of national food consumption data in the view of a pan-European dietary survey. *EFSA Journal*, 2009 7(12): 1435.

Journal proceedings

- **FAO.** 2002. Measurement and Assessment of Food Deprivation and Undernutrition International Scientific Symposium, 26–28 June 2002, Rome. Food and Agriculture Organization of the United Nations.
- **FAO.** 2012. International Scientific Symposium on Food and Nutrition Security information: from Valid Measurement to Effective Decision Making 17–19 January 2012, Rome. Food and Agriculture Organization of the United Nations.
- **Tee, E.S., Dop, M.C. and Winichagoon, P.** 2004. Proceedings of the Workshop on Food-Consumption Surveys in Developing Countries: Future challenges. *Food and Nutrition Bulletin*, 25(4) 407–427.

Interactive tools to guide method selection

- Dietary Assessment Primer, United States National Cancer Institute.
- Diet and physical activity measurement toolkit, Medical Research Council.

Other

- TADA, Technology assisted dietary assessment
- Learn More about Technology in Dietary Assessment, United States National Cancer Institute.
- ASA 24, Automatic Self-administered 24-hour Recall, United States National Cancer Institute.
- The International Household Survey Network
- Register of Validated Short Dietary Assessment Instruments, United States National Cancer Institute.
- Dietary Assessment Calibration/Validation Register, United States National Cancer Institute.
- Assessment of the Reliability and Relevance of the Food Data Collected in National Household Consumption and Expenditure Surveys, International Household Survey Network.

- **FAO.** 2001. Food balance sheets: a handbook Rome. Food and Agriculture Organization of the United Nations.
- Dietary Assessment Research Resources, Epidemiology and Genomics Research Program, United States National Cancer Institute.
- Measurement Error Webinar Series, United States National Cancer Institute.
- **Smith, L.S. and Subandoro, A.** 2007. Measuring Food Security Using Household Expenditure Surveys. Food Security in Practice technical guide series. Washington, DC, International Food Policy Research Institute (IFPRI).
- **Deaton, A.** 1997. The analysis of household surveys: a microeconomic approach to development policy. Washington, DC, The World Bank.
- **FAO.** 2014. Guidelines for assessing nutrition-related Knowledge, Attitudes and Practices Rome. Food and Agriculture Organization of the United Nations. **FAO.** 2015. Guidelines on the collection of information on food processing through food consumption surveys Rome. Food and Agriculture Organization of the United Nations.
- STROBE-nut: An extension of the STROBE statement for nutritional epidemiology
- Nutritools developed through DIET@NET (DIETary Assessment Tools NETwork)
- **Cassidy, C. M.** 1994. Walk a mile in my shoes: culturally sensitive food-habit research. *American Journal of Clinical Nutrition*, 59 (1 Suppl).



©FAO/Marco Salustro

7

REFERENCES

- Abdulla M., Andersson I., Asp N., Berthelsen K., Birkhed D., Dencker I., Johansson C., Jägerstad M., Kolar K., Nair B. (1981) Nutrient intake and health status of vegans. Chemical analyses of diets using the duplicate portion sampling technique. *The American Journal of Clinical Nutrition* 34: 2464–2477.
- Acham H., Oldewage-Theron W., Egal A.A. (2012) Dietary diversity, micronutrient intake and their variation among black women in informal settlements in South Africa: a cross-sectional study. *Intern. J. Nutr. Metab* 4: 24–39.
- Albuquerque F.C.A., Bueno N.B., Clemente A.P.G., Ferriolli E., Florêncio T.M.M.T., Hoffman D., Sawaya A.L. (2015) Association between Adult Stature and Energy Expenditure in Low-Income Women from Northeastern Brazil. *Plos One* 10: e0131891.
- Alemayehu A., Abebe Y., Gibson R. (2011) A 24-h recall does not provide a valid estimate of absolute nutrient intakes for rural women in southern Ethiopia. *Nutrition* 27: 919–924.
- Andrissi L., Mottini G., Sebastiani V., Boldrini L., Giuliani A. (2013) Dietary habits and growth: an urban/rural comparison in the Andean region of Apurimac, Peru. *Annali dell'Istituto Superiore di Sanità* 49: 340–346.
- Arimond M., Ruel M.T. (2004) Dietary Diversity Is Associated with Child Nutritional Status: Evidence from 11 Demographic and Health Surveys. *The Journal of Nutrition* 134: 2579–2585.
- Arimond M., Wiesmann D., Becquey E., Carriquiry A., Daniels M.C., Deitchler M., Fanou-Fogny N., Joseph M.L., Kennedy G., Martin-Prevel Y., Torheim L.E. (2010) Simple Food Group Diversity Indicators Predict Micronutrient Adequacy of Women's Diets in 5 Diverse, Resource-Poor Settings. *The Journal of Nutrition* 140: 2059S–2069S. DOI: 10.3945/jn.110.123414.
- Arsenault J., Hijmans R., Brown K. (2015) *Estimating the Micronutrient Adequacy of the Food Supply Using Food Balance Sheets for Targeted Nutrition-sensitive Agricultural Approaches: Example of Bangladesh*. FANTA
- Arsenault J.E., Yakes E.A., Islam M.M., Hossain M.B., Ahmed T., Hotz C., Lewis B., Rahman A.S., Jamil K.M., Brown K.H. (2013) Very Low Adequacy of Micronutrient Intakes by Young Children and Women in Rural Bangladesh Is Primarily Explained by Low Food Intake and

- Limited Diversity. *The Journal of Nutrition* 143: 197–203. DOI: 10.3945/jn.112.169524.
- Baranowski T. (2013) 24-hour Recall and Diet Record Methods. In: Willet W. Third ed. *Nutrition Epidemiology*. Oxford University Press. Third ed.
- Barr S.I., Murphy S.P., Poos M.I. (2002) Interpreting and using the dietary references intakes in dietary assessment of individuals and groups. *Journal of the American Dietetic Association* 102: 780–788.
- Basiotis P.P., Welsh S.O., Cronin F.J., Kelsay J.L., Mertz W. (1987) Number of days of food intake records required to estimate individual and group nutrient intakes with defined confidence. *The Journal of Nutrition* 117: 1638–41.
- Beasley J.M., Davis A., Riley W.T. (2009) Evaluation of a web-based, pictorial diet history questionnaire. *Public Health Nutrition* 12: 651–659.
- Beaton G.H. (1994) Approaches to analysis of dietary data: relationship between planned analyses and choice of methodology. *The American Journal of Clinical Nutrition* 59: 253S–261S.
- Beaton G.H., Burema J., Ritenbaugh C. (1997) Errors in the interpretation of dietary assessments. *The American Journal of Clinical Nutrition* 65: 1100S–1107S.
- Benefice E., Monroy S.L., Jiménez S., López R. (2006) Nutritional status of Amerindian children from the Beni River (lowland Bolivia) as related to environmental, maternal and dietary factors. *Public Health Nutrition* 9: 327–335.
- Bermudez O.I., Lividini K., Smitz M.-F., Fiedler J.L. (2012) Estimating micronutrient intakes from Household Consumption and Expenditures Surveys (HCES): an example from Bangladesh. *Food and Nutrition Bulletin* 33: S208–S213.
- Bingham S., Gill C., Welch A., Day K., Cassidy A., Khaw K., Sneyd M., Key T., Roe L., Day N. (1994) Comparison of dietary assessment methods in nutritional epidemiology: weighed records v. 24 h recalls, food-frequency questionnaires and estimated-diet records. *British Journal of Nutrition* 72: 619–643.
- Bingham S., Cassidy A., Cole T., Welch A., Runswick S., Black A., Thurnham D., Bates C., Khaw K., Key T. (1995) Validation of weighed records and other methods of dietary assessment using the 24 h urine nitrogen technique and other biological markers. *British Journal of Nutrition* 73: 531–550.
- Bingham S.A., Day N.E. (1997) Using biochemical markers to assess the validity of prospective dietary assessment methods and the effect of energy adjustment. *The American Journal of Clinical Nutrition* 65: 1130S–1137S.
- Biro G., Hulshof K., Ovesen L., Amorim C.J. (2002) Selection of methodology to assess food intake. *European Journal of Clinical Nutrition* 56: S25–32.
- Black A., Bingham S., Johansson G., Coward W. (1997) Validation of dietary intakes of protein and energy against 24 hour urinary N and DLW energy expenditure in middle-aged women, retired men and post-obese subjects: comparisons with validation against presumed energy requirements. *European Journal of Clinical Nutrition* 51: 405–413.
- Black A.E. (2000) Critical evaluation of energy intake using the Goldberg cut-off for energy intake: basal metabolic rate. A practical guide to its calculation, use and limitations. *International Journal of Obesity & Related Metabolic Disorders* 24.
- Bland J.M., Altman D. (1986) Statistical methods for assessing agreement between two methods of clinical measurement. *The Lancet* 327: 307–310.

- Blanton C.A., Moshfegh A.J., Baer D.J., Kretsch M.J. (2006) The USDA Automated Multiple-Pass Method accurately estimates group total energy and nutrient intake. *The Journal of Nutrition* 136: 2594–2599.
- Block G., Hartman A.M. (1989) Issues in reproducibility and validity of dietary studies. *The American Journal of Clinical Nutrition* 50: 1133–1138.
- Bonifacj C., Gerber M., Scali J., Daires J. (1997) Comparison of dietary assessment methods in a southern French population: use of weighed records, estimated-diet records and a food-frequency questionnaire. *European Journal of Clinical Nutrition* 51: 217–231.
- Bountziouka V., Panagiotakos D. (2010) Statistical methods used for the evaluation of reliability and validity of nutrition assessment tools used in medical research. *Current Pharmaceutical Design* 16: 3770–3775.
- Boushey C.J., Harray A.J., Kerr D.A., Schap T.E., Paterson S., Aflague T., Ruiz M.B., Ahmad Z., Delp E.J. (2015) How Willing Are Adolescents to Record Their Dietary Intake? The Mobile Food Record. *JMIR mHealth and uHealth* 3: e47.
- Brink P.J. (1991) Issues of reliability and validity. Qualitative nursing research: A contemporary dialogue: 164–186.
- Burke B.S. (1947) The dietary history as a tool in research. *Journal of the American Dietetic Association* 23: 1041–1046.
- Burrows T.L., Martin R.J., Collins C.E. (2010) A systematic review of the validity of dietary assessment methods in children when compared with the method of doubly labelled water. *Journal of the American Dietetic Association* 110: 1501–1510.
- Cade J., Thompson R., Burley V., Warm D. (2002) Development, validation and utilisation of food-frequency questionnaires—a review. *Public Health Nutrition* 5: 567–587.
- Campbell R.K., Talegawkar S.A., Christian P., LeClerq S.C., Khatry S.K., Wu L.S., West Jr K.P. (2014) Seasonal dietary intakes and socioeconomic status among women in the Terai of Nepal. *Journal of health, population, and nutrition* 32: 198.
- Carnethon M.R., Loria C.M., Hill J.O., Sidney S., Savage P.J., Liu K. (2004) Risk Factors for the Metabolic Syndrome The Coronary Artery Risk Development in Young Adults (CARDIA) study, 1985–2001. *Diabetes care* 27: 2707–2715.
- Carroll R.J., Freedman L.S., Kipnis V. (1998) Measurement error and dietary intake, Mathematical Modeling in Experimental Nutrition, Springer. pp. 139–145.
- Casperson S.L., Sieling J., Moon J., Johnson L., Roemmich J.N., Whigham L. (2015) A Mobile Phone Food Record App to Digitally Capture Dietary Intake for Adolescents in a Free-Living Environment: Usability Study. *JMIR mHealth and uHealth* 3.
- Cassidy C.M. (1994) Walk a mile in my shoes: culturally sensitive food-habit research. *The American Journal of Clinical Nutrition* 59: 190S-197S.
- Chae J., Woo I., Kim S., Maciejewski R., Zhu F., Delp E.J., Boushey C.J., Ebert D.S. (2011) Volume estimation using food specific shape templates in mobile image-based dietary assessment, IS&T/SPIE *Electronic Imaging, International Society for Optics and Photonics*. pp. 78730K-78730K-8.
- Chambers R. (1994) Participatory rural appraisal (PRA): analysis of experience. *World development* 22: 1253–1268.
- Charlton K.E., Steyn K., Levitt N.S., Jonathan D., Zulu J.V., Nel J.H. (2008) Development and validation of a short questionnaire to assess sodium intake. *Public Health Nutrition* 11: 83–94.

- Chinnock A. (2006) Validation of an estimated food record. *Public Health Nutrition* 9: 934–941.
- Clausen T., Charlton K.E., Gobotswang K.S.M., Holmboe-Ottesen G. (2005) Predictors of food variety and dietary diversity among older persons 7-in Botswana. *Nutrition* 21: 86–95. DOI: <http://dx.doi.org/10.1016/j.nut.2004.09.012>.
- Corsi A., Lorens A. (2008) A participatory assessment of dietary patterns and food behaviour in Pohnpei, Federated States of Micronesia. *Asia Pacific Journal of Clinical Nutrition* 17: 309.
- Coulibaly A., O'Brien H.T., Galibois I. (2009) Validation of an FFQ to assess dietary protein intake in type 2 diabetic subjects attending primary health-care services in Mali. *Public Health Nutrition* 12: 644–650.
- Coward W.A. (1998) Contributions of the doubly labelled water method to studies of energy balance in the Third World. *The American Journal of Clinical Nutrition* 68: 962S–969S.
- da Silva R., Bach-Faig A., Quintana B.R., Buckland G., de Almeida M.D.V., Serra-Majem L. (2009) Worldwide variation of adherence to the Mediterranean diet, in 1961–1965 and 2000–2003. *Public Health Nutrition* 12: 1676–1684.
- Dehghan M., Jaramillo P.L., Dueñas R., Anaya L.L., Garcia R.G., Zhang X., Islam S., Merchant A.T. (2012) Development and validation of a quantitative food frequency questionnaire among rural-and urban-dwelling adults in Colombia. *Journal of Nutrition Education and Behavior* 44: 609–613.
- Del Gobbo L.C., Khatibzadeh S., Imamura F., Micha R., Shi P., Smith M., Myers S.S., Mozaffarian D. (2015) Assessing global dietary habits: a comparison of national estimates from the FAO and the Global Dietary Database. *The American Journal of Clinical Nutrition* 101: 1038–1046.
- Dennis B., Stamler J., Buzzard M., Conway R., Elliott P., Moag-Stahlberg A., Okayama A., Okuda N., Robertson C., Robinson F. (2003) INTERMAP: the dietary data—process and quality control. *Journal of human hypertension* 17: 609–622.
- Dodd K.W., Guenther P.M., Freedman L.S., Subar A.F., Kipnis V., Midthune D., Tooze J.A., Krebs-Smith S.M. (2006) Statistical methods for estimating usual intake of nutrients and foods: a review of the theory. *Journal of the American Dietetic Association* 106: 1640–1650.
- Dop M.-C., Milan C., Milan C., N'Diaye A. (1994) The 24-hour recall for Senegalese weanlings: a validation exercise. *European Journal of Clinical Nutrition* 48: 643–653.
- Dop M.C., Mistura L., Pereira C., Martinez C., Cardoso E. (2012) Using HIES data to explore dietary and nutrient intake patterns: a case study of the nutrition transition of Cape Verde. *Food & Nutrition Bulletin*.
- Drewnowski A. (2001) Diet image: a new perspective on the food-frequency questionnaire. *Nutrition Reviews* 59: 370.
- EFSA. (2014) *Guidance on the EU Menu methodology*. European Food Safety Authority. (available at: http://www.efsa.europa.eu/sites/default/files/scientific_output/files/main_documents/3944.pdf Accessed 23 October 2015).
- Fahmida U., Kolopaking R., Santika O., Sriani S., Umar J., Htet M.K., Ferguson E. (2015) Effectiveness in improving knowledge, practices, and intakes of “key problem nutrients” of a complementary feeding intervention developed by using linear programming: experience in Lombok, Indonesia. *The American Journal of Clinical Nutrition* 101: 455–461.
- FAO. (2001) *Food balance sheets. A handbook*. <http://www.fao.org/3/a-x9892e.pdf> Accessed February 2016.

- FAO. (2010) *Guidelines for Measuring Household and Individual Dietary Diversity*. Food and Agricultural Organization. Rome, Italy. http://www.fao.org/fileadmin/user_upload/wa_workshop/docs/FAO-guidelines-dietary-diversity2011.pdf.
- FAO. (2015) FAOSTAT, Food Balances. <http://faostat3.fao.org/home/E>. Accessed May 2016
- FAO/FANTA/IRD. (2014) Meeting to Reach Consensus on a Global Dietary Diversity Indicator for Women, Washington DC. <http://www.fantaproject.org/sites/default/files/resources/WDDP-Meeting-Report-Oct2014.pdf>.
- FAO/FHI360. (2016) Minimum Dietary Diversity for Women: A Guide for Measurement. Rome: FAO. Rome, Italy. <http://www.fao.org/3/a-i5486e.pdf>
- FAO/IFAD/WFP. (2015) *The State of Food Insecurity in the World 2015*. Meeting the 2015 international hunger targets: taking stock of uneven progress. Rome, Italy.
- Fiedler J.L., Martin-Prével Y., Moursi M. (2013) Relative costs of 24-hour recall and household consumption and expenditures surveys for nutrition analysis. *Food and Nutrition Bulletin* 34: 318–330.
- Fiedler J.L., Lividini K., Bermudez O.I., Smitz M.-F. (2012) Household Consumption and Expenditures Surveys (HCES): a primer for food and nutrition analysts in low-and middle-income countries. *Food and Nutrition Bulletin* 33: S170—S184.
- Fontana J.M., Higgins J.A., Schuckers S.C., Bellisle F., Pan Z., Melanson E.L., Neuman M.R., Sazonov E. (2015) Energy intake estimation from counts of chews and swallows. *Appetite* 85: 14–21. DOI: <http://dx.doi.org/10.1016/j.appet.2014.11.003>.
- Forster H., Walsh M.C., Gibney M.J., Brennan L., Gibney E.R. (2015) Personalised nutrition: the role of new dietary assessment methods. *The Proceedings of the Nutrition Society*: 1–10.
- Fowles E.R., Gentry B. (2008) The feasibility of personal digital assistants (PDAs) to collect dietary intake data in low-income pregnant women. *Journal of Nutrition Education and Behavior* 40: 374–377.
- Freedman L.S., Commins J.M., Moler J.E., Arab L., Baer D.J., Kipnis V., Midthune D., Moshfegh A.J., Neuhauser M.L., Prentice R.L. (2014) Pooled results from 5 validation studies of dietary self-report instruments using recovery biomarkers for energy and protein intake. *American Journal of Epidemiology*: kwu116.
- Freedman L.S., Commins J.M., Moler J.E., Willett W., Tinker L.F., Subar A.F., Spiegelman D., Rhodes D., Potischman N., Neuhauser M.L. (2015) Pooled results from 5 validation studies of dietary self-report instruments using recovery biomarkers for potassium and sodium intake. *American Journal of Epidemiology*: kwu325.
- Fujita M., Lo Y.-j., Baranski J.R. (2012) Dietary diversity score is a useful indicator of vitamin A status of adult women in Northern Kenya. *American Journal of Human Biology* 24: 829–834. DOI: 10.1002/ajhb.22327.
- Fukumoto A., Asakura K., Murakami K., Sasaki S., Okubo H., Hirota N., Notsu A., Todoriki H., Miura A., Fukui M. (2013) Within-and between-individual variation in energy and nutrient intake in Japanese adults: effect of age and sex differences on group size and number of records required for adequate dietary assessment. *Journal of Epidemiology* 23: 178–186.
- Gemming L., Utter J., Mhurchu C.N. (2015) Image-Assisted Dietary Assessment: A Systematic Review of the Evidence. *Journal of the Academy of Nutrition and Dietetics* 115: 64–77.

- Gemming L., Doherty A., Kelly P., Utter J., Ni Mhurchu C. (2013) Feasibility of a SenseCam-assisted 24-h recall to reduce under-reporting of energy intake. *European Journal of Clinical Nutrition* 67: 1095–1099. DOI: 10.1038/ejcn.2013.156.
- Gewa C.A., Murphy S.P., Neumann C.G. (2009) A comparison of weighed and recalled intakes for schoolchildren and mothers in rural Kenya. *Public Health Nutrition* 12: 1197–1204.
- Gibson R.S., Cavalli-Sforza. (2012) Using reference nutrient density goals with food balance sheet data to identify likely micronutrient deficits for fortification planning in countries in the Western Pacific region. *Food and Nutrition Bulletin* 33: S214–S220.
- Gibson R.S., Scythes C.A. (1982) Trace element intakes of women. *British Journal of Nutrition* 48: 241–248.
- Gibson R.S., Ferguson E.L. (2008) An interactive 24-hour recall for assessing the adequacy of iron and zinc intakes in developing countries. Washington, DC: IFPRI and CIAT.
- Gibson R.S., Wawer A.A., Fairweather-Tait S.J., Hurst R., Young S.D., Broadley M.R., Chilimba A.D., Ander E.L., Watts M.J., Kalimbira A. (2015) Dietary iron intakes based on food composition data may underestimate the contribution of potentially exchangeable contaminant iron from soil. *Journal of Food Composition and Analysis* 40: 19–23.
- Gittelsohn J. (1989) *Intrahousehold food distribution in rural Nepal*, Doctoral dissertation, University of Connecticut. University Microfilms, Ann Arbor, MI.
- Gonzalez C. (1997) Relative validity and reproducibility of a diet history questionnaire in Spain. III. Biochemical markers. *International Journal of Epidemiology* (United Kingdom).
- Hagfors L., Nilsson I., Sköldstam L., Johansson G. (2005) Fat intake and composition of fatty acids in serum phospholipids in a randomized, controlled, Mediterranean dietary intervention study on patients with rheumatoid arthritis. *Nutrition & Metabolism* 2: 1.
- Hansen A.W., Christensen D.L., Larsson M.W., Eis J., Christensen T., Friis H., Mwaniki D.L., Kilonzo B., Boit M.K., Borch-Johnsen K. (2011) Dietary patterns, food and macronutrient intakes among adults in three ethnic groups in rural Kenya. *Public Health Nutrition* 14: 1671–1679.
- Harrison G.G. (2004) Methodologic considerations in descriptive food-consumption surveys in developing countries. *Food and Nutrition Bulletin* 25: 415–419.
- Hartman A.M., Brown C.C., Palmgren J., Pietinen P., Verkasalo M., Myer D., Virtamo J. (1990) Variability in nutrient and food intakes among older middle-aged men Implications for design of epidemiologic and validation studies using food recording. *American Journal of Epidemiology* 132: 999–1012.
- Hedrick V.E., Dietrich A.M., Estabrooks P.A., Savla J., Serrano E., Davy B.M. (2012) Dietary biomarkers: advances, limitations and future directions. *Nutrition Journal* 11: 1.
- Herforth A. (2015) Access to adequate nutritious food: new indicators to track progress and inform action, in Sahn D., ed. *The Fight Against Hunger and Malnutrition: The Role of Food, Agriculture, and Targeted Policies*, Oxford University Press: 139.
- Hernández-Avila M., Romieu I., Parra S., Hernández-Avila J., Madrigal H., Willett W. (1998) Validity and reproducibility of a food frequency questionnaire to assess dietary intake of women living in Mexico City. *Salud Pública de México* 40: 133–140.
- Hernández-Cordero S., Popkin B.M. (2015) Impact of a Water Intervention on Sugar-Sweetened Beverage Intake Substitution by Water: A Clinical Trial in Overweight and Obese Mexican

- Women. *Annals of Nutrition and Metabolism* 66: 22–25.
- Hill R., Davies P. (2001) The validity of self-reported energy intake as determined using the doubly labelled water technique. *British Journal of Nutrition* 85: 415–430.
- Hoddinott J., Yohannes Y. (2002) *Dietary diversity as a food security indicator. Food consumption and nutrition division discussion paper 136*. IFPRI
- Holm L., Lund T.B., Niva M. (2015) Eating practices and diet quality: a population study of four Nordic countries. *European Journal of Clinical Nutrition*. DOI: 10.1038/ejcn.2015.61.
- Holmes B., Nelson M. (2009) The strengths and weaknesses of dietary survey methods in materially deprived households in England: a discussion paper. *Public Health Nutrition* 12: 1157–1164.
- Holmes B., Dick K., Nelson M. (2008) A comparison of four dietary assessment methods in materially deprived households in England. *Public Health Nutrition* 11: 444–456.
- Hurst R., Siyame E.W., Young S.D., Chilimba A.D., Joy E.J., Black C.R., Ander E.L., Watts M.J., Chilima B., Gondwe J., Kang'ombe D., Stein A.J., Fairweather-Tait S.J., Gibson R.S., Kalimpira A.A., Broadley M.R. (2013) Soil-type influences human selenium status and underlies widespread selenium deficiency risks in Malawi. *Scientific Reports* 3.
- Hutanasu C., Sfarti C., Trifan A., Hutanasu M., Stanciu C. (2009) [Aflatoxin contamination of food: additional risk factor for chronic liver diseases]. *Rev Med Chir Soc Med Nat Iasi* 113: 1061–5.
- Illner A., Freisling H., Boeing H., Huybrechts I., Crispim S., Slimani N. (2012) Review and evaluation of innovative technologies for measuring diet in nutritional epidemiology. *International Journal of Epidemiology* 41: 1187–1203.
- Irwig M.S., El-Sohehy A., Baylin A., Rifai N., Campos H. (2002) Frequent intake of tropical fruits that are rich in β -cryptoxanthin is associated with higher plasma β -cryptoxanthin concentrations in Costa Rican adolescents. *The Journal of Nutrition* 132: 3161–3167.
- Jackson M., Walker S., Cade J., Forrester T., Cruickshank J., Wilks R. (2001) Reproducibility and validity of a quantitative food-frequency questionnaire among Jamaicans of African origin. *Public Health Nutrition* 4: 971–980.
- Jackson M.D., Motswagole B.S., Kwape L.D., Kobue-Lekalake R.I., Rakgantswana T.B., Mongwaketse T., Mokotedi M., Jackson-Malete J. (2013) Validation and reproducibility of an FFQ for use among adults in Botswana. *Public Health Nutrition* 16: 1995–2004.
- Jariseta Z.R., Dary O., Fiedler J.L., Franklin N. (2012) Comparison of estimates of the nutrient density of the diet of women and children in Uganda by Household Consumption and Expenditures Surveys (HCES) and 24-hour recall. *Food and Nutrition Bulletin* 33: S199–S207.
- Jilcott S.B., Keyserling T.C., Samuel-Hodge C.D., Johnston L.F., Gross M.D., Ammerman A.S. (2007) Validation of a brief dietary assessment to guide counselling for cardiovascular disease risk reduction in an underserved population. *Journal of the American Dietetic Association* 107: 246–255.
- Johnson R.K., Goran M.I., Poehlman E.T. (1994) Correlates of over- and underreporting of energy intake in healthy older men and women. *The American Journal of Clinical Nutrition* 59: 1286–1290.
- Kabagambe E.K., Baylin A., Allan D.A., Siles X., Spiegelman D., Campos H. (2001) Application of the method of triads to

- evaluate the performance of food frequency questionnaires and biomarkers as indicators of long-term dietary intake. *American Journal of Epidemiology* 154: 1126–1135.
- Kennedy G., Fanou-Fogny N., Seghieri C., Arimond M., Koreissi Y., Dossa R., Kok F.J., Brouwer I.D. (2010) Food Groups Associated with a Composite Measure of Probability of Adequate Intake of 11 Micronutrients in the Diets of Women in Urban Mali. *The Journal of Nutrition* 140: 2070S-2078S. DOI: 10.3945/jn.110.123612.
- Kennedy G.L., Pedro M.R., Seghieri C., Nantel G., Brouwer I. (2007) Dietary Diversity Score Is a Useful Indicator of Micronutrient Intake in Non-Breast-Feeding Filipino Children. *The Journal of Nutrition* 137: 472–477.
- Khanna N., Boushey C.J., Kerr D., Okos M., Ebert D.S., Delp E.J. (2010) An overview of the technology assisted dietary assessment project at Purdue University, Multimedia (ISM), 2010 IEEE International Symposium on IEEE. pp. 290–295.
- Kigutha H.N. (1997) Assessment of dietary intake in rural communities in Africa: experiences in Kenya. *The American Journal of Clinical Nutrition* 65: 1168S-1172S.
- Kikunaga S., Tin T., Ishibashi G., Wang D.-H., Kira S. (2007) The application of a handheld personal digital assistant with camera and mobile phone card (Wellnavi) to the general population in a dietary survey. *Journal of Nutritional Science and Vitaminology* 53: 109–116.
- Kirkpatrick S.I., Subar A.F., Douglass D., Zimmerman T.P., Thompson F.E., Kahle L.L., George S.M., Dodd K.W., Potischman N. (2014) Performance of the Automated Self-Administered 24-hour Recall relative to a measure of true intakes and to an interviewer-administered 24-h recall. *The American Journal of Clinical Nutrition* 100: 233–240.
- Klipstein-Grobusch K.d., Den Breeijen J., Goldbohm R., Geleijnse J., Hofman A., Grobbee D., Witteman J. (1998) Dietary assessment in the elderly: validation of a semiquantitative food frequency questionnaire. *European Journal of Clinical Nutrition* 52: 588–596.
- Kristal A.R., Peters U., Potter J.D. (2005) Is it time to abandon the food frequency questionnaire? *Cancer Epidemiology Biomarkers & Prevention* 14: 2826–2828.
- Kristal A.R., Feng Z., Coates R.J., Oberman A., George V. (1997) Associations of Race/Ethnicity, Education, and Dietary Intervention with the Validity and Reliability of a Food Frequency Questionnaire The Women’s Health Trial Feasibility Study in Minority Populations. *American Journal of Epidemiology* 146: 856–869.
- Landais E., Gartner A., Bour A., McCullough F., Delpeuch F., Holdsworth M. (2014) Reproducibility and relative validity of a brief quantitative food frequency questionnaire for assessing fruit and vegetable intakes in North-African women. *Journal of Human Nutrition and Dietetics* 27: 152–159.
- Lassen A.D., Poulsen S., Ernst L., Andersen K.K., Biloft-Jensen A., Tetens I. (2010) Evaluation of a digital method to assess evening meal intake in a free-living adult population. *Food & nutrition research* 54.
- Lazarte C.E., Mouratidou T., Kurbanov S., Qulov A., Lee W. (2015) *Project Report on the Integration of the Minimum Dietary Diversity-Women (MDD-W) into the Household Budget Survey (HBS) in Tajikistan*, 2015 FAO, Rome, Italy. http://www.fao.org/fileadmin/user_upload/nutrition/docs/assessment/Report_MDD-W_Tajikistan_Nov2015.pdf.
- Lazarte C.E., Encinas M.E., Alegre C., Granfeldt Y. (2012) Validation of digital photographs, as a tool in 24-h recall, for the improvement of dietary assessment among rural populations

- in developing countries. *Nutrition Journal* 11: 61.
- Leroy J.L., Ruel M., Frongillo E.A., Harris J., Ballard T.J. (2015) Measuring the Food Access Dimension of Food Security A Critical Review and Mapping of Indicators. *Food and Nutrition Bulletin* 36: 167–195.
- Leung S., Chan S., Lui S., Lee W., Davies D. (2000) Growth and nutrition of Hong Kong children aged 0–7 years. *Journal of Paediatrics and Child Health* 36: 56–65.
- Liberato S.C., Kearns T., Ward F., Brimblecombe J. (2015) Use of electronic visual recording to aid assessment of dietary intake of Australian Aboriginal children living in remote communities. *Australian and New Zealand Journal of Public Health: n/a-n/a*. DOI: 10.1111/1753–6405.12373.
- Lightowler H.J., Davies G.J. (1998) Iodine intake and iodine deficiency in vegans as assessed by the duplicate-portion technique and urinary iodine excretion. *British Journal of Nutrition* 80: 529–35.
- Lim S., Sohn W., Burt B.A., Sandretto A.M., Kolker J.L., Marshall T.A., Ismail A.I. (2008) Cariogenicity of soft drinks, milk and fruit juice in low-income african-american children: a longitudinal study. *The Journal of the American Dental Association* 139: 959–967.
- Liu S., Manson J.E., Stampfer M.J., Holmes M.D., Hu F.B., Hankinson S.E., Willett W.C. (2001) Dietary glycemic load assessed by food-frequency questionnaire in relation to plasma high-density-lipoprotein cholesterol and fasting plasma triacylglycerols in postmenopausal women. *The American Journal of Clinical Nutrition* 73: 560–566.
- Livingstone M.B.E., Black A.E. (2003) Markers of the validity of reported energy intake. *The Journal of Nutrition* 133: 895S-920S.
- Lombard M.J., Steyn N.P., Charlton K.E., Senekal M. (2015) Application and interpretation of multiple statistical tests to evaluate validity of dietary intake assessment methods. *Nutrition Journal* 14: 1.
- Malde M., Zerihun L., Julshamn K., Bjorvatn K. (2004) Fluoride, calcium and magnesium intake in children living in a high-fluoride area in Ethiopia. Intake through food. *International Journal of Paediatric Dentistry* 14: 167–174.
- Marr J.W. (1971) Individual dietary surveys: purposes and methods. *World Review of Nutrition and Dietetics* 13: 105–164.
- Martin-Moreno JM, Boyle P, Gorgojo L, Maisonneuve P, Fernandez-Rodriguez JC, Salvini S, W. W. (1993) Development and validation of a food frequency questionnaire in Spain. *International Journal of Epidemiology* 22: 512–519.
- Martin-Prével Y., Allemand P., Wiesmann D., Arimond M., Ballard T., Deitchler M., Dop MC., Kennedy G., Lee WTK., M. M. (2015) FAO, IRD. *Moving Forward on Choosing a Standard Operational Indicator of Women's Dietary Diversity*. FAO, Food and Agriculture Organization. Rome, Italy Rome, Italy. <http://www.fao.org/documents/card/en/c/678ab9d4-e7a8-4388-9f9f-1c709ea47752/>
- Martin-Prevel Y., Arimond M., Allemand P., Wiesmann D., Ballard T.J., Deitchler M., Dop MC., Kennedy G., Lartey A., Lee WTK, Moursi M. (2017) Development of a dichotomous indicator for population-level assessment of dietary diversity of women of reproductive age. *Current Developments in Nutrition* 1:e001701.
- Martin-Moreno J.M., Willett W.C., Gorgojo L., Banegas J.R., Rodriguez-Artalejo F., Fernandez-Rodriguez J.C., Maisonneuve P, Boyle P. (1994) Dietary fat, olive oil intake and breast cancer risk. *International Journal of Cancer* 58: 774–780.

- Martin C.K., Han H., Coulon S.M., Allen H.R., Champagne C.M., Anton S.D. (2009) A novel method to remotely measure food intake of free-living individuals in real time: the remote food photography method. *British Journal of Nutrition* 101: 446–456.
- Martin C.K., Nicklas T., Gunturk B., Correa J.B., Allen H.R., Champagne C. (2014) Measuring food intake with digital photography. *Journal of Human Nutrition and Dietetics*: the official journal of the British Dietetic Association 27: 72–81. DOI: 10.1111/jhn.12014.
- Martin C.K., Correa J.B., Han H., Allen H.R., Rood J.C., Champagne C.M., Gunturk B.K., Bray G.A. (2012) Validity of the Remote Food Photography Method (RFPM) for Estimating Energy and Nutrient Intake in Near Real-Time. *Obesity* 20: 891–899.
- Mattei J., Malik V., Wedick N.M., Hu F.B., Spiegelman D., Willett W.C., Campos H. (2015) Reducing the global burden of type 2 diabetes by improving the quality of staple foods: The Global Nutrition and Epidemiologic Transition Initiative. *Globalization and health* 11: 1.
- McClung H.L., Sigrist L.D., Smith T.J., Karl J.P., Rood J.C., Young A.J., Bathalon G.P. (2009) Monitoring energy intake: a hand-held personal digital assistant provides accuracy comparable to written records. *Journal of the American Dietetic Association* 109: 1241–1245.
- McCullough M.L., Feskanich D., Stampfer M.J., Giovannucci E.L., Rimm E.B., Hu F.B., Spiegelman D., Hunter D.J., Colditz G.A., Willett W.C. (2002) *Diet quality* and major chronic disease risk in men and women: moving toward improved dietary guidance. *The American Journal of Clinical Nutrition* 76: 1261–1271.
- McDonald A., Van Horn L., Slattery M., Hilner J., Bragg C., Caan B., Jacobs Jr D., Liu K., Hubert H., Gernhofer N. (1991) The CARDIA dietary history: development, implementation, and evaluation. *Journal of the American Dietetic Association* 91: 1104–1112.
- McKeown N.M., Day N.E., Welch A.A., Runswick S.A., Luben R.N., Mulligan A.A., McTaggart A., Bingham S.A. (2001) Use of biological markers to validate self-reported dietary intake in a random sample of the European Prospective Investigation into Cancer United Kingdom Norfolk cohort. *The American Journal of Clinical Nutrition* 74: 188–196.
- Mensink G., Haftenberger M., Thamm M. (2001) Original Communications-Validity of DISHES 98, a computerised dietary history interview: Energy and macronutrient intake. *European Journal of Clinical Nutrition* 55: 409–417.
- Merchant A.T., Dehghan M., Chifamba J., Terera G., Yusuf S. (2005) Nutrient estimation from an FFQ developed for a black Zimbabwean population. *Nutrition Journal* 4: 1.
- Moltedo A., Troubat N., Lokshin M., Sajaia Z. (2014) *Analyzing food security using household survey data: streamlined analysis with ADePT software*. World Bank Publications.
- Murphy S., Ruel M., Carriquiry A. (2012) Should Household Consumption and Expenditures Surveys (HCES) be used for nutritional assessment and planning? *Food and Nutrition Bulletin* 33: S235–S241.
- Naska A., Berg M.-A., Cuadrado C., Freisling H., Gedrich K., Gregorič M., Kelleher C., Leskova E., Nelson M., Pace L. (2009) Food balance sheet and household budget survey dietary data and mortality patterns in Europe. *British Journal of Nutrition* 102: 166–171.
- Nelson M. (1997) 8. The validation of dietary assessment. In: Margetts BM, Nelson M (eds.). *Design concepts in nutritional epidemiology*. Second edition, Oxford University Press. New York: 241–272.

- Ngo J., Engelen A., Molag M., Roesle J., García-Segovia P., Serra-Majem L. (2009) A review of the use of information and communication technologies for dietary assessment. *British Journal of Nutrition* 101: S102–S112. DOI: doi: 10.1017/S0007114509990638.
- Nurul-Fadhilah A., Teo P.S., Foo L.H. (2012) Validity and reproducibility of a food frequency questionnaire (FFQ) for dietary assessment in Malay adolescents in Malaysia. *Asia Pacific Journal of Clinical Nutrition* 21: 97.
- Nyambose J., Koski K.G., Tucker K.L. (2002) High intra/interindividual variance ratios for energy and nutrient intakes of pregnant women in rural Malawi show that many days are required to estimate usual intake. *The Journal of Nutrition* 132: 1313–1318.
- Omigbodun O.O., Adediran K.I., Akinyemi J.O., Omigbodun A.O., Adedokun B.O., Esan O. (2010) Gender and rural–urban differences in the nutritional status of in-school adolescents in south-western Nigeria. *Journal of biosocial science* 42: 653–676.
- Oninla S., Owa J., Onayade A., Taiwo O. (2007) Comparative study of nutritional status of urban and rural Nigerian school children. *Journal of Tropical Pediatrics* 53: 39–43.
- Ortega R.M., Pérez-Rodrigo C., López-Sobaler A.M. (2015) Dietary assessment methods: dietary records. *Nutr Hosp* 31: 38–45.
- Patterson R.E., Pietinen P. (2004) In: Gibney MJ, Barrie BM, Kearney JM, Arab L, *Public Health Nutrition*. Third ed. Oxford, Wiley-Blackwell Science.
- Pennington J. (2004) Proceedings of the workshop on food-consumption surveys in developing countries: general approaches to estimation of dietary exposure to contaminants. *Food and Nutrition Bulletin* 25: 420–7.
- Pereira R.A., Araujo M.C., Lopes T.d.S., Yokoo E.M. (2010) How many 24-hour recalls or food records are required to estimate usual energy and nutrient intake? *Cadernos de Saude Publica* 26: 2101–2111.
- Perez Rodrigo C., Moran Fagundez L.J., Riobo Servan P., Aranceta Bartrina J. (2015) Screeners and brief assessment methods. *Nutr Hosp* 3: 91–8.
- Persson V., Winkvist A., Ninuk T., Hartini S., Greiner T., Hakimi M., Stenlund H. (2001) Variability in nutrient intakes among pregnant women in Indonesia: implications for the design of epidemiological studies using the 24-h recall method. *The Journal of Nutrition* 131: 325–330.
- Poslusna K., Ruprich J., de Vries J.H., Jakubikova M., van't Veer P. (2009) Misreporting of energy and micronutrient intake estimated by food records and 24 hour recalls, control and adjustment methods in practice. *British Journal of Nutrition* 101: S73–S85.
- Ptomey L.T., Willis E.A., Goetz J.R., Lee J., Sullivan D.K., Donnelly J.E. (2015) Digital photography improves estimates of dietary intake in adolescents with intellectual and developmental disabilities. *Disability and health journal* 8: 146–150.
- Putnam J.J., Allshouse J.E. (1994) *Food Consumption, Prices, and Expenditures, 1970–93* US Department of Agriculture, Economic Research Service.
- Rankin D., Ellis S., Macintyre U., Hanekom S., Wright H.H. (2011) Dietary intakes assessed by 24-h recalls in peri-urban African adolescents: validity of energy intake compared with estimated energy expenditure. *European Journal of Clinical Nutrition* 65: 910–919.
- Rankin D., MacIntyre U.E., Hanekom S.M., Steyn H.S., Wright H.H. (2012) Reproducibility of two, three, four and five 24-hour recalls in peri-urban African adolescents in the North West

- province. *South African Journal of Clinical Nutrition* 25: 27–32.
- Rao S., Kanade A.N., Yajnik C.S., Fall C.H. (2009) Seasonality in maternal intake and activity influence offspring's birth size among rural Indian mothers—Pune Maternal Nutrition Study. *International Journal of Epidemiology* 38: 1094–1103.
- Rasmussen L.B., Matthiessen J., Biloft-Jensen A., Tetens I. (2007) Characteristics of misreporters of dietary intake and physical activity. *Public Health Nutrition* 10: 230–237.
- Rezali F.W., Chin Y.S., Yusof M., Nisak B. (2012) Obesity-related behaviors of Malaysian adolescents: a sample from Kajang district of Selangor state. *Nutrition research and practice* 6: 458–465.
- Rollo M.E., Ash S., Lyons-Wall P., Russell A. (2011) Trial of a mobile phone method for recording dietary intake in adults with type 2 diabetes: evaluation and implications for future applications. *Journal of telemedicine and telecare* 17: 318–323.
- Rose D., Tschirley D. (2003) Predicting dietary intakes with simple food recall information: a case study from rural Mozambique. *European Journal of Clinical Nutrition* 57: 1212–1221.
- Rose D., Chotard S., Oliveira L., Mock N., Libombo M. (2008) A comparative evaluation of dietary indicators used in food consumption assessments of at-risk populations. *Food and Nutrition Bulletin* 29: 113–122.
- Ruel M., Harris J., Cunningham K. (2013) *Diet Quality in Developing Countries*, in: V. R. Preedy, et al. (Eds.), *Diet Quality*, Springer New York. pp. 239–261.
- Ruel M.T. (2003a) Is dietary diversity an indicator of food security or dietary quality? A review of measurement issues and research needs. *Food and Nutrition Bulletin* 24: 231–2.
- Ruel M.T. (2003b) Operationalizing Dietary Diversity: A Review of Measurement Issues and Research Priorities. *The Journal of Nutrition* 133: 3911S–3926S.
- Sanchez-Llaguno S., Neira-Mosquera J., Pérez-Rodríguez F., Moreno-Rojas R. (2013) Preliminary nutritional assessment of the Ecuadorian diet based on a 24-h food recall survey in Ecuador. *Nutr Hosp* 28: 1646–1656.
- Sanz-Gallardo M.I., Guallar E., Van'T Veer P., Longnecker M.P., Strain J., Martin B.C., Kardinaal A.F., Fernández-Crehuet J., Thamm M., Kohlmeier L. (1999) Determinants of p, p'-dichlorodiphenyldichloroethane (DDE) concentration in adipose tissue in women from five European cities. *Archives of Environmental Health: An International Journal* 54: 277–283.
- Savy M., Martin-Prével Y., Traissac P., Eymard-Duvernay S., Delpeuch F. (2006) Dietary diversity scores and nutritional status of women change during the seasonal food shortage in rural Burkina Faso. *The Journal of Nutrition* 136: 2625–2632.
- Scagliusi F.B., Ferrioli E., Lancha A.H. (2006) Underreporting of energy intake in developing nations. *Nutrition Reviews* 64: 319–330.
- Schakel S.F., Buzzard I.M., Gebhardt S.E. (1997) Procedures for estimating nutrient values for food composition databases. *Journal of Food Composition and Analysis* 10: 102–114.
- Schap T., Zhu F., Delp E., Boushey C. (2014) Merging dietary assessment with the adolescent lifestyle. *Journal of Human Nutrition and Dietetics* 27: 82–88.
- Schofield W. (1984) Predicting basal metabolic rate, new standards and review of previous work. *Human nutrition. Clinical nutrition* 39: 5–41.
- Serra-Majem L., MacLean D., Ribas L., Brulé D., Sekula W., Prattala R., Garcia-Closas R., Yngve A., Lalonde M., Petrasovits A. (2003)

- Comparative analysis of nutrition data from national, household, and individual levels: results from a WHO-CINDI collaborative project in Canada, Finland, Poland, and Spain. *Journal of Epidemiology and Community Health* 57: 74–80.
- Shapiro S.S., Wilk M.B. (1965) An analysis of variance test for normality (complete samples). *Biometrika* 52: 591–611.
- Shim J.-S., Oh K., Kim H.C. (2014) Dietary assessment methods in epidemiologic studies. *Epidemiology and health* 36.
- Siyame E.W., Hurst R., Wawer A.A., Young S.D., Broadley M.R., Chilimba A.D., Ander L.E., Watts M.J., Chilima B., Gondwe J. (2013) A high prevalence of zinc-but not iron-deficiency among women in rural Malawi: a cross-sectional study. *Int J Vitam Nutr Res* 83: 176–87.
- Slimani N., Freisling H., Illner A.K., Huybrechts I. (2015) Methods to Determine Dietary Intake. *Nutrition Research Methodologies*: 48–70.
- Smith L.C., Subandoro A. (2007) *Measuring food security using household expenditure surveys*. International Food Policy Research Institute.
- Stephen A.M., Mak T.N., Fitt E., Nicholson S., Roberts C., Sommerville J. (2013) Innovations in national nutrition surveys. *Proceedings of the Nutrition Society* 72: 77–88.
- Steyn N., Nel J., Nantel G., Kennedy G., Labadarios D. (2006) Food variety and dietary diversity scores in children: are they good indicators of dietary adequacy? *Public Health Nutrition* 9: 644–650. DOI: doi: 10.1079/PHN2005912.
- Stote K.S., Radecki S.V., Moshfegh A.J., Ingwersen L.A., Baer D.J. (2011) The number of 24 h dietary recalls using the US Department of Agriculture’s automated multiple-pass method required to estimate nutrient intake in overweight and obese adults. *Public Health Nutrition* 14: 1736–1742.
- Stumbo P.J. (2013) New technology in dietary assessment: a review of digital methods in improving food record accuracy. *Proceedings of the Nutrition Society* 72: 70–76. DOI: doi: 10.1017/S0029665112002911.
- Subar A.F., Kipnis V., Troiano R.P., Midthune D., Schoeller D.A., Bingham S., Sharbaugh C.O., Trabulsi J., Runswick S., Ballard-Barbash R., Sunshine J., Schatzkin A. (2003) Using Intake Biomarkers to Evaluate the Extent of Dietary Misreporting in a Large Sample of Adults: The OPEN Study. *American Journal of Epidemiology* 158: 1–13. DOI: 10.1093/aje/kwg092.
- Sun M., Fernstrom J.D., Jia W., Hackworth S.A., Yao N., Li Y., Li C., Fernstrom M.H., Sclabassi R.J. (2010) A wearable electronic system for objective dietary assessment. *Journal of the American Dietetic Association* 110: 45–47.
- Sun M., Burke L., Baranowski T., Fernstrom J., Zhang H., Chen H.-C., Bai Y., Li Y., Li C., Yue Y., Li Z., Nie J., Sclabassi R., Mao Z.-H., Jia W. (2015) An Exploratory Study on a Chest-Worn Computer for Evaluation of Diet, Physical Activity and Lifestyle. *Journal of Healthcare Engineering* 6: 1–22. DOI: doi: 10.1260/2040—2295.6.1.1.
- Swindale A., Bilinsky P. (2006) *Household dietary diversity score (HDDS) for measurement of household food access: indicator guide, Version 2*. Food and Nutrition Technical Assistance Project, Academy for Educational Development, Washington DC.
- Thompson F.E., Subar A.F. (2013) Dietary assessment methodology. In A.M. Coulston, C.J. Boushey & M.G. Ferruzzi, eds. *Nutrition in the prevention and treatment of disease*. Third ed. London, Academic Press. (available at: http://appliedresearch.cancer.gov/diet/adi/thompson_subar_dietary_assessment_methodology.pdf Accessed 23 October 2015).
- Timon C.M., Astell A.J., Hwang F., Adlam T.D.,

- Smith T., Maclean L., Spurr D., Forster S.E., Williams E.A. (2015) The validation of a computer-based food record for older adults: the Novel Assessment of Nutrition and Ageing (NANA) method. *British Journal of Nutrition* 113: 654–664.
- Tooze J.A., Krebs-Smith S.M., Troiano R.P., Subar A.F. (2012) The accuracy of the Goldberg method for classifying misreporters of energy intake on a food frequency questionnaire and 24-h recalls: comparison with doubly labelled water. *European Journal of Clinical Nutrition* 66: 569–576.
- Torheim L.E., Barikmo I., Parr C.L., Hatloy A., Ouattara F., Oshaug A. (2003) Validation of food variety as an indicator of *diet quality* assessed with a food frequency questionnaire for Western Mali. *European Journal of Clinical Nutrition* 57: 1283–1291.
- Trabulsi J., Schoeller D.A. (2001) Evaluation of dietary assessment instruments against doubly labelled water, a biomarker of habitual energy intake. *American Journal of Physiology-Endocrinology And Metabolism* 281: E891–E899.
- van Stuijvenberg M.E., Nel J., Schoeman S.E., Lombard C.J., du Plessis L.M., Dhansay M.A. (2015) Low intake of calcium and vitamin D, but not zinc, iron or vitamin A, is associated with stunting in 2- to 5-year-old children. *Nutrition* 31: 841–846.
- Ventura A.K., Loken E., Mitchell D.C., Smiciklas-Wright H., Birch L.L. (2006) Understanding Reporting Bias in the Dietary Recall Data of 11 Year-Old Girls. *Obesity* 14: 1073–1084.
- Vickers A.J. (2005) Parametric versus non-parametric statistics in the analysis of randomized trials with non-normally distributed data. *BMC medical research methodology* 5: 1.
- Wang D.-H., Kogashiwa M., Kira S. (2006) Development of a New Instrument for Evaluating Individuals' Dietary Intakes. *Journal of the American Dietetic Association* 106: 1588–1593.
- Webster-Gandy J., Madden A., Holdsworth M. (2011) *Oxford handbook of nutrition and dietetics*. OUP Oxford.
- Welch J., Dowell S., Johnson C.S. (2007) Feasibility of using a personal digital assistant to self-monitor diet and fluid intake: a pilot study. *Nephrology Nursing Journal* 34: 43.
- Wessells K.R., Singh G.M., Brown K.H. (2012) Estimating the global prevalence of inadequate zinc intake from national food balance sheets: effects of methodological assumptions. *Plos One* 7: e50565.
- WFP. (2008) *Food consumption analysis: Calculation and use of the food consumption score in food security analysis*. Vulnerability Analysis and Mapping. World Food Programme.
- WHO, UNICEF, USAID, UCDAVIS, IFPRI. (2008) *Indicators for Assessing Infant and Young Child Feeding Practices: Part I Definitions*. Geneva: WHO http://apps.who.int/iris/bitstream/10665/43895/1/9789241596664_eng.pdf (retrieved in November, 2015): 3–4.
- WHO/UNICEF. (2010) *Indicators for assessing infant and young child feeding practices. Part 3*, in: W. L. C.-i.-P. Data (Ed.), WHO, Geneva.
- Willett W. (2013a) Correction for the Effects of Measurement Error. In: Willett W. Third ed. *Nutrition Epidemiology*. Oxford University Press. Third ed.
- Willett W. (2013b) Nature of Variation in Diet. In: Willett W. Third ed. *Nutrition Epidemiology*. Oxford University Press. Third ed.
- Willett W., E L. (2013) Reproducibility and Validity of Food-Frequency Questionnaires. In: Willett

- W. Third ed. Nutrition Epidemiology. Oxford University Press. Third ed.
- Willett W.C., Hu F.B. (2007) The food frequency questionnaire. *Cancer Epidemiology Biomarkers & Prevention* 16: 182–183.
- Willett W.C., Manson J.E., Stampfer M.J., Colditz G.A., Rosner B., Speizer F.E., Hennekens C.H. (1995) *Weight, weight change, and coronary heart disease in women: risk within the 'normal' weight range.* *Jama* 273: 461–465.
- Williamson D.A., Allen H.R., Martin P.D., Alfonso A.J., Gerald B., Hunt A. (2003) Comparison of digital photography to weighed and visual estimation of portion sizes. *Journal of the American Dietetic Association* 103: 1139–1145.
- Wong S.S., Boushey C.J., Novotny R., Gustafson D.R. (2008) Evaluation of a computerized food frequency questionnaire to estimate calcium intake of Asian, Hispanic, and non-Hispanic white youth. *Journal of the American Dietetic Association* 108: 539–543.
- Yaroch A.L., Toozé J., Thompson F.E., Blanck H.M., Thompson O.M., Colón-Ramos U., Shaikh A.R., McNutt S., Nebeling L.C. (2012) Evaluation of three short dietary instruments to assess fruit and vegetable intake: The National Cancer Institute's Food Attitudes and Behaviors Survey. *Journal of the Academy of Nutrition and Dietetics* 112: 1570—1577.
- Yon B.A., Johnson R.K., Harvey-Berino J., Gold B.C. (2006) The use of a personal digital assistant for dietary self-monitoring does not improve the validity of self-reports of energy intake. *Journal of the American Dietetic Association* 106: 1256–1259.
- Zhu F., Bosch M., Woo I., Kim S., Boushey C.J., Ebert D.S., Delp E.J. (2010) The Use of Mobile Devices in Aiding Dietary Assessment and Evaluation. *IEEE journal of selected topics in signal processing* 4: 756–766. DOI: 10.1109/jstsp.2010.2051471.
- Zoellner J., Anderson J., Gould S.M. (2005) Comparative validation of a bilingual interactive multimedia dietary assessment tool. *Journal of the American Dietetic Association* 105: 1206–1214.



©FAO/Sebastian Liste

8

APPENDICES

APPENDIX 1: EXAMPLES OF FOOD FREQUENCY QUESTIONNAIRE (FFQ)

INSTRUCTIONS FOR FILLING OUT A QUALITATIVE FFQ

- First, fill out below the personal information (name, ID number and date, etc.).

**Note: If you need to assess only the frequency with which food items or food groups are consumed during a specific time period, you are referring to Table A below which was designed to provide descriptive qualitative information.*

- Mark with an 'X' the frequency with which you consumed the food recorded: never, seldom (less than once a month); number of times per month, week or day.

**Note: you can adapt the questionnaire for local use depending on the food habits of the sample population.*

8.1 EXAMPLE 1A: FILLED-OUT QUALITATIVE FFQ

An FFQ used to evaluate calcium intake in a group of adolescents in Hong Kong* – modified for use by adapting to the Italian food habits.

*(*Lee W.T.K. et al. Generalized Low Bone Mass of Girls with Adolescent Idiopathic Scoliosis is related to Inadequate Calcium Intake and Weight Bearing Physical Activity in Peripubertal Period. Osteoporos Int 2005, 16: 1024–35.)*

Table A - Descriptive qualitative FFQ

FOOD ITEMS	FREQUENCY					
	NEVER	SELDOM (LESS THAN ONCE A MONTH)	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	DAILY
1. Fresh milk						
full-cream		x				
skimmed					x	
high calcium, low fat						
chocolate flavoured						
other flavoured						
others _____						
2. Milk powder						
full-cream						
skimmed						
high calcium low fat						
3. Condensed (sweetened) milk						
reconstituted with water or						
spread on bread						
4. Evaporated milk						
reconstituted with water or						
mixed in beverages						
5. Soymilk						
Plain soymilk				x		

FOOD ITEMS	FREQUENCY					
	NEVER	SELDOM (LESS THAN ONCE A MONTH)	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	DAILY
With calcium				x		
Chocolate flavoured						
Others _____						

6. Cheese

Caciotta		x				
Fontina						
Gorgonzola						
Mozzarella				x		
Parmesan cheese						x
Pecorino			x			
Ricotta			x			
Stracchino				x		

7. Ice-cream, dairy

Fruit flavours		x				
Cream flavours			x			

8. Egg, hen

Boiled egg				x		
Soft-boiled egg		x				
Poached egg		x				
Egg omelette				x		

FOOD ITEMS	FREQUENCY					
	NEVER	SELDOM (LESS THAN ONCE A MONTH)	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	DAILY

9. Rice, raw

Brown rice				x		
White rice		x				

10. Bread

Wholemeal bread						x
White bread						

11. Pasta

Dried pasta (spaghetti, maccheroni, fusilli etc.)						x
Fresh/filled pasta (lasagne, cannelloni, tortellini etc.)				x		

12. Biscuits

Dried biscuits					x	
Cream crackers		x				
Wafers			x			

13. Cake

Cupcake		x				
Sponge cake filling (with cream or chocolate)		x				
other cakes _____						

14. Chocolate

smarties						
----------	--	--	--	--	--	--

FOOD ITEMS	FREQUENCY					
	NEVER	SELDOM (LESS THAN ONCE A MONTH)	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	DAILY
M&M						
Kit-Kat			x			
Mars bar		x				
Toblerone						

15. Legumes, dried, fresh, frozen

white beans		x				
red beans		x				
black eyed peas or cow peas			x			
lentils				x		
chickpeas				x		

16. Green leafy vegetables, cooked

Broccoli		x				
Chicory			x			
Chard				x		
Kale		x				
Spinach					x	
Cauliflower						
Brussels Sprouts						
Dill						

FOOD ITEMS	FREQUENCY					
	NEVER	SELDOM (LESS THAN ONCE A MONTH)	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	DAILY
Beet						

17. Green leafy vegetables, raw

Arugula					x	
Radicchio			x			
Romaine lettuce						x
Watercress		x				
Iceberg lettuce		x				
Endive						
Valerian						

18. Meat, white

Chicken				x		
Turkey			x			
Rabbit						

19. Meat, red

Beef						
Calf		x				
Pork			x			

INSTRUCTIONS FOR FILLING OUT A SEMI-QUANTITATIVE FFQ

- First, fill out below the personal information (name, ID number and date etc.).
- **Note: If you also need to assess the quantity of food intake, the method becomes semi-quantitative and you are referring to the Table B below designed to provide semi-quantitative information.*
- Mark with an 'X' the frequency with which you consumed the food recorded: never, seldom (less than once a month), number of times per month, week or day;
- Record the amount of food consumed under the column 'Amount', using household measurements as reported (cup, teaspoon or slice, etc.) or if you have the weight or volume please specify, and include the units (e.g. grams, oz. millilitre or litre, etc.).
- Write down the brand name of the food when applicable in the final column.

**Note: you can adapt the questionnaire for local use depending on the food habits of the sample population.*

8.2 EXAMPLE 1B: FILLED-OUT SEMI-QUANTITATIVE FFQ

Table B - Semi-quantitative FFQ

FOOD ITEMS	FREQUENCY						AMOUNT	BRAND
	NEVER	SELDOM (LESS THAN ONCE A MONTH)	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	DAILY		
20. Fresh milk								
full-cream		x					1 cup (200ml)	Parmalat
skimmed					x		1 cup (200ml)	Parmalat
high calcium, low fat								
chocolate flavoured								
other flavoured								
others _____								
21. Milk powder								
full-cream								

FOOD ITEMS	FREQUENCY						AMOUNT	BRAND
	NEVER	SELDOM (LESS THAN ONCE A MONTH)	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	DAILY		
skimmed								
high calcium low fat								
22. Condensed (sweetened) milk								
reconstituted with water or								
spread on bread								
23. Evaporated milk								
reconstituted with water or								
mixed in beverages								
24. Soymilk								
Plain soymilk				x			1 cup (200ml)	IsolaBio
With calcium				x			1 cup (200ml)	Alpro
Chocolate flavoured								
Others _____								
25. Cheese								
Caciotta		x					1 slice, thick (50g)	Auricchio

FOOD ITEMS	FREQUENCY						AMOUNT	BRAND
	NEVER	SELDOM (LESS THAN ONCE A MONTH)	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	DAILY		
Fontina								
Gorgonzola								
Mozzarella				x			2 slices, thick (100g)	Galbani
Parmesan cheese						x	1 slice, thick (50g)	Lucini
Pecorino			x				1 slice, thick (50g)	Locatelli
Ricotta			x				2 heaped tablespoons, (80g)	Galbani
Stracchino				x			1 slice, thick (50g)	Granarolo
26. Ice-cream, dairy								
Fruit flavours		x					small cone	
Cream flavours			x				medium cone	
27. Egg, hen								
Boiled egg				x			2 pieces	Ovito
Soft-boiled egg		x					2 pieces	Ovito
Poached egg		x					2 pieces	Ovito
Egg omelette				x			2 pieces	Ovito

FOOD ITEMS	FREQUENCY						AMOUNT	BRAND
	NEVER	SELDOM (LESS THAN ONCE A MONTH)	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	DAILY		
28. Rice, raw								
Brown rice				x			1 cup (80g)	Scotti
White rice		x					1 cup (80g)	Scotti
29. Bread								
Wholemeal bread						x	2 slices, thin (100g)	
White bread								
30. Pasta								
Dried pasta (spaghetti, maccheroni, fusilli etc.)						x	1 cup (80g)	Barilla
Fresh/ filled pasta (lasagne, cannellini, tortellini etc.)				x			1 cup (130g)	De Cecco
31. Biscuits								
Dried biscuits					x		5 pieces (40g)	Gentilini
Cream crackers		x					2 pieces (30g)	Ferrero
Wafers			x				4 pieces (30g)	Locher
32. Cake								
Cupcake		x					1 slice	Jemima

FOOD ITEMS	FREQUENCY						AMOUNT	BRAND
	NEVER	SELDOM (LESS THAN ONCE A MONTH)	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	DAILY		
Sponge cake filling (with cream or chocolate)		x					1 slice	Redman
other cakes								

33. Chocolate

smarties								
M&M								
Kit-Kat			x				½ bar (small, 45g)	Nestlé
Mars bar		x					1 bar (45g)	Mars
Toblerone								

34. Legumes, dried, fresh, frozen

white beans		x					1 cup (50g dried)	Valfrutta
red beans		x					1 cup (50g dried)	Valfrutta
black eyed peas or cow peas			x				1 cup (150g fresh)	Findus
lentils				x			1 cup (50g dried)	Alce Nero
chickpeas				x			1 cup (50g dried)	Alce Nero

FOOD ITEMS	FREQUENCY						AMOUNT	BRAND
	NEVER	SELDOM (LESS THAN ONCE A MONTH)	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	DAILY		

35. Green leafy vegetables, cooked

Broccoli		x					1 cup (200g)	
Chicory			x				1 cup (200g)	
Chard				x			1 cup (200g)	
Kale		x					1 cup (200g)	
Spinach					x		1 cup (200g)	
Cauliflower								
Brussels Sprouts								
Dill								

36. Green leafy vegetables, raw

Arugula					x		1 cup (80 g)	
Radicchio			x				1 cup(80 g)	
Romaine lettuce						x	4 leaves (80g)	
Watercress		x					1 cup (80g)	
Iceberg lettuce		x					1 cup (80g)	
Endive								

FOOD ITEMS	FREQUENCY						AMOUNT	BRAND
	NEVER	SELDOM (LESS THAN ONCE A MONTH)	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	DAILY		

37. Meat, white, raw

Chicken				x			2 slices (200g)	Amadori
Turkey			x				2 slices (200g)	Amadori
Rabbit								

38. Meat, red, raw

Beef								
Calf		x					1 slice (100g)	
Pork			x				2 slices (200g)	

APPENDIX 2: AN EXAMPLE OF BRIEF DIETARY QUESTIONNAIRE

INSTRUCTIONS FOR FILLING OUT A BRIEF DIETARY QUESTIONNAIRE

- First, fill out below the personal information (the name, ID number, group, sex, birth etc.)
- Specify how often you eat the following food items.
- Mark with an 'X' the box for the most appropriate answer for the frequency of the following food group that you consumed.

8.3 EXAMPLE 2: FILLED-OUT BRIEF DIETARY QUESTIONNAIRE

Name: Richard ID Number: 004 Sex: M

Date of Birth: 24th October 1982 Date: 2nd November 2005

Table C - A brief dietary questionnaire

FOOD ITEMS	FREQUENCY							
	NEVER OR RARELY	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	5-6 PER WEEK	DAILY	2-3 PER DAY	5+ PER DAY
Cabbage	x							
Chard			x					
Chicory				x				
Cauliflower		x						
Carrot						x		
Chive		x						
Chinese cabbage	x							
Cress					x			
Beet green		x						
Brussels sprout					x			
Endive				x				
Leaf lettuce						x		

FOOD ITEMS	FREQUENCY							
VEGETABLES	NEVER OR RARELY	1-3 PER MONTH	1-2 PER WEEK	3-4 PER WEEK	5-6 PER WEEK	DAILY	2-3 PER DAY	5+ PER DAY
Leek		x						
Pumpkin		x						
Romaine lettuce						x		
Red pepper				x				
Radicchio					x			
Scallion				x				
Tomato							x	
Turnip	x							
Watercress		x						
Winter squash		x						

APPENDIX 3: AN EXAMPLE OF 24-HOUR RECALL

INSTRUCTIONS FOR FILLING OUT A 24-HOUR RECALL

- First, fill out below the personal information (the name, ID number, sex, birth etc.)
- Report all the foods and beverages consumed in the preceding 24 hours or in the preceding day along with further descriptions of the food where it is requested.
- Start with the first thing eaten in the morning until the last food item consumed before waking up the next morning, reporting for each meal the time along with the place consumed.
- Report for each meal consumed: name of food, food description, household measures (e.g. slices, teaspoons, etc.), unit of measure if possible (e.g. grams, oz. and ml etc.), and finally the kind of preparation methods used and/or ingredients.

8.4 EXAMPLE 3: FILLED-OUT 24-HOUR RECALL*

Name: Simon ID Number: 008 Sex: M

Date of Birth: 13th July 1989 Date: 18th September 2010

Table D - A 24-hour recall

BREAKFAST				Time: 7.15 am Place: home
NAME	FOOD DESCRIPTION	HOUSEHOLD AMOUNT	AMOUNT (g/mL)	PREPARATION/ INGREDIENTS
Soymilk	Soymilk, plain	1 cup	200ml	Hot soymilk
+ coffee	Espresso (Illy)	1 coffee cup	30ml	+ coffee (without sugar)
Bread	Wholemeal bread, with salt	2 slices, thin	100g	Toasted bread
+ butter	Low fat butter, (Granarolo)	1 level tablespoon	10g	
+ jam	Blueberry jam, (Zuegg)	2 heaped teaspoons	20g	Sugar free jam
Biscuits	Dried, Vanilla biscuits	4/5 pieces	30g	Homemade

SNACK (mid-morning)				Time: 10.00 pm Place: work
NAME	FOOD DESCRIPTION	HOUSEHOLD AMOUNT	AMOUNT (g/mL)	PREPARATION/ INGREDIENTS
Apple	Fresh apple	1 medium	150g	Fresh apple with peel
Cappuccino	Full cream milk	1 cup	200ml	Hot cappuccino + sugar
	+ espresso (Illy)	1 coffee cup	30ml	
+ sugar, brown, packed		1 pack	7g	
Crackers	Wholemeal crackers, low fat, (Misura)	1 pack	30g	Wholemeal crackers, low fat, without salt

LUNCH				Time: 13.00 pm Place: work
NAME	FOOD DESCRIPTION	HOUSEHOLD AMOUNT	AMOUNT (g/mL)	PREPARATION/ INGREDIENTS
Dried pasta, (uncooked) + beans, white	Macaroni (Barilla) with beans, white, mature seeds, canned (Valfrutta)	1 cup	60g	Boiled Pasta with pulses (beans)
		1 cup	150g	
		1 tablespoon	10ml	
+ e.v. olive oil				+ e.v. olive oil
+ cheese		2 heaped tablespoons	20g	+ parmesan (cheese)
Salad, raw	Lettuce, romaine, fresh, raw	4 leaves	80g	Fresh salad
+ e.v. olive oil	(De Cecco)	2 tablespoons	20ml	+ e.v. olive oil
+ salt		A pinch		+ salt
Water	Natural water	2 glasses	400ml	

SNACK (mid-afternoon)				Time: 4.00 pm Place: outside
NAME	FOOD DESCRIPTION	HOUSEHOLD AMOUNT	AMOUNT (g/mL)	PREPARATION/ INGREDIENTS
Yogurt	Yogurt, plain Low fat, without sugar (Parmalat)	1 cup	125g	Plain yogurt
+ cereals	Cereals, oats, regular and quick, not fortified, dry (Ecor)	3 heaped tablespoons	30g	+ cereals
+ honey	Acacia honey, (Fior di Loto)	1 heaped teaspoon	8g	+ honey
DINNER				Time: 20.30 pm Place: home
NAME	FOOD DESCRIPTION	HOUSEHOLD AMOUNT	AMOUNT (g/mL)	PREPARATION/ INGREDIENTS
Chicken	Chicken breast, skinless, boneless, raw	2 slices, thin	200g	Roasted chicken
+ e.v. olive oil		1 tablespoon	10ml	Monini
Tubers	Potatoes, uncooked	2 potatoes, small	200g	Roasted potatoes
+ e.v. olive oil	(Monini brand)	1 tablespoon	10ml	
+ salt		A pinch		
Vegetables	Chicory, cooked	1 cup	200g	Sautéed chicory
+ e.v. olive oil	+ e.v. olive oil	1 tablespoon	10ml	
+ salt	+ salt	A pinch		
Water	Natural water	2 glasses	400ml	
Wine	Red wine	½ glass	80ml	
OTHERS				Time: 23.30 pm Place: home
NAME	FOOD DESCRIPTION	HOUSEHOLD AMOUNT	AMOUNT (g/mL)	PREPARATION/ INGREDIENTS
Dried nuts	Almonds	3 pieces	3g	
	Walnuts	3 pieces	12g	

*Note: record adapted using Italian food habits

APPENDIX 4: AN EXAMPLE OF DIETARY HISTORY

INSTRUCTIONS FOR FILLING OUT A DIETARY HISTORY

A dietary history is used to assess a person's usual food intake over a long period of time. The original dietary history consisted of three parts (see below for an example):

- An in-depth interview to assess usual food intake and eating patterns
- A food list
- A three-day record with portion size estimates used as cross-checks
- First, fill out the personal information below (the ID number, group, sex, birth etc.)
- Write in the first box, all the foods and beverages you consumed and in second box the relative amount;

**Note: if the food record is estimated, please use household measurements (cups, slices etc.) if the food record is weighed please use a scale (grams)*

- Specify in the third box the preparation method (boiled, fried, cooked etc.) and if possible please write down the brand name of the food.
- Finally, under the 'frequency' column, mark with an 'X' the box for the most appropriate answer, relating to the frequency with which food items are consumed (e.g. during the last week).

**Note: you can adapt the questionnaire depending on the food habits of the sample population*

8.5 EXAMPLE 4: FILLED-OUT DIETARY HISTORY*

ID Number: 004 (Group: Chan 001) Sex: M Date of Birth: 13th July 1989

Date: 18th September 2010 Day of Week: Tuesday

Table E - A dietary history record

BREAKFAST				FREQUENCY					
				Time: 8.30 am Place: home					
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5 - 6 per wk	Daily
Skimmed milk, without sugar	1 cup (200ml)	Hot skimmed milk	Parmalat					x	
Wholemeal bread, no salt	2 slices, thin (100g)	Toasted bread							

BREAKFAST				Time: 8.30 am Place: home		FREQUENCY					
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
+ jam without sugar	2 heaped teaspoons, (20g)	Apricot jam	Zuegg								
Yogurt, plain	1 cup(125g)	Yogurt, plain, low fat, no sugar	Yomo			x					
+ cereals, corn flakes, plain	6–8 heaped tablespoons, (30 g)	Cereals, ready-to-eat, corn grits, toasted plain	Carrefour								
Sandwich, white bread, commercially prepared	1 sandwich, medium (80 g)	Toasted bread			x						
+ omelette	2 eggs	Cooked egg	Ovito								
+ e.v. olive oil	1 tablespoon, (10ml)		De Cecco								
+ lettuce, romaine, fresh, raw	4 leaves (80g)	Lettuce, fresh, raw									
Biscuits, dried	6–8 pieces (50g)	Dried biscuits with chocolate chips, commercially prepared	Mulino Bianco			x					
+ orange juice	1 glass (200ml)		Santal								
Cappuccino without sugar	1 cup (150ml)	Hot Skimmed Milk	Parlmalat		x						
	+ coffee cup (30ml)	+ coffee	+ espresso (kimbo)								

BREAKFAST				Time: 8.30 am Place: home		FREQUENCY					
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
+ wholemeal croissant filled with honey	1 piece (70g)										
Fruit salad, fresh	2 cups (200g)				x						
+ yogurt, plain, low fat, sugar free	1 cup (125g)		Sterzing-Vipiteno								
English tea	1 cup (150 ml)	Hot tea	Twinings			x					
+ yogurt cake, cinnamon flavoured	2 slices, thin (200g)	Yogurt cake (homemade), prepared from recipe, fat free									

MID-MORNING TEA				Time: 10.30 am Place: school		FREQUENCY					
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
Banana	1 medium (150g)	fresh banana	Chiquita			x					
Apple	1 medium (150g)	Fresh apple	Melinda				x				
Cappuccino without sugar	1 cup (150ml) + coffee cup (30ml)	Hot skimmed milk + coffee Espresso				x					
+ biscuits, buttermilk	3/4 pieces (30g)	Biscuits buttermilk, commercially prepared	Mulino Bianco								

MID-MORNING TEA				Time: 10.30 am Place: school		FREQUENCY					
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
Orange	1 medium (150g)	Riberella				x					
+ rice crackers, Gluten free	1 pack (30g)	Galbusera									
Porridge oats	3 heaped tablespoons (30g)	Rapunzel				x					
+ soymilk, plain	1 cup (150ml)	Isola Bio									

LUNCH				Time: 13.00 pm Place: school		FREQUENCY					
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
Rice, white, long-grain (basmati), raw	1 cup (100g)	Boiled rice	Scotti			x					
+ mushrooms, fresh	1 cup (80g)	+ sliced, sautéed mushrooms									
+ cheese: parmesan cheese	1 heaped tablespoon, (10g)		Bonduelle								
+ e.v. olive oil	1 tablespoon, (10ml)		Monini								

LUNCH				Time: 13.00 pm		Place: school		FREQUENCY					
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily				
+ salt	A pinch												
VEGETABLES													
Lettuce, romaine, fresh, raw	4 leaves (80g)	Raw lettuce								x			
+ e.v. olive oil	1 tablespoon, (10ml)		Monini										
+ salt	A pinch												
Apple Cake	1 slice (100g)	Baked cake	Homemade		x								
DRIED PASTA													
Spaghetti, rigatoni, fusilli	1 cup (100g) (uncooked)	Boiled pasta	De Cecco				x						
FRESH/FILLED PASTA													
Lasagne	1 piece (250g)	Baked pasta	Barilla		x								
Cannelloni (filled with red meat- beef- and cheese- ricotta)	2 pieces (200g)	Baked pasta	Barilla		x								
Tortellini (with spinach and cheese)	1 cup (125g)	Boiled pasta	Rana			x							
OTHERS													
Mashed, dehydrated, potatoes	A pack, commercially prepared (75g)	mashed, dehydrated potatoes, prepared from recipe, with whole milk + water + cheese (cooked in pan)	Buitoni			x							

LUNCH				Time: 13.00 pm Place: school		FREQUENCY					
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
OTHERS											
+ full cream milk	1 heaped cup (300ml)		Parmalat								
+ cheese (parmesan cheese) +salt	3 heaped tablespoons (30g). A pinch		Lucini								
Soup, pulses and wheat, dried	1 cup (100g)	Boiled lentils, chickpeas, spelt, oats	AlceNero			x					
+ e.v. olive oil	1 tablespoon (10ml)	De Cecco									
+ salt	A pinch										
Cheese, mozzarella, whole milk	1 piece (125g)	Fresh Mozzarella	Vallelata		x						
+ pork, cured ham	3 oz. slices (45g)	+ cured ham, pork	Beretta								
+ tomatoes, red, fresh	3 pieces (200g)	+ fresh tomatoes	Cirio								
+ e.v. olive oil	1 tablespoon, (10ml)		De Cecco								
+ salt	A pinch										
VEGETABLES											
Zucchini, with skin, fresh	2-3 pieces (200g)	Sliced, grilled					x				
Eggplant, raw	1 piece (250g)	Sliced, grilled			x						

LUNCH				Time: 13.00 pm		FREQUENCY					
				Place: school							
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
VEGETABLES											
Mushrooms, white, fresh, raw	1 cup (80g)	Sliced, cooked			x						
Pepper, red, fresh, (uncooked)	1 piece (200g)	Sliced, grilled			x						
+ e.v. olive oil	1 tablespoon, (10ml)		Monini								x
+ salt	A pinch										x
Water	2 glasses (400ml)	Natural water									x

AFTERNOON TEA				Time: 4.00 pm		FREQUENCY					
				Place: home							
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
English Tea	1 cup (200ml)	Hot Tea	Sainsbury's brand				x				
+ sugar, brown, unpacked	1 teaspoon sugar (5g)										
+ dried, chocolate biscuits	2 pieces (20g)	Dried, chocolate, biscuits, commercially prepared	Mulino Bianco								

AFTERNOON TEA				Time: 4.00 pm Place: home		FREQUENCY					
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
Pear juice	1 medium glass (200ml)	Fresh pear juice	Yoga			x					
+ yogurt, plain, low fat	1 cup (125g)	+ yogurt, plain, low fat, sugar free	Danone, Vitasnella								
Banana	1 medium (150g)	Fresh fruit, raw	Chiquita			x					
+ cappuccino	1 cup (150ml) + coffee cup (30 ml)	Whole milk + coffee, espresso, (Illy)	Illy								
Apple	1 medium (150g)	Fresh fruit, apple, raw, with skin				x					
+ multigrain crackers, (with rice, buckwheat and spelt)	1 pack (30g)	Misura									
Fruit salad, fresh	1 big cup (200g)	Fresh mixed fruits (orange, apple, banana, kiwi)			x						
DINNER				Time: 8.30 pm Place: outside/home		FREQUENCY					
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
FISH											
Sogliola, fillet, fresh (uncooked)	1 piece (150g)	Baked				x					

DINNER				Time: 8.30 pm		FREQUENCY					
				Place: outside/home							
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
FISH											
Codfish, frozen, fillet (uncooked)	1 piece (150g)	Baked			x						
Salmon, pink, fillet, fresh (uncooked)	1 piece (150g)	Cooked in the pan				x					
+e.v. olive oil	1 tablespoon (10ml)								x		
+ salt	A pinch								x		
+ lemon	A few drops of lemon juice								x		
MEAT											
Chicken, breast, fillet, boneless, skinless (uncooked)	2 slices, thin (200g)	Grilled chicken	Amadori				x				
Chicken, legs, with bone without skin (uncooked)	2 pieces, small (200g)	Grilled chicken	Aia		x						
Turkey, breast, fillet, without skin (uncooked)	2 slices, thin (200g)	Baked turkey	Amadori		x						
Beef, steak (uncooked)	1 slice (130g)	Grilled Beef			x						
Beef, meatballs, (cooked)	2 pieces (120g)				x						
+salt	A pinch								x		

DINNER				Time: 8.30 pm		FREQUENCY					
				Place: outside/home							
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
TUBERS											
Potatoes, peeled	1 cup (200g)	French Fries			x						
VEGETABLES, COOKED											
Spinach, fresh	1 cup (200g)	boiled				x					
Chicory, fresh	1 cup (200g)	sautéed				x					
Chard, fresh	1 cup (200g)	boiled					x				
+ e.v. olive oil	1 tablespoon, (10 ml)		Monini								x
+ salt	A pinch										x
OTHERS											
Eggs, whole, raw	2 eggs, hen (100g)	Boiled egg	Ovito			x					
Bread, oatmeal	2 slices, thin (100g)	Fresh bread						x			
Cheese cake	1 slice, (100g)		Homemade		x						
Water	2 glasses (400ml)	Natural water	Lete								x
Alcoholic beverage, wine, red, Pinot Noir	½ glass (75ml)	Red wine	Antinori		x						

OTHERS				Time: 11.30 pm Place: outside/home		FREQUENCY					
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
Dark chocolate	2 pieces, (square) thin (20g)		Lindt		x						
Muffin	1 piece (80g)	Yogurt Muffin, commercially prepared	OrGran		x						
DRIED NUTS											
walnuts	3 pieces (12g)						x				
almonds	5 pieces (5g)						x				
pistachios	10 pieces(10g)					x					
BISCUITS											
dried biscuits, cinnamon flavoured	4/5 pieces (30g)	Commercially prepared	Gentilini				x				
Chocolate wafer	2 pieces (20g)	Commercially prepared	Loacher			x					
Coca-cola	1 can (330 ml)		Coca-cola		x						
FRESH FRUITS											
Apple, fresh, raw, with skin	1 medium (150g)					x					
Orange, fresh, raw	1 medium (150g)					x					
Pineapple, raw	2 slices, thick, (150g)				x						

OTHERS				Time: 11.30 pm Place: outside/home		FREQUENCY					
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS	1 per mo	2 - 3 per mo	1 - 2 per wk	3 - 4 per wk	5- 6 per wk	Daily		
ICE-CREAM											
Fruit flavours	Small cone (60g)	Commercially prepared	Algida		x						
Cream flavours	Small cone (60g)	Commercially prepared	Algida		x						

*Note: record adapted using Italian food habits

APPENDIX 5: AN EXAMPLE OF FOOD RECORD

INSTRUCTIONS FOR FILLING OUT A THREE DAY ESTIMATED FOOD RECORD

- First, fill out below the personal information (the ID number, group, sex, birth etc.)
- Write in the first box, all the kind of foods and beverages that you have consumed and in second box the relative amount;

** Note: if the food record is estimated, please use household measurements (cups, slices etc.) if the food record is weighed please use a scale (grams). Specify in the third box the preparation method (boiled, fried, cooked etc.) and if possible please write down the brand name of the food.*

- Clearly state the time and place for each meal.

8.6 EXAMPLE 5: FILLED-OUT THREE DAY FOOD RECORD

An example of food record adapted from 'Lee W.T.K. *et al.* 'Prevalence of obesity and its associated risk factors among Chinese students and adult family members attending Chinese Association of Working Chinese School, Woking, Surrey, U.K. (** Note: questionnaire adapted using a Chinese population's food habits living in the UK.*)

ID Number: 001 (Group: Chan 001) Sex: M Date of Birth: 4th Nov 1990
Date: 30th October 2007 Day of week: Tuesday

Table F - A three day food record

BREAKFAST			Time: 07.30 am Place: home
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS
English tea	1 cup (200ml)	1 teaspoon sugar + skimmed milk	
Toast with butter	2 slices		Kingsmill brand
Siu mai	2 pieces		
Yogurt	1 cup (200g)		Muller brand
MID-MORNING TEA			Time: 10.00 am Place: school
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS
Orange juice Banana	1 cup (medium)		Tropicana brand

LUNCH			Time: 12.30 pm Place: school
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS
Sandwich made from:			
- thick white bread	2 slices		Kingsmill brand
- ham	2 slices		
- cheese	1 slice		
- lettuce	3 leaves		
- butter	trace		(olive spread)
Apple	1 (medium)		
Spongy cake	1 (30g)		Cadbury brand
Orange squash	1 glass (large)		Robinson's brand
AFTERNOON TEA			Time: 3.30 am Place: home
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS
English tea	1 cup	1 teaspoon sugar + skimmed milk 1 teaspoon	Tropicana brand
Rich tea biscuits	2 pieces		Sainsbury's brand
DINNER			Time: 7.30 pm Place: outside
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS
Rice	1 small bowl	Boiled	
Chicken wings	2 wings	Cooked	In oyster sauce
Spare-ribs	2 ribs	Oven-baked	In barbeque sauce
Pak-choi	1 portion (100g)	Boiled	

DINNER			Time: 7.30 pm Place: outside
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS
Chinese flowering cabbage fried with beef	(40g)	Fried with black pepper	
Strawberry cheese cake	1 portion (100g)		Tesco brand
Orange juice	1 glass (medium)		Tropicana brand
Water	1 glass		
OTHERS			
FOOD/DRINK	AMOUNT	PREPARATION METHOD	REMARKS
Chocolate	2 fun-size packages		Twix brand
Crisps	1 package (32g)		Walkers brand
Supplement: Complete Multimineral-Multivitamin Formula	1 tablet		Centrum brand

APPENDIX 6: AN EXAMPLE OF MINIMUM DIETARY DIVERSITY – WOMEN (MDD-W) QUESTIONNAIRE

INSTRUCTIONS FOR FILLING OUT A MDD-W QUESTIONNAIRE

- Fill in the MDD-W Code and demographic information, making sure all items are filled in correctly.
- Ask the respondent a series of questions about the foods and drinks that she consumed the previous day or the last 24 hours. The 24-hour cycle lasted from the time she got up in the morning yesterday until the time she got up in the morning today. (Be cautious that some women might eat or drink at night, e.g. lactating mother’s breastfeed infants at night). Both inside and outside home food consumption are counted.
- If the respondent is the one who is responsible for cooking in the family, remind her that you are going to ask about the diet that she actually consumed but not the diets of her family.

8.7 EXAMPLE 6: FILLED-OUT MDD-W QUESTIONNAIRE

MDD-W questionnaire used for data collection in the Sughd region, Tajikistan.

Example from ‘Lazarte *et al.* ‘Integrating the Minimum Dietary Diversity-Women (MDD-W) module into the Household Budget Survey (HBS) in Tajikistan’, 2015.

This project was carried out by the Food and Agriculture Organization of the United Nations (FAO) and the Agency of Statistics under the President of the Republic of Tajikistan (AoS), with financial contributions from the European Union and FAO in the framework of the project “*Improved Global Governance for Hunger Reduction*”.

For more information please visit the Nutrition Assessment team of the FAO Nutrition Division: www.fao.org/nutrition/assessment

STATISTICS AGENCY UNDER THE PRESIDENT OF TAJIKISTAN

Household Budget Survey

Household No (from HBS):

--	--	--	--	--	--	--

Minimum Dietary Diversity-Women

MDD-W Code:

--	--	--	--	--	--	--

Name of the enumerator:	Sophia
Date of interview: (dd/mm/yyyy)	23/05/2013

PART 1 – QUALITATIVE 24-HOUR DIETARY RECALL

- Please describe the foods (meal/ tea break/ break) that you ate or drank yesterday, whether at home or outside the home. Start with the first food or drink of the morning.
- Write down all foods and drinks mentioned. When composite dishes are mentioned, ask for the FULL list of ingredients.
- When the respondent has finished, probe for meal/ tea break/ break not mentioned.

Table G - A 24-hour recall

Time: 7.30 am Place: home	Meal/ Tea break/ Break: Breakfast
Hot Skimmed milk (Parlmalat) + chocolate, dried biscuits (Gentilini); Coffee (espresso, illy) + sugar	
Time: 10.30 am Place: work	Meal/ Tea break/ Break: Break
Yogurt, plain, low fat, sugar free (Sterzing-vipiteno) + orange juice (Santal) Crackers, Wheat, wholemeal, no salt (Misura)	
Time: 13.00 pm	Meal/ Tea break/ Break: Lunch
Dried pasta (Barilla) with tomato sauce + cheese (parmesan) + e.v. olive oil Spinach, fresh, boiled + e.v. olive oil + salt Potatoes, without peel, baked + e.v. olive oil + salt Natural water	
Time: 16.00 pm Place: work	Meal/ Tea break/ Break: Break
English tea (Twinings) + apple, with skin, fresh (Melinda) + Chocolate cake, prepared from recipe without frosting (homemade)	
Time: 20.30 pm Place: home	Meal/ Tea break/ Break: Dinner
Turkey, breast, baked (Amadori) + e.v. olive oil + salt + drops of lemon juice	
Time: 20.30 pm Place: home	Meal/ Tea break/ Break: Dinner
carrots and potatoes, boiled + e.v. olive oil + salt Lettuce, Romaine, raw, fresh + e.v. olive oil + balsamic vinegar + salt Natural and fizzy water	
Time: 23.00 pm Place: home	Meal/ Tea break/ Break: Break
Ice cream (cream flavours), commercially prepared + 2 walnuts	

PART 2 – TABLE OF FOOD GROUPS

- After the respondent recalls all the food and beverages consumed, underline the corresponding foods in the list under the appropriate food group. For any food groups not mentioned, ask the respondent if a food item from this group was consumed. Write '1' in the column next to the food group if at least one food in this group has been underlined.
- Write "0" in the column next to the food group at this stage if no food items have been underlined in the food group.
- Write down the name of the food in the "Remarks" section (found at the end of the questionnaire) that are mentioned but not found in the food groups.
- Check to make sure all ingredients used in mixed dishes have been identified from the person who prepared the food. Alternatively if the food was bought from the market, go and check the ingredients from the shop in the market.

**Note: this table is an example specifically adapted for use in Tajikistan. For the MDD-W model questionnaire, please refer to the "Minimum Dietary Diversity for women - A Guide to Measurement" (FAO, 2016).*

Table H - Table of food groups

QUESTION NUMBER	FOOD GROUP	LOCALLY AVAILABLE FOODS	YES=1
			NO=0
1	Cereals	Wheat, barley [perlofca], buckwheat, maize, rice, pasta, wheat bread, other bread, wheat flour, other flour, pasta products, kirieshki (snack made from flour), pop corn	1
2	White roots and tubers	Potato, turnip (yellow, red), radish	1
3	Vitamin A rich vegetables and tubers	Pumpkin, carrot, red sweet pepper (bulgori), squash	1
4	Dark green leafy vegetables (DGLV)	Spinach, rhubarb, siyoalaf, bargi salat; dill, coriander, mint, parsley, blue basilica, green garlic, green onion, sorrel, wine leaves, jagh-jagh [Consider as DGLV when consumed at least one tablespoon of these vegetable(s) per day. Otherwise, go to Group 16]	1
5	Other vegetables	Cabbage, cauliflower, garlic, cucumber, leek, tomato, onion, eggplant, beetroot, mushrooms fresh and dried, anzur, green beans, green pepper	1
6	Vitamin A rich fruits	Apricot and dried apricot, peach and dried peach, persimmon, cantaloupe melon	0
7	Other fruits	Apple, banana, lemon, watermelon, mandarin, grapes, pears, melon, muskmelon, fruits and berries, dried fruits and berries, raisins, oranges, cherries, figs, plum, pomegranate, prune, quince, raspberries, strawberries, blackberries, mulberries, king mulberries, yellow cherry, plum (orange-colour), sinjid, chelon, dulona (haw), kiwi, pineapple, grapefruit, simorodina, hips	1

QUESTION NUMBER	FOOD GROUP	LOCALLY AVAILABLE FOODS	YES=1
			NO=0
8	Flesh foods and organ meat	Beef, mutton, goat, chukar, rabbit, chicken, goose, turkey, quail, sausages, veal, lamb and chevron, meat of wild animals and games, sausage products and smoked meat, horse, duck, ox tail, liver, kidney, heart, lung,, stomach, intestine, tongue, brain (goat and sheep), spleen [ssiyohlavak] (cow, goat and sheep),	1
9	Eggs	Quail eggs, chicken eggs, goose eggs, turkey eggs, duck eggs	0
10	Fish and sea foods	Fresh and frozen fish, canned fish, smoked fish, dried fish, caviar, crab sticks	0
11	Beans and peas	Mung bean, peas, red beans, white beans, lentils, chickpeas, split peas	0
12	Nuts and seeds	Sesame seed, pistachios, almonds, pumpkin seeds, sunflower seeds, walnuts, peanuts, apricot seeds, hazelnut, pecan	0
13	Milk and milk products	Milk, skim milk, sour milk, yogurt, qurut (dried yogurt), ice-cream, kefir, chaka, kaymak (sour cream), cheese, sour cream/smetana, powdered milk, condensed milk, tvorog, falla (colostrum from cow), goat milk	1
14	Oils and fats	Vegetable oil (sunflower, flax, sesame, cotton, olive), butter, sheep fat, margarine, mayonnaise, ravgani zard (oil prepared from kaimak and butter), mahsar/saflo oil (plant oil), mixed oil (animal fat and vegetable oil), turta (sediment of boiled kaymak), potato chips, cow fat, sheep fat	1
15	Sweets	Sugar, honey, candies, chocolate, cakes, biscuits, jam, halva, baklava, obinabot (crystallised sugar), nishollo, shirinii tut (Tajik snicker, mulberry paste with sugar), pechak	1
16	Spices, condiments, beverages	Black pepper, cumin, ketchup, salt, pripava (adviya), chicken/ beef cubes; balsamic vinegar, vinegar Dill, coriander, mint, parsley, blue basilica, green garlic, green onion, sorrel, rosemary (small green leaves), black sesame seeds, bay leaf. [Consider as condiments when these vegetable(s) are consumed less than one tablespoon a day. Otherwise, go to Group 4]; Coffee, black and green tea, khuch, fruit drinks (compote), boiled water, sweet soda, water	1


REMARKS

PART 3 – TABLE TO DERIVE THE MMD-W SCORE

Aggregation of food groups to calculate the MDD-W score

Table I - The 10 food groups

FOOD/DRINK	AMOUNT	PREPARATION METHOD
1,2	All starchy staples	
11	Beans and peas	
12	Nuts and seeds	
13	All dairy	
8, 10	Flesh foods (including organ meat and miscellaneous small protein)	
9	Eggs	
4	Vitamin A-rich dark green leafy vegetables	
3, 6	Other vitamin A-rich vegetables and fruits	
5	Other vegetables	
7	Other fruits	
MDD-W score (sum of the above scores)		



European
Commission

This publication has been produced with the assistance of the European Union. The contents of this publication are the sole responsibility of FAO and can in no way be taken to reflect the views of the European Union.

Nutrition and Food Systems Division
Economic and Social Development Department
Food and Agriculture Organization of the United Nations
via delle Terme di Caracalla
00153 Rome, Italy
www.fao.org/nutrition/assessment

ISBN 978-92-5-130635-2



9 789251 306352

19940EN/1/06.18